

An Inventory of Irrigation Structures in the Upper Clark Fork River Drainage, Montana



Final Report 2011
USFWS Agreement Number- 601818J270
Will Schreck, Ryan Kreiner, Brad Liermann, and Jason Lindstrom
Montana Fish, Wildlife and Parks

Cover photograph: Headgate of an irrigation ditch on Cottonwood Creek located near Deer Lodge, Montana.

Table of Contents

| | |
|---|----|
| Introduction | 5 |
| Methods | 7 |
| <i>Irrigation Structures</i> | 7 |
| <i>Fish Sampling</i> | 7 |
| <i>Discharge and Temperature Monitoring</i> | 8 |
| <i>Radio Telemetry</i> | 9 |
| Inventory Results | 10 |
| <i>Ranch Creek</i> | 10 |
| <i>Stony Creek</i> | 11 |
| <i>Beaver Creek</i> | 12 |
| <i>Rock Creek</i> | 13 |
| <i>West Fork Rock Creek</i> | 15 |
| <i>Ross Fork Rock Creek</i> | 16 |
| <i>Middle Fork Rock Creek</i> | 18 |
| <i>East Fork Rock Creek</i> | 19 |
| <i>Harvey Creek</i> | 26 |
| <i>South Fork Lower Willow Creek</i> | 27 |
| <i>Gird Creek</i> | 28 |
| <i>Boulder Creek</i> | 29 |
| <i>Gold Creek</i> | 30 |
| <i>Clark Fork River</i> | 34 |
| <i>Cottonwood Creek</i> | 35 |
| <i>Baggs Creek</i> | 39 |

| | |
|--|-----|
| <i>Racetrack Creek</i> | 41 |
| <i>Lost Creek</i> | 44 |
| <i>Warm Springs Creek</i> | 45 |
| <i>Storm Lake Creek</i> | 47 |
| <i>Mill Creek</i> | 49 |
| <i>Browns Gulch</i> | 50 |
| Radio Telemetry Results | 51 |
| Entrainment Summary and Site Prioritization | 53 |
| Mitigation Efforts | 59 |
| Literature Cited | 60 |
| Appendix A | 61 |
| Appendix B | 107 |
| Appendix C | 140 |
| Appendix D | 142 |

Introduction

Agriculture is a prevailing economic presence in the western United States. In Montana alone, the market value of agriculture is nearly \$2 billion per year (Christensen 2005). Given the arid nature of Montana summers, it is often necessary to divert water from rivers and streams to support agriculture. The operation of irrigation structures such as ditches can have dramatic effects on the health of these systems by entraining resident fishes and de-watering the stream channel. In western Montana, native fishes such as westslope cutthroat trout *Oncorhynchus clarkii lewisi* and the federally threatened bull trout *Salvelinus confluentus* are likely at high risk for entrainment into irrigation ditches due to their migratory life histories. These species are especially at risk as post-spawn adults and juveniles out-migrating from spawning or rearing tributaries (Bahn 2007). Also, due to the thermal sensitivity of bull trout and westslope cutthroat trout, elevated stream temperatures accompanying channel de-watering are likely to impact these species.

In 2009, Montana Fish, Wildlife & Parks (FWP) received a grant from the U.S. Fish and Wildlife Service (USFWS) on behalf of the Fisheries Restoration and Irrigation Mitigation Act (FRIMA) to survey the Upper Clark Fork River Basin's irrigation systems. The objectives of this study were to:

1. Inventory diversions on prioritized tributaries in the Upper Clark Fork River Basin to aid in the development of fish passage/screening plans.
2. Provide recommendations for site-specific passage and entrainment solution options for prioritized streams.
3. Design, plan, and implement, as appropriate, identified fish passage and screening projects as funds become available via the accepted FRIMA proposal and ranking process and/or from other funding sources.

The area chosen for this study includes the Clark Fork River and its tributaries from the headwaters of Silver Bow Creek downstream to its confluence with Rock Creek near Clinton, MT (Figure 1). This includes major agricultural regions such as the Deer Lodge and Flint Creek valleys, as well as the Rock Creek valley, which is less impacted by agriculture and irrigation, but contains significant populations of native fish species such as bull trout and westslope cutthroat trout. Irrigation structures inventoried for this study were chosen based on their presumed impact on the overall Clark Fork River fishery, as well as their suspected entrainment of native fishes.

This report incorporates data collected intensively in 2009, 2010 and 2011, as well as past sampling efforts from 2007 and 2008. This report includes a prioritization list of critical diversions resulting from electrofishing surveys from 2007-2011.




Figure 1. Survey locations within the Upper Clark Fork River Basin by year.

Methods

Irrigation Structures

Brief descriptions of the diversion and headgate styles were recorded to provide information useful for future planning and mitigation. During this inventory, common diversion styles included gravel and boulder berms or pin and plank structures, and common headgate styles included metal slide gates (Waterman C-10 and R-5) and wooden gates with stopboards (See MDNRC 2001). Waterman C-10 headgates are prefabricated metal structures, adjustable with screw-down gates, and R-5 headgates are similar in construction, but are unable to seal, therefore allowing leakage past the gate when closed.

During 2010 and 2011, Montana Trout Unlimited (TU) measured physical characteristics of most diversions visited by FWP in the Upper Clark Fork River Basin. The TU inventory focused on stream habitat, stream morphology and fish passage, and provided dimension and velocity measurements at most diversions (TU 2011).

Fish Sampling

Electrofishing was used to obtain species composition, length frequency, and catch-per-unit-effort (CPUE) of the fishes entrained in irrigation ditches. At many sites, water clarity was low and it was most effective to electrofish in an upstream direction. In locations where avoidance was a concern, block nets were placed at the upstream end of the section or the headgate functioned as a barrier and no block net was installed. When avoidance was less of a concern or access was limited, no block net was installed. Single-pass electrofishing runs were made through 100 m sections of the ditches, unless length was restricted by de-watering or access. Sections were often located immediately below headgates, but additional sections downstream were electrofished on certain ditches to further document entrainment. During the inventory, these additional downstream surveys indicated fish densities are greatly reduced in sections further from the headgates. Thus, fish densities documented immediately below headgates rarely represented densities throughout the ditches.

In ditches expected to contain high numbers of entrained fish, multiple-pass depletions were conducted to generate fish population estimates with confidence intervals. In these situations, block-nets were used, multiple passes were made through sections, and captured fish were removed and placed aside in live-cars. Population estimates were calculated using the maximum likelihood estimator provided by Montana Fish, Wildlife & Parks' FA+ fisheries analysis software.

Fishes were collected with a Smith-Root LR-24 Electrofisher. Fishes were identified to species, measured, weighed and released. Data were recorded on standard FWP electrofishing data sheets using FWP species abbreviations. These abbreviations are also used in the tables presented in the results section. Following are the abbreviations and the species they represent.

BULL = bull trout
EB = eastern brook trout *Salvelinus fontinalis* (referred to as brook trout)
LC SU = largescale sucker *Catostomus macrocheilus*
LL = brown trout *Salmo trutta*
LN SU = longnose sucker *Catostomus catostomus*
LN DC = longnose dace *Rhinichthys cataractae*
MWF = mountain whitefish *Prosopium williamsoni*
ONC = unidentifiable *Oncorhynchus* species
RB = rainbow trout *Oncorhynchus mykiss*
RS SH = redbside shiner *Richardsonius balteatu*
SL COT = slimy sculpin *Cottus cognatus*
SU = unspecified sucker
WCT = westslope cutthroat trout

Data Summary

Fishery data was summarized by species and included the number of fish captured catch-per-unit-effort (CPUE) or depletion estimates (standardized to number of fish per 100 m), mean and range of fish lengths, and percent species composition. Tables summarizing this information have been included for each ditch surveyed. Additionally, length-frequency histograms were constructed for trout species with ≥ 5 individuals collected to illustrate length distributions. These data are provided as an appendix (Appendix A). For the purpose of this report, ditches were named for the approximate river mile (RM) location of the diversion and headgate. River mile designations were calculated using FWP's River Mile Index GIS Layer. Photographs of most irrigation structures visited are also provided as an appendix (Appendix B). Fish population data for streams discussed in this report were taken from Lindstrom et al. 2008, Liermann et al. 2009, Lindstrom 2011, or unpublished electrofishing data collected in 2009.

In most ditches, not all slimy sculpin or other non-salmonid species (e.g., longnose sucker, largescale sucker and longnose dace) were captured or enumerated. Thus, these species were categorized as abundant, common or rare. Species with ≥ 51 individuals were considered abundant, 11-50 individuals considered common, and 1-10 individuals considered rare. The presence of an "A" in the tables of this report signifies the species was abundant, "C" signifies common, and "R" signifies rare.

The Entrainment Summary and Site Prioritization section of this report includes a summary of the ditches surveyed and priority ratings (Table 27). These ratings are based upon the species and abundance of fishes entrained in the ditch, with emphasis on native species. Ditches were assigned high, medium or low ratings.

Discharge and Temperature Monitoring

At most survey locations, stream discharges were measured using a Marsh-McBirney flow meter and a top-set wading rod. When feasible, discharge was measured within the ditch and the adjacent stream. Mean velocities were obtained over a 20 sec period using

the fixed point average (FPA) setting and discharge was calculated as cubic feet per second (cfs). During the summers of 2010 and 2011, a Montana Trout Unlimited (TU) crew collected many of the measurements in Powell and Deer Lodge counties. These estimates were summarized in a report submitted to the Watershed Restoration Coalition of the Upper Clark Fork (WRC), and several were included in this report (TU 2011).

Since the withdrawal of water for irrigation may impact downstream water temperatures, stream temperatures were monitored in a few of the streams chosen for this inventory. In 2007 and 2009, one or more thermographs were deployed in target drainages (ONSET Computer Corp, Model: HOBO Water Temp Pro V2). To most accurately determine the effects of de-watering on stream temperature, thermographs were deployed above and below major diversions in some streams. In other streams, only one thermograph was deployed, generally near the mouth of the stream, below all diversions. Temperatures were measured in degrees Celsius (°C) every half hour, with temperatures above 15 °C considered harmful to bull trout and westslope cutthroat trout (Behnke 2002). Results are displayed in an appendix (Appendix C).

Radio Telemetry

In April 2009, a radio telemetry project was initiated within the Upper Clark Fork River. This study consisted of implanting trout captured from the mainstem Clark Fork with radio tags and observing their movements throughout the system (Mayfield and McMahon 2010). Fish species selected for tagging in this study were roughly proportional to overall species composition in the Clark Fork River. In 2009, 100 trout comprised of 72 brown trout, 15 westslope cutthroat trout, 6 suspected westslope cutthroat/rainbow trout hybrids, 6 rainbow trout, and 1 bull trout were implanted with radio tags. In 2010, 149 trout were tagged consisting of 104 brown trout, 33 westslope cutthroat trout, 6 cutthroat/rainbow trout hybrids, 3 bull trout and 3 rainbow trout. In 2011, an additional 19 trout were tagged consisting of 9 brown trout, 8 westslope cutthroat trout, and 2 cutthroat/rainbow trout hybrids. Three of the objectives for this study were to identify 1) critical spawning and rearing habitat, 2) possible impediments to both up- and downstream fish migration, and 3) possible limiting factors and causes of mortality to trout in the Upper Clark Fork River. Any results from the telemetry study pertinent to this project's objectives are displayed in the Radio Telemetry Results section of this report. Re-location data were collected for the study until December 2011.

Inventory Results

Ranch Creek

Ranch Creek is a tributary to Rock Creek and enters the drainage at approximately RM 11.5 (Figure 1). The entire upper portion of the creek is located on lands administered by the Lolo National Forest. Most of the lower portion runs through private land with the exception of the U.S. Forest Service (USFS) campground located near the confluence of Grizzly Creek. Land use on the lower creek is limited to a few private residences and small horse pastures. Through past electrofishing surveys, native westslope cutthroat trout and bull trout are known to inhabit Ranch Creek. Migratory bull trout from Rock Creek are also known to spawn in Ranch Creek.

Two small (<2 CFS) ditches draw water from Ranch Creek on its lower end, below the Grizzly Creek campground (Figure 1). Electrofishing surveys were completed in these ditches in August 2009 and 2010. The upper ditch, located at RM 1.1, is fitted with a metal slide gate and water is diverted into this ditch by a gravel berm (Appendix B). The diversion does not serve as a barrier to fish passage. Two 100 m sections on upper ditch (RM 1.1) were surveyed, one near the headgate and one approximately one mile downstream at Norton USFS campground. In 2009, a total of 11 fish were captured below the headgate of the upper ditch (Table 1, Appendix A). Eight of these fish were westslope cutthroat trout, along with two brook trout and one brown trout. Slimy sculpin were also present but rare in this section. No fish were captured in the lower section of this ditch near the Norton campground. In 2010, a total of 26 fish were collected in 100 m below the headgate of the upper ditch, comprised of fourteen westslope cutthroat trout, four brown trout, four unidentifiable *Oncorhynchus* spp, three brook trout and one rainbow trout. In 2010, slimy sculpin were abundant in this section. Again in 2010, no fish were captured at the lower section on this ditch (Norton campground). The lower of the two ditches, located at RM 0.1, has no formal headgate, but water enters this ditch from a side channel on Ranch Creek. On both occasions the lower ditch (RM 0.1) was surveyed below its point of diversion near the Norton campground. In 2009, five westslope cutthroat trout were collected along with two brook trout (Table 1, Appendix A). Slimy sculpin were also abundant in the section. In 2010, ten brown trout, eight brook trout, three *Oncorhynchus* species, and one rainbow trout were collected (Table 1, Appendix A). Slimy sculpin were again abundant.

Although no bull trout were observed in these ditches in 2009 or 2010, past research found a bull trout entrained in the lower ditch (RM 0.1) on lower Ranch Creek. Approximately ten years ago, a radio-tagged adult bull trout was entrained during a radio telemetry study and presumably died in the ditch (radio tagged fish made movements in the ditch but the radio transmitter was later recovered in the ditch- Brad Liermann, Montana Fish, Wildlife and Parks, personal communication).

Table 1. Electrofishing data collected during 2009 and 2010 in ditches on Ranch Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 1.1 (Headgate) | 2009 | WCT | 8 | 8 | 121 | 78-187 | 73 |
| | | LL | 1 | 1 | 96 | 96 | 9 |
| | | EB | 2 | 2 | 83 | 50-115 | 18 |
| | 2010 | WCT | 14 | 14 | 95 | 35-161 | 54 |
| | | LL | 4 | 4 | 86 | 46-135 | 15 |
| | | RB | 1 | 1 | 77 | 77 | 4 |
| | | ONC | 4 | 4 | 80 | 71-92 | 15 |
| | | EB | 3 | 3 | 132 | 115-163 | 12 |
| | | SL COT | A | - | - | - | - |
| | 2009 | NO | FISH | - | - | - | - |
| | | NO | FISH | - | - | - | - |
| RM 0.1 (Headgate) | 2009 | WCT | 5 | 5 | 91 | 78-112 | 71 |
| | | EB | 2 | 2 | 163 | 133-193 | 29 |
| | | SL COT | R | - | - | - | - |
| | 2010 | LL | 10 | 10 | 138 | 78-180 | 45 |
| | | RB | 1 | 1 | 130 | 130 | 5 |
| | | ONC | 3 | 3 | 58 | 32-109 | 14 |
| | | EB | 8 | 8 | 157 | 124-235 | 36 |
| | | SL COT | A | - | - | - | - |

On August 10, 2010, discharges were measured within both ditches and upstream of the upper ditch at RM 1.1 in mainstem Ranch Creek. The ditch at RM 1.1 was receiving 1.2 cfs, approximately 4% of the mainstem discharge at 28.6 cfs. The ditch at RM 0.1 near the mouth of Ranch Creek was receiving 0.8 cfs, approximately 3% of the remaining mainstem discharge (estimated at 27.4 cfs).

Stony Creek

Stony Creek is a tributary to Rock Creek and enters the drainage at approximately RM 38.4 (Figure 1). Roughly one mile of lower Stony Creek flows through private land, while upper Stony Creek flows through National Forest land. Forest Road 241 (FR 241) follows the lower creek for nearly five miles, while the remainder of the drainage is roadless. Fish populations in upper Stony Creek are comprised entirely of native westslope cutthroat trout and bull trout, while a mix of native and non-native species are present in the lower portion of the drainage. Past electrofishing surveys found non-native brown trout and rainbow trout in lower Stony Creek, with visual evidence of hybridization between westslope cutthroat trout and rainbow trout (Liermann et al. 2008).

Fluvial bull trout from Rock Creek are also known to spawn in Stony Creek. In June 2009, a radio-tagged westslope cutthroat trout from the Clark Fork River was relocated at the mouth of Stony Creek and may have spawned in the creek.

One irrigation ditch exists on Stony Creek (Figure 1). The headgate of this ditch is located just above the USFS Stony Creek campground at RM 0.3 in the lower portion of the drainage. This ditch is fitted with a metal slide gate and water is diverted by a gravel berm (Appendix B). The diversion does not likely serve as a barrier to fish passage. In August 2009, a three-pass depletion was conducted from the headgate to a block-net 100 m downstream. A total of 53 westslope cutthroat trout and 53 brown trout were collected in this ditch (Table 2, Appendix A). The population estimate for westslope cutthroat trout was 55.0 fish per 100 m (53.0-59.3; 95% confidence interval (CI)) and for brown trout, 54.0 fish per 100 m (53.0-57.4; 95% CI).

Table 2. Electrofishing data collected during 2009 in a ditch on Stony Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (Depletion) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|----------------------------|------------------|-------------------|------------------|
| RM 0.3 (Headgate) | 2009 | WCT | 53 | 55.0 | 73 | 27-139 | 50 |
| | | LL | 53 | 54.0 | 52 | 45-60 | 50 |

Discharges within the ditch and upstream of the diversion on Stony Creek were measured on August 27, 2009. On this date, the ditch was receiving 0.8 cfs, approximately 7% of the mainstem discharge of 12.5 cfs.

Beaver Creek

Beaver Creek is a tributary to Upper Willow Creek and enters the drainage at approximately RM 12.8 (Figure 1). Beaver Creek begins on the Beaverhead-Deerlodge National Forest, flows through approximately 1.5 miles of state administered land, and crosses a short section of private land near its confluence with Upper Willow Creek. The creek is accessible by gated FR 4325, which provides vehicle access for administrative purposes, as well as walk-in traffic for recreation. Based on electrofishing surveys conducted in 2009, fish populations in Beaver Creek are comprised primarily of westslope cutthroat trout and brook trout; however, one brown trout was captured.

One irrigation ditch was observed drawing water from Beaver Creek at approximately RM 0.9 (Figure 1). This ditch is fitted with a metal slide gate and water is diverted from Beaver Creek by a check dam spanning the channel (Appendix B). The check dam may serve as a barrier to fish passage during low flows. This ditch was electrofished for 100 m below the headgate in 2009 (Table 3). Both brook trout and westslope cutthroat trout were found entrained in this ditch. In total, 16 westslope cutthroat trout and 23 brook trout were collected.

Table 3. Electrofishing data collected during 2009 in the ditch on Beaver Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 0.9 (Headgate) | 2009 | WCT | 16 | 16 | 79 | 58-117 | 41 |
| | | EB | 23 | 23 | 58 | 46-120 | 59 |

Rock Creek

Rock Creek is a tributary to the Clark Fork River and enters the drainage at approximately RM 239.0 (Figure 1). Four forks join to form mainstem Rock Creek, including the West, Ross, Middle, and East Forks. The upper reaches of these forks are located on land administered by the Beaverhead-Deerlodge National Forest, while the lower reaches are located on private land. The upper portion of mainstem Rock Creek runs through private land, and the lower portion crosses a combination of Lolo National Forest and private land. Land use within the upper Rock Creek watershed is primarily agriculture, with historical mining present. Land use lower in the watershed is primarily recreation (National Forest) and private residences. Rock Creek provides a world renowned fishery and a valued recreational resource supported primarily by brown trout, but rainbow trout, and native westslope cutthroat trout and bull trout are also present.

In 2010 and 2011, electrofishing surveys were completed in a total of three irrigation ditches on mainstem Rock Creek (Figure 1). In August 2010, ditches located at RM 50.6 and 50.5 (just below the confluence of the West and Middle Forks at RM 51.9) on Rock Creek were surveyed. These ditches are both fitted with metal slide gates and rock weirs, and supply water to grazing land on a guest ranch and adjacent downstream landowners (Appendix B). Neither diversion is likely to be a barrier to fish passage. In 2010, two sections were sampled on the upper ditch at RM 50.6 with the first being located immediately below the headgate and the second 0.5 mi downstream. One bull trout, two westslope cutthroat, and five brown trout were collected in the first 100 m below the headgate (Table 4). Mountain whitefish were also present but rare in this section. No fish were collected in the section 0.5 mi downstream of the headgate. In 2010, three sections were sampled on the lower of the two ditches at RM 50.5 with the first being located immediately below the headgate, the second 0.5 mi downstream and the lowest 1.6 mi downstream. Three westslope cutthroat trout and 14 brown trout were collected in the first 100 m below the headgate (Table 4, Appendix A). Mountain whitefish, longnose dace and slimy sculpin were also present but rare in this section. No fish were collected 0.5 mi downstream of the headgate, likely due to very limited holding habitat in the reach. Two species were captured in the section 1.6 mi downstream including, mountain whitefish and longnose dace (Table 4). In late August 2011, the ditch at RM 50.5 was re-surveyed in two locations to further document the composition of species entrained. A total of six westslope cutthroat trout, nine brown trout and one brook trout were collected in the first 100 m below the headgate (Table 4, Appendix A). Mountain whitefish, longnose dace and slimy sculpin were also collected below the headgate. One brown trout was collected in a 100 m section 0.5 mi downstream of the headgate.

In 2011, an additional ditch was surveyed on upper Rock Creek downstream from the two previously visited. The ditch, located at RM 49.5, is fitted with wooden gate and receives water diverted from a side-channel by a pin and plank diversion. The potential for the diversion to be a barrier is unknown since the planks were removed at the time of the survey. Two sections were electrofished on the lowest ditch, one immediately below the headgate and another 0.8 mi downstream. In a 100 m section below the headgate, one westslope cutthroat, nine brown trout, as well as several mountain whitefish, longnose dace and slimy sculpin were collected (Table 4). Two cutthroat trout, four brown trout and numerous mountain whitefish were collected 0.8 mi downstream of the headgate in a 170 m section. Catch-per-unit-effort estimates were calculated with a 170 m section. The trout collected in the lower section were large adult cutthroat (235-355mm) and brown trout (235-351mm), and represent losses to the recreational fishery.

Table 4. Electrofishing data collected during 2010 and 2011 in ditches on Rock Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 50.6 (Headgate) | 2010 | BULL | 1 | 1 | 197 | 197 | 13 |
| | | WCT | 2 | 2 | 198 | 121-275 | 25 |
| | | LL | 5 | 5 | 100 | 51-135 | 62 |
| | | MWF | R | - | - | - | - |
| RM 50.6 (0.5 mi) | 2010 | NO | FISH | - | - | - | - |
| RM 50.5 (Headgate) | 2010 | WCT | 3 | 3 | 115 | 91-113 | 18 |
| | | LL | 14 | 14 | 88 | 47-279 | 82 |
| | | MWF | R | - | - | - | - |
| | | LN DC | R | - | - | - | - |
| | | SL COT | R | - | - | - | - |
| RM 50.5 (0.5 mi) | 2010 | NO | FISH | - | - | - | - |
| RM 50.5 (1.6 mi) | 2010 | MWF | R | - | - | - | - |
| | | LN DC | R | - | - | - | - |
| RM 50.5 (Headgate) | 2011 | WCT | 6 | 6 | 89 | 39-112 | 38 |
| | | LL | 9 | 9 | 87 | 60-174 | 56 |
| | | EB | 1 | 1 | 47 | 47 | 6 |
| | | MWF | R | - | - | - | - |
| | | LN DC | R | - | - | - | - |
| | | SL COT | R | - | - | - | - |
| RM 50.5 (0.5 mi) | 2011 | LL | 1 | 1 | 135 | 135 | 100 |

Table 4 (continued). Electrofishing data collected during 2010 and 2011 in ditches on Rock Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|----------------------------|-----------------------------|---------------------|----------------------|---------------------|
| RM 49.5 (Headgate) | 2011 | WCT | 1 | 1 | 115 | 115 | 10 |
| | | LL | 9 | 9 | 73 | 47-204 | 90 |
| | | MWF | C | - | - | - | - |
| | | LN DC | R | - | - | - | - |
| | | SL COT | R | - | - | - | - |
| RM 49.5 (0.8 mi) | 2011 | WCT | 2 | 1.2* | 295 | 235-355 | 33 |
| | | LL | 4 | 2.4* | 314 | 235-351 | 67 |
| | | MWF | C | - | - | - | - |

Discharge measurements collected just below the headgates on August 19, 2010, established the ditch at RM 50.6 was receiving 5.8 cfs and the ditch at RM 50.5 was receiving 18.1 cfs.

West Fork Rock Creek

West Fork Rock Creek is a tributary to Rock Creek, and joins the Middle Fork Rock Creek at approximately RM 51.3 to form mainstem Rock Creek (Figure 1). West Fork Rock Creek originates in the Sapphire Mountains on the Beaverhead-Deerlodge National Forest and the majority of the upper drainage is located within National Forest lands. Land ownership in the lower drainage is a mix of USFS, State of Montana, and private lands. Grazing occurs on private lands, as well as National Forest and State of Montana land, through grazing allotments. Historical logging is also evident in portions of the drainage. For most of its length, West Fork Rock Creek is followed by Montana Highway 38 (Skalkaho Highway). Fish populations in upper West Fork Rock Creek are comprised primarily of native bull trout, westslope cutthroat trout, and mountain whitefish, while lower in the drainage brown trout as well as native longnose sucker and longnose dace become more abundant (Liermann et al. 2009).

In August of 2010, a ditch on lower West Fork Rock Creek was surveyed (Figure 1). This ditch is located at approximately RM 4.2, and has been used to flood irrigate a large private pasture approximately one mile downstream. The ditch is fitted with a wooden gate, regulated by wooden planks and a gravel berm diverts water into this ditch (Appendix B). The diversion does not serve as a barrier to fish passage. This ditch was highly inefficient at the time of the survey, with several breaks in the walls of the ditch. On August 25 the ditch held very little water and a total of 38 fishes were collected in a 30 m section below the headgate (Table 5, Appendix A). All fish collected were sub-adult and consisted of 11 westslope cutthroat trout, one brown trout and 26 unidentifiable *Oncorhynchus* species. Slimy sculpin were present but rare in this ditch.

Table 5. Electrofishing data collected during 2010 in a ditch on the West Fork Rock Creek. (*- CPUE extrapolated from a 30 m section).

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 4.2 (Headgate) | 2010 | WCT | 11 | 37* | 81 | 73-95 | 29 |
| | | LL | 1 | 3* | 52 | 52 | 3 |
| | | ONC | 26 | 87* | 36 | 27-48 | 68 |
| | | SL COT | R | - | - | - | - |

Although discharge was not measured in 2010, visual observations suggested the ditch was receiving less than 0.5 cfs, likely due to the low discharge of the West Fork as well as the disrepair of the diversion and headgate.

Ross Fork Rock Creek

Ross Fork Rock Creek is a tributary to West Fork Rock Creek and enters the drainage just above the confluence of the West and Middle Forks of Rock Creek at approximately RM 0.2 (Figure 1). The Ross Fork originates on National Forest land and the upper portion of the drainage is managed as roadless. Roads are present in the middle portion of the drainage on National Forest land, where historical logging is evident. Lower Ross Fork Rock Creek flows through several cattle ranches and additional grazing occurs on National Forest land through leased grazing allotments. Lower in the drainage water is diverted from the Ross Fork to flood irrigate hay pastures. In the upper reaches of Ross Fork Rock Creek, fish populations are comprised almost entirely of native westslope cutthroat trout and bull trout, with migratory bull trout from Rock Creek known to spawn in the upper Ross Fork. In the lower reaches, non-native brown trout and brook trout comprise a higher proportion of the fish community, and de-watering accompanied by elevated stream temperatures is common, presumably due to irrigation withdrawals (Liermann et al. 2009).

At least seven diversions are known to draw water from mainstem Ross Fork Rock Creek. These diversions are primarily operated to flood irrigate pastures in the lower portion of the Ross Fork valley. Several of these ditches are fitted with metal slide gates and most draw water via gravel/boulder berms (Appendix B). The upstream most diversion was located on National Forest land at approximately RM 9.4 (Figure 1). This ditch delivered water several miles downstream to a private ranch, but in 2007 appeared to be highly inefficient. Although this ditch received a substantial amount of water at the diversion, it was nearly dry at its intersection with Angelico Creek, several miles upstream of the intended water users. Five more diversions were located downstream between RM 6.5 and 5.5. These diversions were located in close succession and diverted similar volumes of water. Although no measurements were collected, the mainstem appeared to be dewatered by approximately 50% of the expected average summer level below this series of diversions.

The lowest diversion on the Ross Fork, was located below RM 3.5 and when observed was diverting the majority of the water remaining in the Ross Fork. Visual estimates suggested the discharge in the mainstem below this site was less than 10% of average summer levels. This ditch was not electrofished due to a lack of landowner permission. Visual observation of the stream flows below these diversions in 2007, suggests these withdrawals greatly impacted in-stream flows.

Two thermographs were used to monitor temperatures on the Ross Fork in 2007, one above the majority of the diversions at RM 6.7, and one near the mouth of a small tributary at RM 0.1. At the upper site, stream temperature exceeded 20°C on four days with a maximum-recorded temperature of 20.7°C. At the lower site, temperatures exceeded 20°C on 24 days with a maximum recorded of 24.4°C. These measurements suggest water temperatures are influenced by irrigation withdrawals within this 6.5-mile stretch. These elevated temperatures likely negatively impact native bull trout and westslope cutthroat trout.

Electrofishing surveys were completed in six of the ditches on the Ross Fork in 2007. The upper ditch (RM 9.4) was sampled approximately 0.5 miles upstream of the crossing of FR 5060 over Ross Fork Rock Creek. In a 100 m section, four westslope cutthroat trout, one brown trout, and one brook trout were collected (Table 6). Longnose dace and longnose sucker were also present, but rare. Surveys of the five ditches between RM 6.5 and RM 5.5 measured 100 m in length and began within 50 meters of their headgates. Entrainment of westslope cutthroat trout was documented in all ditches and bull trout entrainment was observed in two (Table 6, Appendix A). Other salmonids collected in these ditches included brown trout, brook trout and mountain whitefish. Native longnose sucker and longnose dace were collected in the ditch located at RM 6.0, and slimy sculpin were present in the ditches located at RM's 6.5, 6.0, and 5.5.

Table 6. Electrofishing data collected during 2007 in ditches on Ross Fork Rock Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 9.4 (Headgate) | 2007 | WCT | 4 | 4 | 98 | 30-211 | 27 |
| | | LL | 1 | 1 | 138 | 138 | 7 |
| | | EB | 1 | 1 | 173 | 173 | 7 |
| | | LN DC | R | - | - | - | - |
| | | LN SU | R | - | - | - | - |
| RM 6.5 (Headgate) | 2007 | WCT | 24 | 24 | 44 | 37-56 | 73 |
| | | LL | 9 | 9 | 67 | 52-78 | 27 |
| | | SL COT | C | - | - | - | - |
| RM 6.2 (Headgate) | 2007 | WCT | 22 | 22 | 86 | 32-189 | 42 |
| | | LL | 31 | 31 | 63 | 50-72 | 58 |

Table 6 (continued). Electrofishing data collected during 2007 in ditches on Ross Fork Rock Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 6.1 (Headgate) | 2007 | BULL | 1 | 1 | 43 | 43 | 50 |
| | | WCT | 1 | 1 | 237 | 237 | 50 |
| RM 6.0 (Headgate) | 2007 | WCT | 16 | 16 | 54 | 41-146 | 49 |
| | | LL | 17 | 17 | 64 | 50-150 | 51 |
| | | MWF | R | - | - | - | - |
| | | LN DC | R | - | - | - | - |
| | | LN SU | R | - | - | - | - |
| | | SL COT | R | - | - | - | - |
| RM 5.5 (Headgate) | 2007 | BULL | 1 | 1 | 207 | 207 | 6 |
| | | WCT | 7 | 7 | 105 | 41-180 | 41 |
| | | LL | 8 | 8 | 64 | 55-71 | 47 |
| | | EB | 1 | 1 | 64 | 64 | 6 |
| | | MWF | R | - | - | - | - |
| | | SL COT | A | - | - | - | - |

Middle Fork Rock Creek

Middle Fork Rock Creek and West Fork Rock Creek join to form mainstem Rock Creek just upstream of Skalkaho Bridge at approximately RM 51.4 (Figure 1). The Middle Fork Rock Creek drainage begins in the Anaconda-Pintler Wilderness Area and fish habitat in the upper portion of the drainage is excellent. Land ownership below the wilderness boundary remains USFS; however, additional land uses include cattle grazing and historical logging. In the lower portion of the drainage, land ownership is private and land use is restricted primarily to cattle ranching. In upper Middle Fork Rock Creek, fish populations are comprised of native westslope cutthroat trout and bull trout (Liermann et al. 2009). Large migratory bull trout have also been collected in the upper reaches and are known to spawn in the mainstem Middle Fork and several of its tributaries. Lower in the drainage non-native brook and brown trout comprise more of the community, but westslope cutthroat trout and bull trout remain present. Native mountain whitefish and longnose dace are also found in the lower reaches of the Middle Fork.

In early August 2010, electrofishing surveys were completed in two 100 m sections on a ditch diverting water from Middle Fork Rock Creek at RM 0.1 (Figure 1). The headgate of this ditch is located just upstream of Skalkaho Bridge and the confluence of West and Middle Fork Rock Creek. This ditch is fitted with a metal slide gate and a small gravel berm diverts water from a side-channel of Middle Fork Rock Creek (Appendix B). The diversion does not serve as a barrier to fish passage. This ditch is routed under Montana State Highway 38 and travels through several culverts before supplying water to center pivots on a guest ranch downstream. The first section electrofished began approximately

0.2 mi below the headgate. Nine trout were collected in the 100 m section, consisting of two bull trout, two westslope cutthroat trout and five brown trout (Table 7, Appendix A). Mountain whitefish were also present in the section, but rare. The second section electrofished was located downstream of a culvert, approximately 0.5 mi below the headgate. Only three trout were collected in this section, one westslope cutthroat trout and two brown trout, but native mountain whitefish and longnose sucker were also present.

Table 7. Electrofishing data collected during 2010 in a ditch on Middle Fork Rock Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|-----------------|---------|-------------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 0.1 (0.2 mi) | 2010 | BULL | 2 | 2 | 221 | 187-255 | 22 |
| | | WCT | 2 | 2 | 140 | 134-146 | 22 |
| | | LL | 5 | 5 | 122 | 64-168 | 56 |
| | | MWF | R | - | - | - | - |
| RM 0.1 (0.5 mi) | 2010 | WCT | 1 | 1 | 125 | 125 | 33 |
| | | LL | 2 | 2 | 166 | 154-177 | 67 |
| | | MWF | R | - | - | - | - |
| | | LN SU | R | - | - | - | - |

Discharge measurements collected on August 19, 2010, established the ditch at RM 0.1 was receiving 3.7 cfs. This ditch draws from a medium-sized side channel of the Middle Fork and at the time of measurement, flows in the side channel were low and debris had accumulated at the entrance to the headgate.

East Fork Rock Creek

East Fork Rock Creek is a tributary to Middle Fork Rock Creek and enters the drainage at approximately RM 2.1 (Figure 1). East Fork Rock Creek begins in the Anaconda-Pintler Wilderness Area and is fed by several tributaries before leaving the wilderness and entering East Fork Reservoir at approximately RM 10.6. The primary purpose of East Fork Reservoir is to store spring runoff for irrigation in the Flint Creek valley via a trans-basin diversion (Flint Creek Main Canal) and the reservoir is managed by Montana Department of Natural Resources and Conservation (DNRC). Releases from the East Fork dam are quite high during the irrigation season (greater than 100 cfs during May-October) and low during the remainder of the year (< 1 cfs during November through April). The diversion into the canal is located below the dam on East Fork Rock Creek at RM 8.6. During the irrigation season, a majority of the East Fork Rock Creek is diverted into this canal. The diverted water travels 2.7 mi, where it is siphoned from the west side of the valley to the east side and eventually gains the necessary elevation to enter the Trout Creek drainage. This water is then delivered to the Flint Creek valley via Trout Creek.

Fish populations in East Fork Rock Creek vary above and below the dam. The reservoir and East Fork Rock Creek above the reservoir are strongholds for native bull trout. Westslope cutthroat trout are currently stocked into the reservoir by FWP and are the most abundant species in the reservoir. Native bull trout are also relatively abundant in East Fork Reservoir while non-native brook and rainbow trout are also present in limited numbers. Conversely, below the dam, fish populations are dominated by brook trout and brown trout with a limited number of bull trout and westslope cutthroat trout. Larger bull trout and westslope cutthroat trout are often observed in the spillway below the dam and it is likely these fish have been entrained through the dam.

Electrofishing surveys have been completed in the Flint Creek Main Canal on several occasions between 2007 and 2011 (Table 8, Appendix A). Sampling was routinely conducted in October after the headgate was closed, due to the high volume and velocity of water in the canal during irrigation season. Each year, a section was electrofished directly below the headgate and varied from 50-100 m in length. Since large numbers of age-0 trout were often captured in this section, length data was not always collected on all fish. In 2007 and 2009, only a sub-set of the fish collected were measured, while in 2008 and 2010, all fish were measured. In 2011, the canal was dry 50 m below the headgate and thus the section electrofished was reduced to 50 m. Also, fish were identified and enumerated, but not measure to total length in 2011.

A single bull trout was captured during 2007 and 2009 in the canal, while three were collected in 2010 and two were collected in 2011 (Table 8). Numerous westslope cutthroat trout have been collected at every visit. A majority of the native bull trout and westslope cutthroat trout captured in the canal are believed to have out-migrated from the reservoir due to their low abundance in East Fork Rock Creek below the dam. Non-native brook, brown and rainbow trout have also been collected in the canal (Table 8). In 2009 and 2011, a section approximately one mile below the headgate near the Lost Sapphire subdivision was surveyed. Two small westslope cutthroat trout were collected from this section in 2009 and two small brook trout were collected in 2011. In 2007, a pool at the siphon release valve was also electrofished. Fish are stranded in this pool after the siphon is shut down for the winter. Both native bull trout and cutthroat trout were collected from this pool along with brook trout, brown trout, and rainbow trout. Fish collected from the canal were released into mainstem East Fork Rock Creek.

Table 8. Electrofishing data collected from 2007 to 2011 in sections of the Flint Creek Main Canal (arranged upstream to downstream). (*- Indicates average or range based on the measurement of a sub-set of fish).

| Section Name | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|--------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| Headgate | 2007 | BULL | 1 | 1.6 | 90 | 90 | <1 |
| | | WCT | 108 | 177 | n/a | 50-70* | 66 |
| | | RB | 19 | 31 | n/a | 75-90* | 12 |
| | | EB | 35 | 57 | n/a | 75-200* | 21 |

Table 8 (continued). Electrofishing data collected from 2007 to 2011 in sections of the Flint Creek Main Canal (arranged upstream to downstream). (*- Indicates average or range based on the measurement of a sub-set of fish).

| Section Name | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|---------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| Headgate | 2008 | WCT | 22 | 22 | 121 | 41-384 | 11 |
| | | LL | 6 | 6 | 312 | 287-351 | 3 |
| | | RB | 4 | 4 | 230 | 152-318 | 1 |
| | | EB | 176 | 176 | 143 | 55-322 | 85 |
| Headgate | 2009 | BULL | 1 | 1 | 460 | 460 | <1 |
| | | WCT | 13 | 13 | 197 | 106-435 | 5 |
| | | LL | 5 | 5 | 334* | 307-375* | 2 |
| | | RB | 4 | 4 | 201 | 156-261 | 2 |
| | | ONC | 171 | 171 | 49* | 43-60* | 71 |
| | | EB | 46 | 46 | 136* | 62-292* | 19 |
| Headgate | 2010 | BULL | 3 | 3 | 110 | 91-121 | 1 |
| | | WCT | 32 | 32 | 127 | 62-178 | 9 |
| | | LL | 40 | 40 | 119 | 62-348 | 12 |
| | | RB | 3 | 3 | 201 | 135-316 | 1 |
| | | ONC | 49 | 49 | 50 | 40-61 | 14 |
| | | EB | 213 | 213 | 119 | 62-279 | 63 |
| | | SL COT | A | - | - | - | - |
| Headgate | 2011 | BULL | 2 | n/a | n/a | 200-480 | 2 |
| | | WCT | 5 | n/a | n/a | n/a | 4 |
| | | LL | 16 | n/a | n/a | n/a | 13 |
| | | EB | 105 | n/a | n/a | n/a | 82 |
| | | SL COT | C | - | - | - | - |
| Lost Sapphire | 2009 | WCT | 2 | 2 | 55* | 50-60* | 100 |
| Lost Sapphire | 2011 | EB | 2 | 2 | 69 | 51-87 | 100 |
| Siphon | 2007 | BULL | 1 | n/a | 90 | 90 | 4 |
| | | WCT | 6 | n/a | n/a | 60-400* | 32 |
| | | LL | 3 | n/a | n/a | 225-325* | 16 |
| | | RB | 2 | n/a | 450* | 450* | 11 |
| | | EB | 7 | n/a | n/a | 55-325* | 37 |

Recent releases from East Fork Dam however, are generally greater than 100 cfs during the irrigation season and commonly less than 1 cfs during the off-season. In August 2008

and 2009, discharges measured in Flint Creek Main Canal just below the headgate, were 104.6 cfs, and 90.0 cfs, respectively. In 2010 discharges were measured in both the canal and East Fork Rock Creek below the diversion. On August 25, 2010 an estimated 115.9 cfs was being diverted into Flint Creek Main Canal, while East Fork Rock Creek received 4.1 cfs. A minimum instream flow of 5 cfs during irrigation season was provided by the Flint Creek Water Users as mitigation for replacing the East Fork Siphon in 2008. On August 25, DNRC was notified measurements collected in the East Fork indicated instream flows were below this agreed amount and the diversion was adjusted to meet the flow requirement. In 2011, discharges were measured on several occasions in East Fork Rock Creek and once within the canal. On August 17, 2011 an estimated 104.1 cfs was being diverted into Flint Creek Main Canal, and East Fork Rock Creek received 8.2 cfs. On September 12 and October 18, 2011 East Fork Rock Creek was receiving 4.8 cfs and 2.6 cfs, respectively. In October 2007, the discharge of East Fork Rock Creek above the reservoir, just downstream of a spring-fed tributary, was 9.7 cfs. This estimate may approximate the historical discharge on the East Fork Rock Creek near the headgate of the Flint Creek Main Canal during a relatively normal snowpack year (this assumes minimal up-welling in the reservoir reach which is unknown).

Recent discussion between canal owners/operators (DNRC/Flint Creek Water Users) and fishery biologists may result in actions to mitigate the impacts of East Fork Reservoir operations on fishes in mainstem East Fork Rock Creek below the dam and populations within East Fork Reservoir. The absence of spring scouring flows and the lack of adequate instream flows appears to be altering fish habitat and affecting species composition, as well as abundance of salmonids. Also, the entrainment of fish into the canal represents direct losses to fish populations in the East Fork below the reservoir. To mitigate these issues, additional instream flows and a spring flushing flow may be provided to the mainstem and a fish screen installed at the headgate of the canal. In 2009-2011, additional sites were sampled on East Fork Rock Creek and in 2010 and 2011, on Ross Fork Rock Creek, to obtain baseline data before these actions are potentially implemented. The sites on Ross Fork Rock Creek will serve as references or control sites to account for regional changes in fish populations. This pre-data will help in assessing the effectiveness and benefits of additional flows and fish screening, if these mitigation actions are taken. On East Fork Rock Creek, multiple-pass depletion estimates were conducted at three sites downstream of the diversion; 1) on a section near the USFS campground approximately 0.2 mi below the diversion, 2) on a private ranch approximately 3.9 mi below the diversion, and 3) on a section on state owned land approximately 7.4 mi below the diversion. On Ross Fork Rock Creek, depletion estimates were conducted on two sites, both located on U.S. Forest Service land, at approximately RM 8.9 and RM 7.0. The results from electrofishing the East Fork and the Ross Fork are summarized in Table 9 below, with sample sites arranged from upstream to downstream.

Table 9. Electrofishing results from East Fork Rock Creek and Ross Fork Rock Creek. Population estimates of fish only ≥ 75 mm. Corral Gulch section (2007) located just upstream of the State Section (2010 and 2011). (* - indicates section name changed from Christiansen Ranch to Angelico Creek).

| Water Body | Section Name | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (Depletion) | 95% CI Range | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|------------|---------------|--------------|---------|-------------------------|----------------------------|--------------|------------------|-------------------|------------------|
| East Fork | Below dam | 2009 | LL | 15 | 14.0 | 13.0-18.2 | 135 | 71-255 | 3 |
| | | | EB | 511 | 473.0 | 462.9-483.1 | 115 | 56-244 | 97 |
| East Fork | Below dam | 2010 | WCT | 13 | 13.0 | 13.0-14.2 | 118 | 97-136 | 2 |
| | | | LL | 30 | 24.0 | 20.0-35.4 | 116 | 65-256 | 3 |
| | | | EB | 862 | 710.0 | 659.9-760.1 | 95 | 50-223 | 95 |
| East Fork | Below dam | 2011 | WCT | 9 | 9.0 | 9.0-9.5 | 140 | 95-200 | 1 |
| | | | LL | 36 | 36.0 | 36.0-37.8 | 131 | 63-190 | 6 |
| | | | EB | 599 | 607.0 | 600.1-613.9 | 120 | 63-1408 | 93 |
| East Fork | Cadieux Ranch | 2007 | BULL | 1 | 1.0 | n/a | 173 | n/a | <1 |
| | | | WCT | 2 | 2.0 | 2.0-2.8 | 292 | 259-324 | 1 |
| | | | LL | 96 | 97.0 | 96.0-100.1 | 175 | 61-337 | 66 |
| | | | EB | 46 | 47.0 | 46.0-50.2 | 146 | 75-274 | 32 |
| | | | MWF | 1 | 1(CPUE) | n/a | 102 | n/a | <1 |
| East Fork | Cadieux Ranch | 2009 | WCT | 4 | 4.0 | 4.0-4.4 | 249 | 131-318 | 2 |
| | | | LL | 127 | 94.0 | 94.0-94.7 | 168 | 61-338 | 68 |
| | | | EB | 55 | 47.0 | 46.0-50.2 | 122 | 58-268 | 29 |
| | | | MWF | 1 | 1 (CPUE) | n/a | 317 | n/a | <1 |
| East Fork | Cadieux Ranch | 2010 | WCT | 2 | 2.0 | 2.0-4.0 | 295 | 280-309 | 2 |
| | | | LL | 105 | 99.0 | 94.0-106.0 | 170 | 105-355 | 73 |
| | | | EB | 36 | 34.0 | 31.0-40.9 | 138 | 48-250 | 25 |

Table 9 (continued). Electrofishing results from East Fork Rock Creek and Ross Fork Rock Creek. Population estimates of fish only ≥ 75 mm. Corral Gulch section (2007) located just upstream of the State Section (2010 and 2011). (* - indicates section name changed from Christiansen Ranch to Angelico Creek).

| Water Body | Section Name | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (Depletion) | 95% CI Range | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|------------|---------------|--------------|---------|-------------------------|----------------------------|--------------|------------------|-------------------|------------------|
| East Fork | Cadieux Ranch | 2011 | BULL | 2 | 2.0 | n/a | 194 | 186-202 | 2 |
| | | | WCT | 7 | 7.0 | n/a | 276 | 206-347 | 6 |
| | | | LL | 78 | 105.0 | 78.0-139.0 | 195 | 76-397 | 69 |
| | | | EB | 26 | 29.0 | 26.0-36.3 | 91 | 76-206 | 23 |
| East Fork | Corral Gulch | 2007 | BULL | 1 | 1.0 | n/a | 194 | n/a | 1 |
| | | | WCT | 1 | 1.0 | n/a | 173 | n/a | 1 |
| | | | LL | 28 | 28.0 | 28.0-30.2 | 136 | 63-328 | 29 |
| | | | EB | 15 | 15.0 | 15.0-17.5 | 153 | 82-230 | 15 |
| | | | MWF | 52 | 52(CPUE) | n/a | 126 | 76-222 | 54 |
| East Fork | State Section | 2010 | BULL | 1 | 1.0 | n/a | 150 | n/a | <1 |
| | | | LL | 130 | 143.0 | 117.1-168.9 | 122 | 68-434 | 90 |
| | | | EB | 13 | 12.0 | 12.0-13.8 | 134 | 74-206 | 9 |
| East Fork | State Section | 2011 | BULL | 2 | 2.0 | 2.0-2.8 | 167 | 159-176 | 2 |
| | | | WCT | 5 | 5.0 | 5.0-6.5 | 237 | 123-325 | 5 |
| | | | LL | 73 | 90.0 | 73.0-111.6 | 126 | 62-379 | 77 |
| | | | EB | 15 | 15.0 | n/a | 91 | 72-110 | 16 |
| Ross Fork | Above FR 70 | 2010 | BULL | 6 | 6.0 | 6.0-6.7 | 166 | 136-219 | 6 |
| | | | WCT | 30 | 25.0 | 25.0-26.0 | 164 | 86-269 | 29 |
| | | | LL | 34 | 20.0 | 20.0-20.7 | 147 | 55-316 | 33 |
| | | | ONC | 4 | 4 (CPUE) | n/a | 51 | 48-53 | 4 |
| | | | EB | 28 | 24.0 | 24.0-25.5 | 145 | 73-240 | 28 |

Table 9 (continued). Electrofishing results from East Fork Rock Creek and Ross Fork Rock Creek. Population estimates of fish only ≥ 75 mm. Corral Gulch section (2007) located just upstream of the State Section (2010 and 2011). (* - indicates section name changed from Christiansen Ranch to Angelico Creek).

| Water Body | Section Name | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (Depletion) | 95% CI Range | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|------------|--------------|--------------|---------|-------------------------|----------------------------|--------------|------------------|-------------------|------------------|
| Ross Fork | Above FR 70 | 2011 | BULL | 10 | 10.0 | 10.0-10.8 | 149 | 127-170 | 10 |
| | | | WCT | 34 | 34.0 | 34.0-36.1 | 184 | 95-320 | 31 |
| | | | LL | 30 | 30.0 | 30.0-31.6 | 154 | 82-264 | 28 |
| | | | EB | 34 | 34.0 | 34.0-35.3 | 152 | 98-231 | 31 |
| Ross Fork | Angelico Cr* | 2010 | BULL | 4 | 4.0 | n/a | 216 | 162-290 | 5 |
| | | | WCT | 39 | 40.0 | 38.0-45.5 | 157 | 62-285 | 45 |
| | | | LL | 22 | 15.0 | 14.0-19.4 | 143 | 56-408 | 25 |
| | | | ONC | 16 | 16 (CPUE) | n/a | 54 | 42-63 | 18 |
| | | | EB | 4 | 4.0 | n/a | 133 | 75-195 | 5 |
| | | | MWF | 2 | 2 (CPUE) | n/a | 209 | 92-326 | 2 |
| Ross Fork | Angelico Cr* | 2011 | BULL | 7 | 7.0 | 7.0-8.1 | 192 | 90-347 | 8 |
| | | | WCT | 53 | 57.0 | 53.0-64.0 | 155 | 47-292 | 57 |
| | | | LL | 20 | 20.0 | 20.0-20.7 | 148 | 49-298 | 22 |
| | | | EB | 12 | 12.0 | 12.0-13.0 | 155 | 123-197 | 13 |

Harvey Creek

Harvey Creek is a tributary to the Clark Fork River and enters the drainage at approximately RM 257.3 (Figure 1). Land ownership in the upper Harvey Creek drainage is primarily National Forest, while land ownership in the lower portion of the drainage consists of National Forest, Stimson Lumber, and private lands. Primary land uses in the drainage are cattle grazing and timber harvest. At RM 0.2, just upstream from the mouth of Harvey Creek a large fish migration barrier exists and has restricted the movement of non-native fishes into the remainder of the drainage. Fish populations in Harvey Creek above the barrier are comprised exclusively of native bull trout and westslope cutthroat trout (Liermann et al. 2009). Bull trout densities are higher in the upper portions of the drainage and were absent approximately 0.5 mi upstream of the barrier.

In 2010 and 2011, electrofishing surveys were completed in a ditch on lower Harvey Creek (Figure 1). The ditch is located at RM 0.1, just upstream from the mouth of Harvey, but also receives water diverted from the Clark Fork River via an upstream diversion. Water is diverted from the Clark Fork River upstream of the mouth of Harvey Creek and mixes with Harvey Creek at the diversion allowing water from both sources to be diverted into the ditch. This ditch is fitted with a metal slide gate, and a diversion of timber and boulders diverts water from Harvey Creek and water from the Clark Fork River. According to the movements of radio-tagged fish during 2009, 2010 and 2011 the diversion does not likely serve as a barrier to fish passage. This ditch was selected because a radio-tagged bull trout was found dead in the ditch approximately 0.8 mi below the headgate in 2009 (see Radio Telemetry Results). In late August 2010, two surveys were completed in the ditch. In a 100 m section below the headgate, 29 brown trout and 8 unidentifiable *Oncorhynchus* species were collected (Table 10). Slimy sculpin were also present, but rare. The second section was located approximately 1.0 mi below the headgate. A small number of fish were collected in this section, including one unidentifiable *Oncorhynchus* species and a few longnose suckers. Due to the addition of water from the Clark Fork River it is difficult to determine whether these fish originated from Harvey Creek or the Clark Fork River.

In late July 2011, the section directly below the headgate was electrofished to further document entrainment. In the 100 m section below the headgate 28 westslope cutthroat trout, 2 brown trout were collected in addition to mountain whitefish, longnose dace, longnose suckers, redbreast shiner and slimy sculpin (Table 10). In late August 2011, both the section directly below the headgate and the 1.0 mi section were re-surveyed, but significantly fewer fish were captured, however the headgate was closed and little water was entering the ditch. In the 100 m below the headgate only one westslope cutthroat trout and several slimy sculpin were collected. In a 100 m section approximately 1.0 mi downstream of the headgate, one longnose sucker was collected.

Table 10. Electrofishing data collected during 2010 and 2011 in a ditch on Harvey Creek. ((2) – denotes the second survey in 2011 during late August)

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|----------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 0.01 (Headgate) | 2010 | LL | 29 | 29 | 80 | 66-93 | 78 |
| | | ONC | 8 | 8 | 52 | 39-60 | 22 |
| | | SL COT | C | - | - | - | - |
| RM 0.01 (1.0 mi) | 2010 | ONC | 1 | 1 | 46 | 46 | 100 |
| | | LN SU | R | - | - | - | - |
| RM 0.01 (Headgate) | 2011 | WCT | 28 | 28 | 158 | 92-326 | 93 |
| | | LL | 2 | 2 | 131 | 69-193 | 7 |
| | | MWF | R | - | - | - | - |
| | | LN DC | R | - | - | - | - |
| | | LN SU | R | - | - | - | - |
| | | RS SH | R | - | - | - | - |
| | | SL COT | A | - | - | - | - |
| RM 0.01 (Headgate) | 2011 (2) | WCT | 1 | 1 | 115 | 115 | 100 |
| | | SL COT | C | - | - | - | - |
| RM 0.01 (1.0 mi) | 2011 (2) | LN SU | R | - | - | - | - |

The discharges of Harvey Creek upstream of the diversion and within the ditch just below the headgate were measured on August 30, 2010. The ditch was receiving 9.5 cfs, approximately 185% of the discharge of Harvey Creek at 5.1 cfs. Thus, the joining ditch contributed at least 4.5 cfs of Clark Fork River water to the ditch. This ditch intersects Harvey Creek approximately 20 m upstream of its mouth before it enters the Clark Fork River. On September 2, 2010, the Clark Fork River ditch was carrying 9.7 cfs.

On August 26, 2011, the ditch was receiving 9.7 cfs from Harvey Creek (and the Clark Fork River), while the discharge of Harvey Creek was an estimated 10.0 cfs.

South Fork Lower Willow Creek

South Fork Lower Willow Creek is a tributary of Lower Willow Creek and enters Lower Willow Creek Reservoir at approximate RM 9.4 (Figure 1). Land ownership above the reservoir is divided between private agricultural, USFS, and private timber land. Cattle grazing is occurring throughout the drainage on both private and USFS lands, and signs of past timber harvest and mining also exist. Fish populations above the dam are comprised primarily of westslope cutthroat trout, with some tributaries also containing native longnose sucker (Lindstrom et al. 2008). Genetic analyses conducted on several tributary populations of westslope cutthroat trout above the dam, have found these

populations to be genetically pure. It is hypothesized the Lower Willow Creek dam is preserving the genetic purity of this large population of westslope cutthroat trout by inhibiting upstream migration of non-native rainbow trout. Non-native brook trout are also present above the dam however, their distribution appears limited to the North and West Fork Lower Willow Creek.

At least one irrigation ditch exists above the Lower Willow Creek Reservoir (Figure 1). The ditch draws water from the South Fork Lower Willow Creek above its confluence with Cottonwood Creek at approximately RM 2.9, and routes water to a ranch several miles downstream near mainstem Flint Creek. Electrofishing surveys were completed in this ditch during 2007 and 2009. In 2007, a 200 m section was electrofished approximately two miles below the headgate (Table 11). At this site, two similar-sized westslope cutthroat trout were collected along with 38 longnose suckers. In 2009, a three-pass depletion was conducted on a section beginning 100 m downstream of the headgate and extending upstream to the headgate. A total of 18 westslope cutthroat trout and 53 longnose suckers were collected in the ditch (Table 11, Appendix A). There were an estimated 19.0 (18.0-22.9, 95% C.I.) westslope cutthroat trout per 100 m in this section of the ditch.

Table 11. Electrofishing data collected during 2007 and 2009 in a ditch on South Fork Lower Willow Creek. Entries are arranged in the table chronologically.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (Depletion) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|----------------------|--------------|---------|-------------------------|----------------------------|------------------|-------------------|------------------|
| RM 2.9 (2.0 mi) | 2007 | WCT | 2 | 1 (CPUE) | 142.5 | 142-143 | 5 |
| | | LN SU | 38 | 19 (CPUE) | - | - | 95 |
| RM 2.9 (Headgate) | 2009 | WCT | 18 | 19.0 | 92 | 37-185 | 25 |
| | | LN SU | 53 | - | 62 | 25-177 | 75 |

Discharge was measured in mainstem South Fork Lower Willow Creek and within the ditch on August 24, 2009. The ditch was receiving 4.0 cfs, approximately 98% of the mainstem discharge at 4.1 cfs. Below the diversion, the creek was de-watered and likely leading to increased water temperatures, stressful for westslope cutthroat trout (Appendix B). In 2007, a thermograph was positioned in South Fork Lower Willow Creek just above its confluence with the reservoir. The maximum daily temperatures exceeded 20 °C for 23 days with a maximum recorded temperature was 25.9 °C (Appendix C).

Gird Creek

Gird Creek is a tributary to Flint Creek, and enters the drainage at approximately RM 13.7 (Figure 1). Most of upper Gird Creek is located within the Beaverhead-Deerlodge National Forest, while the lower portion runs through private lands. Cattle grazing is the primary land use on private land as well as National Forest land via grazing allotments. Recent electrofishing surveys found Gird Creek to support only westslope cutthroat trout (FWP unpublished data).

In 2009, one irrigation ditch was observed on lower Gird Creek near RM 1.1 (Figure 1). This ditch is fitted with a metal slide gate and a check dam spanning the channel redirects water into the ditch (Appendix B). The diversion may serve as a barrier to fish passage at low discharges. One section extending 100 m below the headgate was electrofished on this ditch in 2009 and no fish were captured. This ditch was visited again in 2010, but was not in operation at the time.

Boulder Creek

Boulder Creek is a tributary to Flint Creek, and enters the drainage near Maxville, MT at approximately RM 15.7 (Figure 1). The upper portion of Boulder Creek is primarily National Forest land, while the lower portion (below Princeton, MT) is a mix of private land and National Forest Land. Historical land use in the drainage was mining and timber harvest, with abandoned mines visible on many tributaries and some sections of Boulder Creek. Current land use is timber harvest, small-scale hydropower and private residences. Boulder Creek supports the only known population of bull trout in the Flint Creek drainage. In upper Boulder Creek, fish populations are comprised entirely of bull trout and westslope cutthroat trout. Brook trout first appear in the middle section near the confluence of Copper Creek. In the lower portion of the drainage, westslope cutthroat trout as well as brown trout are abundant while bull trout are found in relatively low densities (Lindstrom et al. 2008).

Three ditches draw water from the lower portion of Boulder Creek (Figure 1). Electrofishing surveys were completed in two of these ditches during 2009 and 2010, and the third during 2010. All three ditches have headgates and boulder berms as diversions. None of the diversions are likely barriers to fish passage. The furthest upstream ditch at approximately RM 0.9 is fitted with a metal slide gate, and diverts the second-most volume of water. In 2009, one bull trout, five westslope cutthroat trout and five brown trout were collected the first 100 m below the headgate (Table 12, Appendix A). In 2010, no bull trout were observed, but six westslope cutthroat and nine brown trout were collected, along with mountain whitefish and slimy sculpin. The next ditch downstream, located at approximately RM 0.8, is also fitted with a metal slide gate and diverts the most volume of water. In 2009, two sections were sampled in this ditch, one near the headgate and one approximately a mile downstream near its confluence with Gird Creek. Fish were only collected near the headgate, with seven westslope cutthroat trout captured in a 100 m section (Table 12, Appendix A). In 2010, two additional species were observed entrained in the ditch. In addition to six westslope cutthroat trout, thirty-nine brown trout and one brook trout were collected. More juvenile trout were observed but escaped capture due to their small size. The lowest ditch, located at approximately RM 0.2 is in great disrepair and highly inefficient based on visual observations. The headgate is no longer functional and water enters through a naturally cut side channel. Little water is retained in the ditch due to several gaps in the walls of the ditch. Only 40.5 m was open to electrofishing, before the ditch enters a large pipe. In this section a total of eight trout were collected, including six westslope cutthroat trout and two brown trout (Table 12, Appendix A). Slimy sculpin were also present, but rare.

Table 12. Electrofishing data collected during 2009 and 2010 in two ditches on Boulder Creek (*- Indicates estimates were extrapolated from a 40.5 m section).

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|----------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 0.9 (Headgate) | 2009 | BULL | 1 | 1 | 186 | 186 | 9 |
| | | WCT | 5 | 5 | 141 | 60-269 | 45 |
| | | LL | 5 | 5 | 201 | 92-292 | 45 |
| RM 0.9 (Headgate) | 2010 | WCT | 6 | 6 | 101 | 67-139 | 40 |
| | | LL | 9 | 9 | 66 | 41-151 | 60 |
| | | MWF | R | - | - | - | - |
| | | SL COT | R | - | - | - | - |
| RM 0.8 (Headgate) | 2009 | WCT | 7 | 7 | 60 | 31-125 | 100 |
| RM 0.8 (Headgate) | 2010 | WCT | 6 | 6 | 142 | 33-277 | 13 |
| | | LL | 39 | 39 | 42 | 30-54 | 85 |
| | | EB | 1 | 1 | 135 | 135 | 2 |
| RM 0.8 (Gird Ck) | 2009 | NO | FISH | - | - | - | - |
| RM 0.2 (Headgate) | 2010 | WCT | 6 | 15* | 114 | 70-180 | 75 |
| | | LL | 2 | 5* | 144 | 60-227 | 25 |
| | | SL COT | R | - | - | - | - |

In 2010, discharge was measured in all three ditches. On August 10, the ditches at RM 0.9 and RM 0.8 were receiving 2.4 cfs and 14.1 cfs, respectively, and the United States Geological Survey (USGS) gauging station (downstream of the ditches) recorded 33 cfs. Based on these flow measurements, the mainstem discharge upstream of the two withdrawals was estimated as 49.5 cfs (USGS gage data plus ditch discharge data). Based on these estimates, the ditch at RM 0.9 was receiving approximately 5% of the mainstem discharge and the ditch at RM 0.8 was receiving approximately 30% of the remaining mainstem discharge. The ditch at RM 0.2 was electrofished the following week on August 18 and was receiving an estimated 0.9 cfs. The USGS station upstream recorded 25 cfs in Boulder Creek, and thus it is estimated the ditch was receiving less than 4% of the mainstem discharge.

Gold Creek

Gold Creek is a tributary to the Clark Fork River and enters the drainage at approximately RM 291.6 (Figure 1). Upper Gold Creek is located on National Forest land, but the remainder of the watershed is privately owned. Land use consists of hay production, cattle grazing, and timber harvest, but historical mining is evident. Fish

populations in Gold Creek vary from the upper reaches on National Forest land to the lower reaches on heavily grazed private lands. Westslope cutthroat trout is the only species present at upper sites where the creek is best described as a high gradient mountain stream (Lindstrom et al. 2008). In the middle section, as the creek transitions from the mountains to agricultural land, westslope cutthroat trout are still dominant, but brown trout begin to appear. Conversely, in the lower section as Gold Creek nears the Clark Fork River, brown trout are the dominant species with few westslope cutthroat trout present.

In 2007, five irrigation ditches were observed on lower Gold Creek. Four of these diversions upstream of RM 3.6 drew a significant amount of water from Gold Creek. Sediment accumulation and low in-stream flow was notable below these diversions. The stream channel was observed to be dry for a distance below the lowest of these diversions in 2007. The extent of this dewatering was not determined, but it is likely that it represents a significant seasonal migration barrier to fish in Gold Creek (Lindstrom et al. 2008). It is also likely such reductions in stream discharge result in increased water temperatures. In 2007, thermographs were placed in Gold Creek at approximate RM 0.5 and RM 5.7 (Appendix C). Above the diversions (RM 5.7), water temperatures only exceeded 15°C on ten occasions with a maximum recorded temperature of 16.1°C. At the lower site however (RM 0.5), temperatures in Gold Creek exceeded 15°C on 65 days including eight days in which they exceeded 20°C. Maximum recorded temperature at this site was 21.2°C.

Electrofishing surveys were completed in one ditch on Gold Creek at RM 3.6 during 2007 and 2009 (Table 13, Appendix A). Multiple-pass depletions were conducted at each of three locations on the ditch. Fish densities were highest near the headgate, with westslope cutthroat trout and brown trout captured both years. In 2007, depletion estimates at the headgate site were 7.3 (7.3-8.7; 95% CI) westslope cutthroat trout per 100 m and 40.7 (40.7-47.0; 95% CI) brown trout per 100 m. In 2009, fewer fish were collected and population estimates were 1.0 cutthroat trout and 22.0 brown trout per 100 m. At the lower sites, very few fish were captured in 2007 and no fish were captured in 2009 (Table 13). It is likely more fish occur at these lower sites, but high water velocities and lack of holding water force fish downstream until the ditch empties into an agricultural field.

Table 13. Electrofishing data collected during 2007 and 2009 in a ditch on Gold Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (Depletion) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|----------------------------|-------------------------------|---------------------|----------------------|---------------------|
| RM 3.6 (Headgate) | 2007 | WCT | 11 | 7.3 | 104 | 77-217 | 17 |
| | | LL | 55 | 40.7 | 79 | 38-184 | 83 |
| RM 3.6 (Headgate) | 2009 | WCT | 1 | 1.0 | 86 | 86 | 4 |
| | | LL | 21 | 22.0 | 88 | 45-320 | 96 |

Table 13 (continued). Electrofishing data collected during 2007 and 2009 in a ditch on Gold Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (Depletion) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|----------------------------|------------------|-------------------|------------------|
| RM 3.6 (1.8 mi) | 2007 | WCT | 2 | 2.0 | 175 | 167-183 | 66 |
| | | LL | 1 | 1.0 | 118 | 118 | 33 |
| RM 3.6 (1.8 mi) | 2009 | NO | FISH | - | - | - | - |
| RM 3.6 (2.1 mi) | 2007 | LL | 1 | 1.0 | 157 | 157 | 100 |
| RM 3.6 (2.1 mi) | 2009 | NO | FISH | - | - | - | - |

Discharge was measured as 6.4 cfs in the meadow section of the ditch at RM 3.6 on August 3, 2009.

In 2010, electrofishing surveys were completed in six additional ditches on Gold Creek. These ditches were located on private land from RM 1.1 to 6.1, with all diverting water from the mainstem using a variety of methods (Figure 1, Appendix B). In each ditch, one 100 m section was electrofished below the point of diversion. The furthest upstream ditch at RM 6.1 has no formal headgate or diversion (Appendix B). The ditch is closed off via a gravel push-up dam and the elevation of Gold Creek is similar to the ditch, so with the natural gradient water flows into the ditch. In the 100 m below the start of the ditch, fifteen westslope cutthroat trout and eight brown trout were collected (Table 15, Appendix A). The next ditch downstream, at RM 5.2, is fitted with a metal slide gate and a boulder berm diverts water from Gold Creek (Appendix B). Westslope cutthroat trout and brown trout were also present below the headgate in this ditch, but brown trout were more abundant. A total of eight westslope cutthroat trout and 11 brown trout were collected (Table 14, Appendix A). The ditch at RM 3.9 has no formal headgate or diversion structure (Appendix B). Similar to the ditch at RM 6.1 channel and ditch elevations are similar so water enters the ditch via the natural gradient. At the time of survey, this ditch was blocked by a combination of plywood, fenceposts, and debris, allowing minimal flow into the ditch. In the 100 m section below the opening of the ditch, 16 westslope cutthroat trout, 120 brown trout and six unidentifiable *Oncorhynchus* species were collected. Slimy sculpin were also present but rare in the section (Table 14, Appendix A). The ditch at RM 3.8 is fitted with a wooden gate and a boulder berm reinforced with a large wooden plank serves as the diversion structure (Appendix B). Within this ditch, 19 westslope cutthroat trout, 158 brown trout and four *Oncorhynchus* species were collected, along with several slimy sculpin (Table 14, Appendix A). The ditch at RM 3.7 has no formal headgate, but receives water diverted from Gold Creek via a boulder berm spanning the channel (Appendix B). A total of 28 westslope cutthroat trout and 41 brown trout were collected in the 100 m section. Again slimy sculpin were

common (Table 14, Appendix A). The ditch at RM 3.6 is fitted with a metal slide gate and a gravel push-up dam diverts water from Gold Creek (Appendix B). At the time of the survey the headgate was closed and the push-up dam had been removed. The 100 m section below the headgate was still wetted and 32 westslope cutthroat trout, 145 brown trout, and eight unidentified *Oncorhynchus* species were collected from the remaining pools. Slimy sculpin were also common in this ditch (Table 14, Appendix A). The lowest of the ditches, at RM 1.1, is fitted with a metal slide gate and a boulder berm reinforced with timber and Visqueen diverts water. At this location, a horizontal flat-plate fish screen and a secondary headgate has been installed within the ditch below the headgate (Appendix B). A total of 161 brown trout were collected in the 70 m above the fish screen (Table 14, Appendix A). At the time of the survey, water was flowing over the screen and the structure was clear of debris.

Table 14. Electrofishing data collected during 2010 in ditches on Gold Creek (* - Indicates average or range based on the measurement of a sub-set of fish, and ¹ - indicates estimates were extrapolated from a 70 m section).

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|----------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 6.1 (Headgate) | 2010 | WCT | 15 | 15 | 116 | 46-167 | 65 |
| | | LL | 8 | 8 | 134 | 61-231 | 35 |
| RM 5.2 (Headgate) | 2010 | WCT | 8 | 8 | 165 | 102-245 | 42 |
| | | LL | 11 | 11 | 84 | 47-256 | 58 |
| RM 3.9 (Headgate) | 2010 | WCT | 16 | 16 | 93 | 82-107 | 11 |
| | | LL | 120 | 120 | 66* | 51-125* | 85 |
| | | ONC | 6 | 6 | 33 | 31-37 | 4 |
| | | SL COT | R | - | - | - | - |
| RM 3.8 (Headgate) | 2010 | WCT | 19 | 19 | 108 | 77-183 | 11 |
| | | LL | 158 | 158 | 90* | 41-216* | 87 |
| | | ONC | 4 | 4 | 39 | 34-45 | 2 |
| | | SL COT | C | - | - | - | - |
| RM 3.7 (Headgate) | 2010 | WCT | 28 | 28 | 106 | 72-190 | 41 |
| | | LL | 41 | 41 | 121* | 61-159* | 59 |
| | | SL COT | C | - | - | - | - |
| RM 3.6 (Headgate) | 2010 | WCT | 32 | 32 | 94 | 81-137 | 17 |
| | | LL | 145 | 145 | 76* | 48-140 | 79 |
| | | ONC | 8 | 8 | 37* | 30-42* | 4 |
| | | SL COT | C | - | - | - | - |
| RM 1.1 (Headgate) | 2010 | LL | 161 | 230 ¹ | 103* | 55-206* | 100 |

In August 2010, measurements of discharge were collected above, within, and below the seven ditches on Gold Creek by Montana TU. On August 7, the discharges of the ditches at RM's 3.9, 3.8, 3.7, and 3.6 were measured, and the discharges in ditches at RM's 6.1, 5.2 and 1.1 were measured the following day. The ditch at RM 6.1 was receiving 3.2 cfs, approximately 17% of the mainstem discharge at 18.3 cfs. The ditch at 5.2 was receiving 8.3 cfs, approximately 62% of the mainstem discharge at 13.3 cfs. The ditch at RM 3.9 was receiving less than 1 cfs, which was approximately 9% of the mainstem discharge at 9.7 cfs. The ditch at RM 3.8 was receiving 1.3 cfs, approximately 13% of the mainstem discharge at 10.4 cfs. The ditch at RM 3.7 was receiving 4.9 cfs, approximately 61% of the mainstem discharge at 8.1 cfs. The ditch at RM 3.6 was receiving less than 1 cfs, less than 27% of the mainstem discharge at 3.3 cfs. However, the push-up dam diversion for the Lower Wall City ditch had been removed at the time measurements were collected. Approximately 5.2 cfs was flowing through the ditch at RM 1.1, but the center-pivot was not in operation and all water was returning to Gold Creek.

Clark Fork River

The Clark Fork River is a major tributary of the Columbia River, flowing through Montana and Idaho. The headwaters of the Clark Fork River, beginning as Silver Bow Creek, are located near the town of Butte, MT (Figure 1). Land use within the upper Clark Fork River basin consists primarily of agriculture, but includes timber harvest and historical mining. Although crop production is limited, irrigation is extensive due to the prevalence of cattle production. Within the upper Clark Fork River, brown trout are the dominate trout species, and westslope cutthroat and rainbow trout are present in lower densities. Although rare, bull trout are also present below Drummond, MT.

In the August of 2010, electrofishing surveys were completed in a ditch diverting water from the Clark Fork River at RM 314.0 at the town of Deer Lodge, MT (Figure 1). This ditch, known as the Kohrs-Manning ditch, provides water to several private water users and the Grant Kohrs Ranch administered by the U.S. National Park Service. This large ditch is fitted with two metal slide gates and water is diverted from the Clark Fork River with a boulder diversion (Appendix B). The diversion on the Clark Fork River is not a barrier to fish passage. The ditch intersects the extreme lower end of Cottonwood Creek approximately 0.2 mi below the headgate. At this junction a diversion near the mouth of Cottonwood Creek redirects water from the creek into the Kohrs-Manning ditch (Appendix B). At the time of the survey the headgates of the Kohrs-Manning ditch were turned down and little water was entering the ditch. Also, little to no water was advancing past the diversion at the mouth of Cottonwood Creek, and nearly all the water flowing out from Cottonwood Creek was entering the Kohrs-Manning ditch. The diversion at the mouth of Cottonwood Creek is certainly a fish barrier. Three 100 m sections were sampled in the ditch, one immediately below the headgate, one below the ditch's intersection with Cottonwood Creek and one approximately 1.6 mi below the headgate. No native trout were found entrained in the Kohrs-Manning ditch, but several other native fishes were observed (Table 15, Appendix A). In the section below the headgate 84 brown trout were collected, unidentified native sucker species and reddsides were found to be abundant, and mountain whitefish, longnose dace, and slimy

sculpin were also present, but rare. Below the Cottonwood Creek intersection, 20 brown trout and a few native sucker were collected. Approximately 1.6 mi below the headgate, although only one adult brown trout was collected, native sucker and reidside shiner were found to be more common. Due to the location and design of the Cottonwood Creek diversion, it is difficult to determine where fishes captured below the ditch's intersection with the creek may have originated.

Table 15. Electrofishing data collected during 2010 in Kohrs-Manning ditch on the Clark Fork River. Longnose sucker and largescale sucker (native spp.) were combined as unspecified sucker (SU) due to their abundance.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|-----------------|---------|-------------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 314.0 (Headgate) | 2010 | LL | 84 | 84 | 107 | 68-207 | 100 |
| | | MWF | R | - | - | - | - |
| | | LN DC | R | - | - | - | - |
| | | SU | A | - | - | - | - |
| | | RS SH | A | - | - | - | - |
| | | SL COT | R | - | - | - | - |
| RM 314.0 (0.25 mi) | 2010 | LL | 20 | 20 | 100 | 71-134 | 100 |
| | | SU | R | - | - | - | - |
| RM 314.0 (1.6 mi) | 2010 | LL | 1 | 1 | 272 | 272 | 100 |
| | | SU | C | - | - | - | - |
| | | RS SH | C | - | - | - | - |

In September 2010, discharges were measured by Montana TU, a few days before the sections were electrofished. On September 13, the discharge in the ditch below the headgate was approximately 1.0 cfs, and discharge of Cottonwood Creek was 6.0 cfs. The discharge in the ditch just below the intersection with Cottonwood Creek was 7.6 cfs, indicating all water from the creek was making its way into the Kohrs-Manning ditch.

Cottonwood Creek

Cottonwood Creek drains for over nine miles before reaching the Clark Fork River at RM 313.8 near the town of Deer Lodge (Figure 1). Lands along Cottonwood Creek are dominated by private agricultural lands, and rural and urban residences. Public ownership along the stream is limited to the lower 0.3 miles of channel through the Grant Kohrs Ranch administered by the National Park Service, and the upper 0.3 miles of stream through U.S. Forest Service land. Cottonwood Creek is extensively channelized through Deer Lodge (RM 0.3 to RM 1.4). The dominant land uses in the Cottonwood Creek drainage are cattle grazing, hay production, urbanization, timber harvest, and historic mining. Fish populations in Cottonwood Creek vary from its upper reaches to the mouth. In 2007 and 2009 surveys, westslope cutthroat trout were only found in high densities in the upper reaches of Cottonwood Creek (RM 6.9; Lindstrom et al. 2008;

unpublished data). On lower sections of the creek, cutthroat trout were rare or absent. Brook trout were common throughout the upper and middle reaches, and brown trout numbers were highest near the mouth. Other species observed were slimy sculpin, longnose sucker and redbreast shiner, which were only present near the mouth of the creek.

Multiple ditches and irrigation structures are located throughout the middle to lower reaches of Cottonwood Creek. Seven of these diversions and ditches were observed between approximately RM 6.0 and RM 2.5 (Figure 1). These structures varied in terms of size, construction, and amount of water diverted (Appendix B). The effects of these structures on Cottonwood Creek water temperatures and fish populations are likely high. In 2007, thermographs were placed in Cottonwood Creek at RM 7.0 and RM 0.5 (Appendix C). Water temperatures were found to be notably cooler at the upper site. At RM 7.0, maximum temperature recorded did not exceed 17.8°C, while at RM 0.5 it exceeded 20°C on 20 days with a maximum recorded temperature of 24.1°C. On June 30, 2009, Cottonwood Creek was found to be dry for several hundred meters below the diversion at RM 4.0 (Appendix B).

Cottonwood Creek was visited on five occasions during the summer of 2009 to assess the effects of the diversion structures on fish populations in Cottonwood Creek. A study was conducted to establish if any structures were fish passage barriers. At the initial visit on June 23, 2009, it was visually determined that six of the 11 diversions were not likely barriers, and mark/recapture studies were conducted on the remaining five. Fish were collected above the diversions, given a unique mark, and transported downstream of each diversion. In the subsequent weeks, extensive electrofishing was performed above the diversions to identify movement across potential barriers. Table 16 below describes characteristics of these diversions and results from this study.

Table 16. Physical characteristics of selected diversion structures on Cottonwood Creek with results from mark/recapture fish passage barrier assessment conducted during 2009.

| Diversion Location (Name) | Diversion Type | Jump Pool | Jump Height | Fish Relocated Below | Fish Captured Above |
|---------------------------|------------------------|-----------|-------------|----------------------|---------------------|
| RM 6.0 (Dippold/Pruyn) | Pin and Plank | <1" | 1' 5.5" | 25 | 4 |
| RM 5.5 (Olsen) | Concrete Dam | 3' 0" | 2' 7" | 30 | 1 |
| RM 5.4 (Pruyn) | Pin and plank | <1" | 2' 1" | 44 | 1 |
| RM 3.5 (Burt) | Concrete rip-rap | n/a | n/a | 17 | 2 |
| RM 3.0 (Lower Applegate) | Dirt, Sod, and rip-rap | n/a | n/a | 25 | 6 |

In summary, none of the structures were found to be complete barriers to fish migration. The diversions at RM 6.0 and RM 5.4 were both pin and plank structures, and appeared to be barriers when all planks were in place. However, on July 15, 2009, at least one plank from each of these diversions had been removed which provided easy passage for fish. The diversion at RM 5.5 was the tallest, most permanent structure and considered most likely to be a seasonal barrier in 2009 (Appendix B). However, on July 15 a 153 mm EB placed below the diversion at RM 5.4 was recaptured above the diversion at RM 5.5. Regardless, in the fall of 2011, the diversion at RM 5.5 was replaced with a passage friendly rock v-weir structure. In 2009, multiple fish also navigated past the two lowest diversions at RM 3.5 and RM 3.0. A final barrier observed during 2009, was the several hundred meters of dry creek bed below the diversion at RM 3.5. This stretch was certainly a barrier for the seven or more days when dry, and may be dry for longer periods during low-water years. Furthermore, when surveyed on June 30, hundreds of trout fry were stranded in pools within this section of creek. Approximately 20 of these fry were positively identified as brook trout, but it is likely a small percentage were westslope cutthroat trout given their presence in this reach of Cottonwood Creek.

Also during 2009, electrofishing surveys were completed in four ditches on Cottonwood Creek to document entrainment (Figure 1). Due to fluctuating irrigation demands, certain ditches were closed off at various times throughout the field season. The ditch at RM 6.1 was only sampled for 61 m below the headgate due to private property constraints. The ditch at RM 4.0 was dry except for a small pool directly below the headgate. The two lower ditches were shocked from the headgate downstream 100 m. No other ditches were sampled either due to lack of landowner permission or water on the intended sampling date. All ditches sampled contained fish with the lower three containing non-native brook trout and the lowest also containing a brown trout (Table 17, Appendix A). The upper ditch (RM 6.1) contained one brook trout and multiple (approximately 20) smaller trout (<30 mm), one of which was identified as a westslope cutthroat trout. These fish were too small for the dip net and escaped capture; however, all were assumed to be age-0 westslope cutthroat trout due to their size. Additionally, a 135 mm westslope cutthroat trout was shocked in the ditch structure just above the headgate. The ditch at RM 3.5 also contained a single adult Columbia spotted frog (*Rana luteiventris*).

In 2011, electrofishing surveys were completed on two additional ditches and one ditch was re-surveyed. The furthest upstream ditch, at RM 9.0, is fitted with a wooden gate, regulated by wooden planks and a boulder diversion spanning the channel (Appendix B). One westslope cutthroat trout and one brook trout were captured in first 100 m below the headgate. Slimy sculpin were also captured, but were in low abundance. The second ditch, located at RM 5.6, is fitted with a metal slide gate and a constructed channel directs water from Cottonwood Creek into the ditch (Appendix B). Two westslope cutthroat trout, 34 brook trout and several slimy sculpin were collected in the first 100 m below the headgate of the ditch (Appendix A, Table 17). The results of electrofishing the ditch at RM 3.0 in 2011 were very similar to those in 2009, with one brown trout and six brook trout captured in 100 m below the headgate. It was noted the diversion for the RM 3.0 ditch had recently been re-built with sod, boulders and slabs of concrete.

Table 17. Electrofishing data collected in ditches on Cottonwood Creek during 2009 and 2011.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-----------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 9.0 (Headgate) | 2011 | WCT | 1 | 1 | 106 | 106 | 50 |
| | | EB | 1 | 1 | 86 | 86 | 50 |
| | | SL COT | R | - | - | - | - |
| RM 6.1 (Headgate) | 2009 | WCT | 1 | n/a | 27 | 27 | 50 |
| | | EB | 1 | 1.6 | 120 | 120 | 50 |
| RM 5.6 (Headgate) | 2011 | WCT | 2 | 2 | 76 | 69-82 | 6 |
| | | EB | 34 | 34 | 98 | 34-257 | 94 |
| | | SL COT | R | - | - | - | - |
| RM 4.0 (Headgate) | 2009 | EB | 8 | n/a | 70 | 43-122 | 100 |
| RM 3.5 (Burt) | 2009 | EB | 11 | 11 | 111 | 91-187 | 100 |
| RM 3.0 (Lower Applegate) | 2009 | LL | 1 | 1 | 156 | 156 | 11 |
| | | EB | 8 | 8 | 143 | 60-156 | 89 |
| RM 3.0 (Lower Applegate) | 2011 | LL | 1 | 1 | 142 | 142 | 14 |
| | | EB | 6 | 6 | 92 | 51-160 | 86 |

In late August and early September 2010, discharges were measured by Montana TU at several of the sites electrofished in 2009. On August 30, the headgate on the Dippold/Pruyn ditch at RM 6.0 was closed, but nearly 1 cfs continued to enter the ditch. According to measurements upstream of 4.4 cfs and downstream of 4.0 cfs, the ditch was receiving 9% or less of the mainstem discharge. The Olsen ditch at RM 5.5 was receiving 0.2 cfs on September 14th, approximately 4% of the mainstem discharge of 4.9 cfs. The headgate on the Pruyn ditch at RM 5.4 was closed on September 14th, but an estimated 1.5 cfs continued to enter the ditch. The Burt ditch at RM 3.5 was receiving <1 cfs on August 24, approximately 6% or less of the mainstem's estimated discharge of 1.8 cfs. The Lower Applegate ditch at RM 3.0 was receiving 2.6 cfs on August 24, approximately 93% of the remaining mainstem discharge of 2.8 cfs.

In 2011, discharges were measured by Montana TU at several of the sites electrofished. The ditch at RM 5.6 was receiving 25.3 cfs from Cottonwood Creek. The ditch at RM 5.6 may also receive water from Baggs Creek when a diversion upstream in Baggs Creek is not under operation.

Baggs Creek

Baggs Creek is a tributary to Cottonwood Creek flowing for approximately 8 miles before entering Cottonwood Creek. The upper portion of Baggs Creek is located on U.S. Forest Service land, while the lower 2 miles flows through private property. The Baggs Creek watershed is primarily used for grazing and recreation. Fish surveys completed in Baggs Creek during 2007 and 2008 found the fish community in an upper section (RM 5.4) is comprised entirely of westslope cutthroat trout, but the community shifts toward brook trout dominated in a lower section (RM 0.5; Lindstrom et al. 2008; Liermann et al. 2009). During the 2007 survey a waterfall was discovered at RM 5.3. Brief electrofishing surveys above and below the feature indicate it is a barrier to upstream migration.

In July 2011, two irrigation ditches diverting water from Baggs Creek were visited. The upstream ditch at RM 0.4 is fitted with a wooden gate regulated by the installation of wooden planks and a diversion comprised of concrete slab, timber and Visqueen. The stream was nearly de-watered below the diversion, likely serving as a fish passage barrier when in operation. The RM 0.4 ditch was accessed on private land and three westslope cutthroat trout and 12 brook trout were collected in first 100 m below the headgate (Table 18, Appendix A). The lower ditch, located just downstream at RM 0.35, had recently been closed and only pools remained below the headgate. Within a section approximately 60 m in length six small westslope cutthroat trout (57-90mm) and one small brook trout (44mm) were collected from these pools. The lower ditch was accessed from a county road below the headgate.

Table 18. Electrofishing data collected in a ditch on Baggs Creek during 2011.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 0.4 (Headgate) | 2011 | WCT | 3 | 3 | 161 | 100-222 | 20 |
| | | EB | 12 | 12 | 138 | 112-193 | 80 |

Discharge measurements were also collected at the upper ditch by Montana TU and FWP in 2011. On July 10, the ditch at RM 0.4 was receiving 8.1 cfs.

Dempsey Creek

Dempsey Creek is a tributary to the Clark Fork River and drains for over 16 miles before reaching the Clark Fork River near RM 322.8 (Figure 1). The lower 10 miles of the stream flows through privately owned lands with the exception of property managed by the Montana State Prison (between RM 5.3 and 3.6). The upper extent of the watershed lies entirely on high elevation lands managed by the USFS. The primary land uses in the drainage are irrigated hay production (lower half of drainage), livestock grazing, and National Forest recreation (upper portion of drainage). Dempsey Creek is heavily used for irrigation, and diversions are common throughout the lower portion of the drainage. There are also a number of high elevation lakes in the headwaters of the drainage, some

regulated to provide summer flows for downstream irrigators. Fish populations in Dempsey Creek vary from its upper to lower reaches. The upper reaches support mostly brook trout with a few westslope cutthroat trout also present, while the lower reaches support mainly brown trout (Liermann et al. 2009).

In early September 2010, four irrigation ditches on lower Dempsey Creek were sampled within land managed by the Montana State Prison (Figure 1). Electrofishing surveys were completed in 100 m sections on all of the ditches. The furthest upstream ditch, at approximately RM 5.1, is fitted with a wooden slide gate and a check dam which spans the creek channel (Appendix B). The check dam does not likely serve as barrier. The headgate was opened the day before electrofishing was conducted and although no fish were captured below the headgate, one small (<80 mm) fish was observed. The next ditch downstream at approximately RM 4.8 is fitted with metal slide gate and a boulder diversion, reinforced by natural vegetation, diverts water from Dempsey Creek. The diversion does not serve as a barrier. At the time of survey, the headgate was closed and no water was being diverted from Dempsey Creek. The remaining pools in the 100 m below the headgate however, contained 85 brown trout, four brook trout and several slimy sculpin (Table 19, Appendix A). It is likely the ditch at RM 4.8 was in operation more regularly earlier in the summer, allowing for the observed entrainment of brown trout, brook trout and slimy sculpin. The ditch at RM 4.7 has no formal headgate, but is fitted with a concrete check-dam to divert water from Dempsey Creek. At the time of the survey, no planks were installed in the check dam and the ditch was closed off with Visqueen. The check dam constricts the channel and may have presented a velocity barrier at the time of the survey. No fish were collected in the first 100 m below the opening of the ditch. The lowest ditch, at RM 4.3, was in operation. This ditch has no formal headgate, but is fitted with a wooden check-dam. This check dam also constricts the channel and creates conditions difficult for fish to pass the structure. No fish were collected in this ditch; however, this ditch had only been opened for a short time.

In early August 2011, two additional ditches on lower Dempsey Creek were surveyed (RM's 2.0 and 1.0). The upper ditch at RM 2.0 has no formal headgate and a temporary diversion had been constructed of lumber and Visqueen at the time of the survey (Appendix B). The diversion does not serve as a fish passage barrier. One red side shiner was the only fish captured in a 100 m section below the head of the ditch (Table 19). The ditch at RM 1.0 is fitted with a metal slide gate and a temporary diversion with timber, wood planks and plastic sheeting (Appendix B). The diversion is not a permanent barrier, but may periodically inhibit migration. In 100 m below the headgate one brook trout was collected along with a two longnose suckers and seven reidside shiners.

Table 19. Electrofishing data collected during 2010 and 2011 in ditches on Dempsey Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|-----------------|---------|-------------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 5.1 (Headgate) | 2010 | NO | FISH | - | - | - | - |
| RM 4.8 (Headgate) | 2010 | LL | 85 | 85 | 173 | 73-356 | 96 |
| | | EB | 4 | 4 | 153 | 136-171 | 4 |
| | | SL COT | R | - | - | - | - |
| RM 4.7 (Headgate) | 2010 | NO | FISH | - | - | - | - |
| RM 4.3 (Headgate) | 2010 | NO | FISH | - | - | - | - |
| RM 2.0 (Head) | 2011 | RS SH | R | - | - | - | - |
| RM 1.0 (Headgate) | 2011 | EB | 1 | 1 | 280 | 280 | 100 |
| | | LN SU | R | - | - | - | - |
| | | RS SH | R | - | - | - | - |

In September 2010, discharge measurements were collected by Montana TU technicians on the same date ditches were electrofished. On September 10, the ditch at RM 5.1 was receiving 0.8 cfs, approximately 8% of the mainstem discharge of 9.5 cfs. The ditches at RM's 4.8 and 4.7 were not receiving any water, and the ditch at RM 4.3 was receiving 10.5 cfs, approximately 56% of the mainstem discharge of 18.6 cfs.

In 2011, discharge measurements were collected by Montana TU on the same day electrofishing surveys were performed. On August 10, the ditch at RM 2.0 was receiving 5.8 cfs, approximately 46% of the mainstem discharge of 12.6 cfs.

Racetrack Creek

Racetrack Creek is a tributary to the upper Clark Fork River and drains for over 23 miles before the entering the Clark Fork River at RM 326.5 (Figure 1). Land ownership along the lower 12 miles of Racetrack Creek is primarily private, and land use is agriculture. The remainder of the stream flows through U.S. Forest Service administered lands that dominate the upper portion of the watershed. Numerous mountain lakes are present in the headwaters of Racetrack Creek and its tributaries. Several of the larger lakes have dams that provide storage for downstream irrigators. The primary land uses in the lower portion of the drainage are hay and crop production, cattle grazing, and rural residences. Land use in the upper portion of the watershed is dominated by motorized and non-

motorized recreation on public lands. Fish populations in the upper reaches of Racetrack Creek are comprised of *Oncorhynchus* species including individuals possessing the phenotypic appearance of westslope cutthroat trout, rainbow trout, and hybrids of the two (Lindstrom et al. 2008). In middle reaches of the creek, the fish community is comprised of brown trout, brook trout, and *Oncorhynchus* species. In the lower reaches, irrigation withdrawals and low summer flows can de-water the channel.

Electrofishing surveys were completed in three ditches on Racetrack Creek in September 2010 (Figure 1). The headgates of two ditches were accessible on State of Montana land at approximately RM 7.4 (Figure 1). The third ditch is located downstream on private property at RM 3.5. The upper ditch at RM 7.5, located just upstream of the lower ditch, is fitted with a metal slide gate and a boulder diversion. The diversion is not a barrier to fish passage. Two westslope cutthroat trout and 15 brown trout were collected in the first 100 m below the headgate. Slimy sculpin were also present but rare in this section (Table 20, Appendix A). The second ditch at RM 7.4 is fitted with a metal slide gate and a gravel berm diversion. This diversion also does not appear to be a fish passage barrier. Twenty brown trout and two unidentifiable *Oncorhynchus* species were collected in the first 100 m below the headgate. Slimy sculpin were also present, but rare (Table 20, Appendix A). At the time of these surveys, the discharge of Racetrack Creek was high due to recent rainfall, and resulted in high velocities and turbidity in the ditches. The lower ditch at RM 3.0 (named RM 3.5 in 2010), was also fitted with a metal slide gate and a gravel berm diversion, reinforced with concrete and Visqueen. A total of 40 trout were collected below the headgate, 39 brown trout and one brook trout. Slimy sculpin were also present, but rare in the section (Table 20, Appendix A).

In late July 2011, electrofishing surveys were completed in three ditches previously unsurveyed and one ditch surveyed in 2010. All four were located on lower Racetrack Creek at RM's 3.5, 3.0, 2.6 and 2.0. The ditch at RM 3.5 is fitted with a wooden sliding gate and a gravel berm spanning the channel (Appendix B). The berm does not serve as a fish passage barrier. Ten brown trout were captured and additional individuals were observed in a 100 m section below the headgate (Table 20). Slimy sculpin were present, but rare. The ditch at RM 3.0 had been surveyed in 2010, but the ditch was receiving more water in 2011 due to recent repairs to the diversion (Appendix B). Due to the recent construction the diversion may inhibit migration at lower discharges. Although one brown trout was collected in a 100 m section below the headgate high velocities made it difficult to effectively capture fish (Table 20). One large additional brown trout (>16") was observed but not captured due to these conditions. The ditch at RM 2.6 is fitted with a large metal slide gate and water was diverted from Racetrack Creek with a recently constructed barb of rock and sod (Appendix B). The diversion does not serve as a fish passage barrier. A large volume of sediment was observed in front of the headgate, likely the result of spring runoff. In the first 100 m below the headgate, 24 brown trout and 1 brook trout were captured (Table 20). The ditch at RM 2.0 did not have a formal headgate and water was diverted from Racetrack Creek with temporary diversion of timber and sheet metal. The diversion does not likely inhibit migration, but the general dewatering of the section may be a concern during the irrigation season. Five small brown trout (100-155mm) were collected in a 100 m section of the ditch (Table 20).

Table 20. Electrofishing data collected during 2010 and 2011 in ditches on Racetrack Creek. Ditch RM 3.0 was renamed from RM 3.5 in 2010 (marked with an asterisk).

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|----------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 7.5 (Headgate) | 2010 | WCT | 2 | 2 | 128 | 101-155 | 12 |
| | | LL | 15 | 15 | 73 | 43-204 | 88 |
| | | SL COT | R | - | - | - | - |
| RM 7.4 (Headgate) | 2010 | LL | 20 | 20 | 97 | 43-321 | 91 |
| | | ONC | 2 | 2 | 227 | 222-231 | 9 |
| | | SL COT | R | - | - | - | - |
| RM 3.5 (Headgate) | 2011 | LL | 10 | 10 | 151 | 106-262 | 100 |
| | | SL COT | R | - | - | - | - |
| RM 3.0* (Headgate) | 2010 | LL | 39 | 39 | 127 | 70-197 | 98 |
| | | EB | 1 | 1 | 173 | 173 | 2 |
| | | SL COT | R | - | - | - | - |
| RM 3.0 (Headgate) | 2011 | LL | 1 | 1 | 139 | 139 | 100 |
| RM 2.6 (Headgate) | 2011 | LL | 24 | 24 | 66 | 40-149 | 96 |
| | | EB | 1 | 1 | 75 | 75 | 4 |
| RM 2.0 (Headgate) | 2011 | LL | 5 | 5 | 125 | 100-155 | 100 |

In September 2010, discharges were collected by Montana TU while the ditches were surveyed. On September 9th, the upper ditch at RM 7.5 was receiving 8.1 cfs, approximately 8% of the mainstem discharge of 100.1 cfs, while the lower ditch was receiving 7.2 cfs, or 8% of the remaining mainstem discharge estimated at 92 cfs. The same date, the ditch at RM 3.0 (name changed from RM 3.5) was receiving 3.3 cfs, approximately 33% of the mainstem discharge of 10.2 cfs. Once again, at the time of the 2010 surveys, the discharge of Racetrack Creek was abnormally high due to recent rainfall.

In 2011, Montana TU also measured discharges in most of the ditches surveyed on Racetrack Creek. On July 27th, the ditch at RM 3.5 was receiving 4.7 cfs, approximately 22% of the mainstem discharge. The ditch at RM 3.0 was receiving 8.3 cfs, approximately 39% of the mainstem discharge estimated at 21.3 cfs. The ditch at RM 2.6 was receiving 3.8 cfs, approximately 29% of the remaining mainstem discharge estimated at 12.9 cfs.

Lost Creek

Lost Creek is a tributary to the Clark Fork River and drains for approximately 23 miles before reaching the Clark Fork River at RM 334.7 (Figure 1). The lower 16 miles of the stream flows primarily across private land, while the upper portion of the drainage flows on State of Montana and National Forest land. Also, Lost Creek State Park is situated along the creek from RM 16.2 to RM 18.2. Land use in the upper portion of the watershed is mostly non-motorized recreation, while cattle grazing, rural residences, and irrigated hay production are primary land uses in the lower portion of the drainage. Lost Creek is a principal source of irrigation water for adjacent hay production, and withdrawals often diminish summer flows in several reaches (e.g. upstream of the Galen Highway crossing, and near Interstate 90). A small irrigation impoundment, known as Dutchman Pond, is located on Lost Creek at RM 7.7, and the Gardiner Ditch (originating on Warm Springs Creek) bisects the stream near RM 10.3. Both locations are probable upstream fish passage barriers. Additionally, a natural waterfall located in Lost Creek State Park at RM 17.6 is a likely barrier to upstream movement. Fish populations vary from the upper reaches of Lost Creek to its mouth. Above the natural barrier the trout community is comprised of brook trout and westslope cutthroat, which show heavy hybridization with Yellowstone cutthroat trout (Liermann et al. 2009). In the middle reaches of the stream below the falls, brook trout, westslope cutthroat trout, and brown trout are all present, while in the lowest reaches of Lost Creek, brown trout dominate the trout community.

In August 2010, electrofishing surveys were completed in two ditches located on Lost Creek at RM 9.2 and 7.6 (Figure 1). The ditch at RM 9.2 is fitted with a metal slide gate and a pin and plank diversion spanning the channel. A Denil fish ladder is also in place along-side the diversion. Two sections of this ditch were sampled, one 25 m below the headgate and one approximately 1.6 mi below the headgate. One adult brown trout was the only fish collected just below the headgate and no fish were collected 1.0 mi below (Table 21). The ditch at RM 7.6 is fitted with two metal slide gates and draws water from Dutchman Pond. In the first 100 m below the headgate seven brown trout and nine brook trout were collected, but in the section approximately 0.8 mi below, only one brown trout was present (Table 21, Appendix A).

In early August 2011, electrofishing surveys were completed in three additional ditches on lower Lost Creek located at RM's 4.5, 4.0 and 1.0 (Figure 1). The ditch at RM 4.5 is fitted with a metal slide gate, but has no formal diversion structure (Appendix B). Twenty-one brown trout, and numerous reidside shiner and slimy sculpin were collected in the first 100 m below the headgate (Table 21). The next diversion downstream at RM 4.0 is fitted with a wooden gate and a temporary diversion constructed of lumber, Visqueen and rock (Appendix B). At the time of the survey the headgate was in great disrepair and it would not have been possible to regulate the volume of water diverted. The diversion however, was recently reinforced and nearly dewatered Lost Creek. At the time of the survey the diversion was a fish passage barrier, but may not be in place outside the irrigation season. Forty brown trout and numerous slimy sculpin were collected in the first 100 m below the headgate of the ditch (Table 21). The furthest

downstream ditch at RM 1.0 is fitted with a metal slide gate and receives water from an outside bend of Lost Creek without a formal diversion (Appendix B). Similar to the upstream diversions brown trout were the only trout collected in the ditch, but native longnose suckers were also present. A total of thirty-five brown trout and six longnose suckers were collected in the 100 m below the headgate (Table 21).

Table 21. Electrofishing data collected during 2010 and 2011 in ditches on Lost Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|-----------------|---------|-------------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 9.2 (Headgate) | 2010 | LL | 1 | 1 | 191 | 191 | 100 |
| RM 9.2 (1.6 mi) | 2010 | NO | FISH | - | - | - | - |
| RM 7.6 (Headgate) | 2010 | LL | 7 | 7 | 214 | 90-418 | 44 |
| | | EB | 9 | 9 | 197 | 128-283 | 56 |
| RM 7.6 (0.8 mi) | 2010 | LL | 1 | 1 | 181 | 181 | 100 |
| RM 4.5 (Headgate) | 2011 | LL | 21 | 21 | 84 | 66-160 | 100 |
| | | RS SH | C | - | - | - | - |
| | | SL COT | C | - | - | - | - |
| RM 4.0 (Headgate) | 2011 | LL | 40 | 40 | 109 | 60-265 | 100 |
| | | SL COT | C | - | - | - | - |
| RM 1.0 (Headgate) | 2011 | LL | 35 | 35 | 149 | 66-274 | 100 |
| | | LN SU | R | - | - | - | - |

Discharges were collected by Montana TU in 2010, and on September 1, the upper ditch at RM 9.2 was receiving 5.6 cfs, approximately 89% of the estimated mainstem discharge of 6.3 cfs. No discharge measurements were collected in the ditch at RM 7.6.

On August 3, 2011 Montana TU measured discharges associated with diversions at RM(s) 4.5, 4.0, and 1.0. The ditch at RM 4.5 was receiving 8.6 cfs, approximately 45% of the mainstem discharge estimated at 19.1 cfs. The ditch at RM 4.0 was receiving less than 1.5 cfs, and the ditch at RM 1.0 was receiving 2.5 cfs.

Warm Springs Creek

Warm Springs Creek is a large headwater tributary to the Clark Fork River and drains for approximately 32 miles before reaching the Clark Fork River at RM 339.4 near Warm

Springs, MT (Figure 1). While most of the lower reaches of Warm Springs Creek flow through private land, including the town of Anaconda, several State of Montana Wildlife Management Areas are also located along the stream. Warm Springs Creek transitions to lands managed by the USFS around RM 22.9, along with several private in-holdings. Land uses in the watershed are varied, and include rural and urban residences, cattle grazing, timber harvest, recreation, and historic mining. The trout community throughout much of upper Warm Springs Creek (above Myers Dam) is comprised largely of westslope cutthroat trout and bull trout, with brook trout, rainbow trout and the occasional brown trout also present (Lindstrom et al. 2008). In the lower reaches of Warm Springs Creek (below Myers Dam), brown trout dominate the trout community with *Oncorhynchus* species and brook trout also occurring in lower numbers.

In 2010, electrofishing surveys were completed in three sections of a ditch at RM 10.0, on Warm Springs Creek (Figure 1). This ditch, which is known as the Gardiner Ditch, is fitted with a large metal slide gate and a pin and plank diversion. The planks were removed at the time of the survey and it was not possible to determine if the diversion is a barrier when in use. In the 100 m section, beginning 175 m below the headgate, one bull trout and two brown trout were collected, along with several slimy sculpin (Table 22). Approximately 2.0 mi downstream of the headgate, 27 brown trout were collected in the survey section along with a few slimy sculpin (Table 22, Appendix A). This section was located just below an intersection with Lost Creek and it is uncertain where fish originated (either Warm Springs Creek or Lost Creek). The diversion on Lost Creek at this intersection is a prominent barrier to fish migration up Lost Creek. The third 100 m section was located approximately 4.9 mi below the headgate. Five brown trout were collected in this reach (Table 22, Appendix A).

One additional diversion on lower Warm Springs Creek was surveyed in 2011. The diversion, located at RM 1.8, provides water to the Waterfowl Management Area (WMA) administered by FWP. The ditch receives water through a vertical perforated culvert and no formal diversion is in place (Appendix B). A total of 48 brown trout were captured below the headgate of the WMA ditch, along with numerous slimy sculpin and a few longnose suckers (Table 22, Appendix A).

Table 22. Electrofishing data collected during 2010 and 2011 in two ditches on Warm Springs Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 10.0 (175 m) | 2010 | BULL | 1 | 1 | 282 | 282 | 33 |
| | | LL | 2 | 2 | 107 | 71-143 | 67 |
| | | SL COT | R | - | - | - | - |
| RM 10.0 (2.0 mi) | 2010 | LL | 27 | 27 | 133 | 54-240 | 100 |
| | | SL COT | R | - | - | - | - |

Table 22 (continued). Electrofishing data collected during 2010 and 2011 in two ditches on Warm Springs Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|-----------------|---------|-------------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 10.0 (4.9 mi) | 2010 | LL | 5 | 5 | 230 | 159-280 | 100 |
| RM 1.8 (Headgate) | 2011 | LL | 48 | 48 | 121 | 52-352 | 100 |
| | | LN SU | R | - | - | - | - |
| | | SL COT | C | - | - | - | - |

Discharges related to the ditch at RM 10.0 were not measured in 2010, but visual estimates at the time of the survey suggest the ditch was receiving 15-25% of the mainstem discharge.

Storm Lake Creek

Bull trout populations in the upper Warm Springs Creek drainage are fragmented by several irrigation diversions and impassable structures. One of the more prominent migration barriers is located at the downstream terminus of Storm Lake Creek. At this location a water control and bypass structure diverts Storm Lake Creek to Silver Lake with the option of bypassing water into a canal leading directly to Warm Springs Creek. The design of this structure has produced a vertical fish barrier that blocks virtually all upstream migration (Appendix B). Barriers such as these are a concern as they inhibit individuals from returning to their natal habitats for spawning and can constrain expression of a migratory life history (DeHaan et al. 2010).

In fall of 2010, electrofishing surveys were completed below the Storm Lake Creek diversion structure to evaluate whether bull trout known to occur in Silver Lake were gathering below the structure in an attempt to move up into Storm Lake Creek to spawn. Fishes were collected with a backpack electrofisher in the 80 m canal section between the structure and Silver Lake. All bull trout collected were measured, weighed, given a unique fin clip for the date, and phenotypically identified as either a pure bull trout or a bull trout/brook trout hybrid. Fish identified as pure bull trout were then manually passed over the barrier into Storm Lake Creek. All other fish collected at the site were released below the structure. On ten occasions, over a month long period, a total of seventeen bull trout ranging from 192-570 mm were collected below the structure and moved over (Table 23, Appendix A).

Table 23. Length (mm) and weight (g) of bull trout captured downstream of the water control structure barrier on Storm Lake Creek in the fall of 2010.

| Month | Day | Length | Weight | Comment |
|-----------|-----|--------|--------|-------------------------|
| August | 5 | 437 | 698 | Hook found in throat |
| | 16 | 461 | 887 | |
| | 16 | 482 | 903 | |
| | 16 | 487 | 1075 | |
| | 16 | 529 | 1415 | |
| | 16 | 539 | 1330 | |
| | 18 | 447 | 859 | |
| | 18 | 473 | 947 | |
| | 18 | 495 | 1117 | |
| | 19 | 447 | 808 | |
| | 23 | 436 | 688 | Recapture from 8/5 |
| | 23 | 491 | 1010 | |
| | 23 | 547 | 1350 | |
| | 23 | 570 | 1613 | |
| | 25 | - | - | No bull trout collected |
| | 30 | 539 | 1276 | Recapture from 8/16 |
| September | 2 | 192 | 62 | |
| | 2 | 307 | 246 | |
| | 2 | 528 | 1241 | |
| | 9 | 424 | 676 | |
| | 9 | 530 | 1375 | Recapture from 8/16 |
| | 14 | - | - | No bull trout collected |

In 2011, electrofishing surveys were again completed below the Storm Lake Creek diversion structure using similar techniques described above. However, in this year, each adult fish phenotypically identified as a bull trout was surgically implanted with a passive integrated transponder (PIT) tag. A PIT reader station was installed in the stream channel approximately 0.8 miles upstream of the diversion structure. The objective of this was to determine if fish passed at the diversion were moving upstream. On eleven occasions, between mid-June and early October, a total of eight bull trout ranging from 283-540 mm were collected below the structure and moved over (Table 24). Of these fish, four were recaptured below the diversion one or more times. This suggests that fish frequently fall back over the drop structure after being released. Nevertheless, at least three of the eight bull trout tagged with a PIT tag passed through the PIT reader station 0.8 miles upstream. This work has shown that there is a definite need to provide fish passage at this structure to allow Silver Lake bull trout to return to Storm Lake Creek to spawn.

Table 24. Length (mm) and weight (g) of bull trout captured downstream of the water control structure barrier on Storm Lake Creek in the summer and fall of 2011.

| Month | Day | Length | Weight | PIT Tag # | Comment |
|-------|------|--------|--------|-----------|---|
| June | 14 | - | - | - | No bull trout collected |
| | 29 | 186 | 49 | n/a | Sub-adult; did not pass |
| Aug | 9 | 283 | 205 | 165717283 | |
| | 11 | 479 | 943 | 165717280 | Through PIT Station Aug. 22 @ 16:57 |
| | 15 | 515 | 1266 | 165717286 | |
| | 15 | 540 | 1386 | 165717284 | Hooking scar |
| | 15 | 502 | 1122 | 165717288 | Spinal deformity |
| | 15 | 500 | 1072 | 165717281 | Hooking scar |
| | 22 | 517 | 1058 | 165717285 | Hooking scar, fresh wounds |
| | 22 | 254 | 146 | n/a | Hybrid – did not pass |
| | 25 | 501 | 1094 | 165717288 | Recapture from Aug. 15 |
| | 25 | 487 | 898 | 165717290 | Hooking scar w/ opercula damage |
| | Sept | 1 | 489 | 165717290 | Recapture from Aug. 25; Through PIT Station on Sept. 17 @ 17:40 |
| | | 1 | 499 | 165717288 | Recapture from Aug. 25; Through PIT Station Sept. 13 @ 22:44 |
| | | 6 | - | - | No bull trout collected |
| Oct | 12 | - | - | - | No bull trout collected |
| | 3 | 539 | n/a | 165717284 | Recapture from Aug. 15; Ripe male |
| | 3 | 508 | n/a | 165717286 | Recapture from Aug. 15; Ripe female |

Mill Creek

Mill Creek is located within the Silver Bow Creek watershed, but a bypass channel (the Mill-Willow Bypass) allows Mill and Willow Creeks to circumvent the Warm Springs settling ponds and Silver Bow Creek, which are contaminated with mining wastes. In the upper portion of the Mill Creek drainage dominant land uses are grazing and timber harvest. Towards the middle of the drainage there are a number of permanent and recreational residences located near the stream. The lower portion of the stream flows through lands owned by Atlantic Richfield (ARCO) and much of this land has, or is going through active remediation to promote vegetation growth on soils previously contaminated from past copper smelting activities at nearby Anaconda (Liermann et al. 2009). Fish surveys in 2008 found fish the community in upper sections to be comprised entirely of westslope cutthroat trout and below a waterfall at RM 11.0, the fish community begins to shift towards brown trout dominated (Liermann et al. 2009). In addition to westslope cutthroat trout and brown trout, brook trout, longnose suckers and slimy sculpin have been collected in Mill Creek.

In early August 2011, electrofishing surveys were completed in two ditches diverting water from Mill Creek (Figure 1). The upper ditch at RM 5.8 is fitted with a wooden gate regulated by the installation of wooden planks, and no formal diversion is in place (Appendix B). In the first 100 m below the headgate, six brown trout were collected, along with longnose suckers and slimy sculpin (Table 25). The lower ditch at RM 5.1 is

also fitted with a wooden gate, but water is diverted into the ditch by a gravel/boulder berm. The diversion is not a barrier to fish passage. In the first 100 m below the headgate, three brown trout were collected and slimy sculpin were numerous (Table 25).

Table 25. Electrofishing data collected during 2011 in two ditches on Mill Creek.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|----------------------|--------------|---------|-------------------------|-----------------------|------------------|-------------------|------------------|
| RM 5.8 (Headgate) | 2011 | LL | 6 | 6 | 105 | 55-120 | 100 |
| | | LN SU | R | - | - | - | - |
| | | SL COT | R | - | - | - | - |
| RM 5.1 (Headgate) | 2011 | LL | 3 | 3 | 151 | 127-199 | 100 |
| | | SL COT | C | | | | |

In 2011, Montana TU measured discharges in both ditches surveyed. On July 28, the ditch at RM 5.8 was receiving 3.9 cfs from Mill Creek and the ditch at RM 5.1 was receiving 6.8 cfs.

Browns Gulch

Browns Gulch is the largest tributary to Silver Bow Creek and drains for approximately 18 miles before reaching Silver Bow Creek near Ramsay, MT (Figure 1). Connectivity between Browns Gulch and Silver Bow Creek is relatively good, but can be compromised by irrigation withdrawals in the lower reaches. Land ownership along Browns Gulch is dominated by private land, but some National Forest land is present in the upper extent of the watershed. Land uses in the drainage are primarily irrigated hay production and livestock grazing; however, timber harvest also occurs in the upper extent of the drainage. Irrigation diversions are common throughout the middle and lower reaches of the stream. Fish populations in Browns Gulch vary little throughout the drainage. The trout community is dominated by brook trout, with westslope cutthroat trout also present in low densities (Lindstrom 2011). Longnose sucker and slimy sculpin have also been observed in the lower reaches of Browns Gulch.

In late August 2010, electrofishing surveys were completed in multiple sections on two ditches carrying water from Browns Gulch (Figure 1). These ditches supply water for flood irrigation and center pivots supporting hay production and grazing in the lower portion of the drainage. The upper ditch at RM 4.3, is fitted with a wooden gate regulated by the installation of wooden planks and water is diverted from the mainstem via a pin and plank structure spanning the channel (Appendix B). The diversion is a potential barrier due to the height of the plunge. Two sections were sampled on this ditch, one 105 m below the headgate, and one 0.4 mi below the headgate. Only one species, native long nose sucker, was collected in both sections (Table 26). The lower ditch at RM 4.1 is similarly fitted with a wooden gate and diversion structure. The diversion is in disrepair, but may serve as a barrier at lower discharges. Three sites were sampled on this ditch below the headgate at 20 m, 0.8 mi and 1.3 mi. Similar to the

upper diversion, longnose sucker were present, but rare in the first two sections and no fish were collected 1.5 mi below the headgate (Table 26).

Table 26. Electrofishing data collected during 2010 in two ditches on Browns Gulch.

| Ditch Name (Section) | Year Sampled | Species | Number of Fish Captured | Fish per 100 m (CPUE) | Mean Length (mm) | Length Range (mm) | Species Comp (%) |
|-------------------------|--------------|---------|----------------------------|-----------------------------|------------------------|-------------------------|------------------------|
| RM 4.3 (105 m) | 2010 | LN SU | R | - | - | - | - |
| RM 4.3 (0.4 mi) | 2010 | LN SU | R | - | - | - | - |
| RM 4.1 (20 m) | 2010 | LN SU | R | - | - | - | - |
| RM 4.1 (0.8 mi) | 2010 | LN SU | R | - | - | - | - |
| RM 4.1 (1.3 mi) | 2010 | NO | FISH | - | - | - | - |

Radio Telemetry Results

Based on radio telemetry data from 2009-2011, most of the fish tagged in the Upper Clark Fork River were not affected directly by irrigation structures (Mayfield and McMahon 2010, 2011). However, this does not consider the effects irrigation may have on the river through de-watering. Reduced discharges may lessen the dilution of toxic metals in the river, and increased stream temperatures, may contribute to algal blooms (*Cladophora species*) which lower dissolved oxygen concentrations.

One fish directly impacted by irrigation was a bull trout tagged in 2009. This individual was found dead in mid-June 2009 approximately 0.75 mi below the headgate of the ditch at RM 0.01 on Harvey Creek. The radio-tagged bull trout had been located in the lower portion of Harvey Creek for over a month prior to its expiration. It is believed Harvey Creek was the bull trout's natal stream, and this individual was returning to spawn. However, when the fish reached the impassable barrier at RM 0.2 it remained in the lower portion of the creek until water levels dropped. Falling water levels may have caused the fish to exit the creek and it likely entered the ditch on its way back to the Clark Fork River. Given the extremely low population size of bull trout in the Upper Clark Fork River, any loss of adult bull trout is considered significant.

Another fish potentially impacted by irrigation diversions in 2009, was located in Flint Creek. In early August 2009, a radio-tagged brown trout entered Flint Creek presumably in the early stages of a spawning migration. While the final destination of this fish was Upper Flint Creek near Maxville, MT (RM 15-16) in October, it remained in lower Flint

Creek (RM 3-4) for two months despite low flows and warm stream temperatures in this section of the creek. Several irrigation diversions exist on lower Flint Creek including one adjacent to Hall, MT (RM 5.5). It is possible this fish was unable to navigate past one of these diversions until the numerous diversions withdrawing water from Flint Creek were closed and sufficient flow returned to the mainstem.

In 2010, few radio-tagged fish were directly impacted by irrigation, but one westslope cutthroat trout was observed in an irrigation ditch located on Cottonwood Creek near Deer Lodge, MT. This individual was located alive in the ditch, but the radio tag was later recovered near a heron rookery. This individual may have been captured due to increased vulnerability in the ditch, or scavenged upon expiration.

Again in 2011, few radio-tagged fish were directly impacted by irrigation, but two westslope cutthroat trout may have been lost due to a diversion on Brock Creek. Brock Creek is a small tributary to mainstem Clark Fork River located near Phosphate, MT. Water is diverted into a large culvert approximately one-half mile upstream from the mouth of Brock Creek. The water is used for irrigation in a hay field adjacent to Brock Creek. In the spring of 2011, three cutthroat entered Brock Creek and moved upstream past the culvert to spawn. On their return downstream, two individuals were entrained into the irrigation culvert and eventually perished. The third individual also died, but this likely occurred before the diversion was in operation.

Flint Creek Diversion

The diversion on Flint Creek near Hall, MT diverts water into a small ditch located 10 meters upstream on river right (Appendix B). During the telemetry study, tagged fish were able to navigate upstream past the diversion. However, as mentioned in 2009 one brown trout was documented below the diversion for two months before moving upstream. In 2010, two brown trout radio-tagged on lower Flint Creek moved upstream past the diversion, one in early September and the other in late September. It is unknown whether the planks were in place during September when these individuals passed the diversion, although irrigation is common in the Flint Creek Valley during September. The radio tag from the early fish was recovered near Maxville, MT in early October and the second fish moved into Douglas Creek a small tributary to Flint Creek, likely to spawn. This individual migrated back downstream in mid-December after the planks were removed from diversion. This data suggests that the Hall diversion allows passage seasonally, but additional work would be necessary to determine whether passage is possible when the planks are in place and the diversion is operational.

Although fish entrainment data was not collected at the site, a barrier evaluation was performed on the Hall diversion in 2010. The physical characteristics of the diversion are detailed below.

Location: downstream of the bridge on County Route 512 east of Hall, MT
(46.58592, -113.18121)

Date Surveyed: August 24, 2010

Type of Diversion: Pin and plank with concrete apron
Spillway Type: Over diversion
Diversion Span: 100% of the channel
Estimated amount of water diverted at the time of survey: 15%
Maximum water velocity through the diversion: 9.05 ft/s
Minimum water velocity through the diversion: 4.90 ft/s
Water surface difference (plunge height): 2.5 ft
Maximum plunge pool depth: 2.5 ft
Maximum tailwater control depth: 1.9 ft
Plunge pool distance from outlet: 1.7-2.2 ft
Tailwater control distance from outlet: 17.3-30.0 ft
Bankfull width at inlet: 47.8 ft
Bankfull width at outlet: 62.7 ft
Upstream fish passage structure: No
Fish Screen: No
Lockable headgate: Yes
Flow measuring device: None observed

Entrainment Summary and Site Prioritization

Electrofishing surveys were conducted on 60 irrigation ditches throughout the Upper Clark Fork River Drainage (Figure 2). Valuable data has been collected at all sites and this information may be used to implement management changes in East Fork Rock Creek and Storm Lake Creek.

Irrigation ditches were assigned ratings of high, medium, or low priority for mitigation depending on the species and abundance of fishes entrained (Table 27). A “high” rating indicates greater than 20 westslope cutthroat trout or at least one bull trout were collected in the ditch. A “medium” rating indicates fewer than 20 westslope cutthroat were collected in the ditch or other native fish such as a slimy sculpin or longnose sucker were present. A “low” rating indicates no fish or only non-native species were collected in the ditch. Ratings were adjusted from medium to high if a combination of native salmonids and other native species were collected or if the potential to entrain bull trout was high, such as when bull trout were observed in neighboring ditches.

The entrainment rating for the lower ditch on Ranch Creek (RM 0.1) was adjusted to “high” due to a radio-tagged adult bull trout located in the ditch approximately ten years ago (Brad Liermann, Montana Fish, Wildlife and Parks, personal communication). Entrainment ratings were adjusted from “medium” to “high” for the five ditches surveyed in 2007 on lower Ross Fork Rock Creek. All headgates are located in close proximity to one another and bull trout were found in two of the five. Since these ditches had comparable diversion structures and drew similar volumes of water, it was theorized bull trout would have an equal likelihood of finding their way into any of the ditches. The ditch on lower Harvey Creek (RM 0.01) was adjusted to “high” due to a radio-tagged adult bull trout entrained in the ditch during 2009. The ditch at RM 0.8 on Boulder Creek was also rated as “high” since a bull trout was collected in the ditch just upstream at RM

0.9. Given their proximity and similarities in design, it was sensible to assign consistent ratings. Although westslope cutthroat trout and slimy sculpin were collected in the ditch at RM 0.2 on Boulder Creek, the ditch was assigned a rating of "low" due to great disrepair.

Many of the high priority ditches are located in middle to upper sections of streams where native fish densities tend to be greater (Figure 2). Of the ditches surveyed in Granite County, 14 were designated as high priority (Table 27). These diversions are located on lower Ranch Creek, upper Rock Creek, lower Middle Fork Rock Creek, upper Ross Fork Rock Creek, upper East Fork Rock Creek, Lower Harvey Creek, and lower Boulder Creek. In Powell County, two ditches are designated as high priority, and both are located on lower Gold Creek. In Deer Lodge County, one high priority ditch is located on Warm Springs Creek.

During the inventory several diversions were noted as potential fish passage barriers. Two diversions were observed on Cottonwood Creek, one at the mouth of Cottonwood Creek and one at RM 3.0. Also, two wooden diversions on Dempsey Creek may serve as velocity barriers, one at RM 4.3 and one at RM 4.7. One prominent barrier on Lost Creek is located at the intersection with a ditch from Warm Springs Creek (RM 10.0, Gardiner ditch). This diversion presents a barrier to upstream migration as well as out-migrating fishes. At the intersection water from Warm Springs Creek is diverted down the Lost Creek channel, while Lost Creek water is diverted into the ditch.

In 2010 and 2011, Montana TU inventoried irrigation diversions in the Upper Clark Fork River Drainage. The results of the inventory were compiled in 2011 (TU 2011). See this report for a summary of the diversions encountered and their dimensions. Since diversion names may differ between the reports, a cross-referencing table was included in the appendix of this report (Appendix D).

Table 27. Streams, locations, and entrainment ratings for ditches surveyed in the Upper Clark Fork River Basin from 2007-2011.

| Water Body | County | Location | Latitude | Longitude | Year(s) Surveyed | Rating |
|-------------------------|---------|----------|----------|------------|------------------|--------|
| Ranch Creek | Granite | RM 1.1 | 46.58778 | -113.66974 | 2009, 2010 | Medium |
| Ranch Creek | Granite | RM 0.1 | 46.58982 | -113.67128 | 2009, 2010 | High |
| Stony Creek | Granite | RM 0.3 | 46.34804 | -113.61127 | 2009 | High |
| Beaver Creek | Granite | RM 0.9 | 46.46953 | -113.51035 | 2009 | Medium |
| Rock Creek | Granite | RM 50.6 | 46.24004 | -113.51767 | 2010 | High |
| Rock Creek | Granite | RM 50.5 | 46.24091 | -113.51863 | 2010, 2011 | High |
| Rock Creek | Granite | RM 49.5 | 46.25475 | -113.51905 | 2011 | Medium |
| West Fork Rock Creek | Granite | RM 4.2 | 46.24443 | -113.59107 | 2010 | Medium |
| Ross Fork Rock Creek | Granite | RM 9.4 | 46.14201 | -113.59182 | 2007 | Medium |
| Ross Fork Rock Creek | Granite | RM 6.5 | 46.15901 | -113.56498 | 2007 | High |
| Ross Fork Rock Creek | Granite | RM 6.2 | 46.16042 | -113.56088 | 2007 | High |
| Ross Fork Rock Creek | Granite | RM 6.1 | 46.16096 | -113.56088 | 2007 | High |
| Ross Fork Rock Creek | Granite | RM 6.0 | 46.16124 | -113.56154 | 2007 | High |
| Ross Fork Rock Creek | Granite | RM 5.5 | 46.16489 | -113.56498 | 2007 | High |
| Middle Fork Rock Creek | Granite | RM 0.1 | 46.22779 | -113.51920 | 2010 | High |
| East Fork Rock Creek | Granite | RM 8.6 | 46.13346 | -113.38535 | 2007-2011 | High |
| Harvey Creek | Granite | RM 0.01 | 46.70633 | -113.37312 | 2010, 2011 | High |
| South Fork Lower Willow | Granite | RM 2.9 | 46.52176 | -113.35271 | 2007, 2009 | Medium |
| Gird Creek | Granite | RM 1.5 | 46.49423 | -113.21259 | 2009 | Low |
| Boulder Creek | Granite | RM 0.9 | 46.46613 | -113.23254 | 2009, 2010 | High |
| Boulder Creek | Granite | RM 0.8 | 46.46690 | -113.23292 | 2009, 2010 | High |

Table 27 (continued). Streams, locations, and entrainment ratings for ditches surveyed in the Upper Clark Fork River Basin from 2007-2011.

| Water Body | County | Location | Latitude | Longitude | Year(s) Surveyed | Rating |
|------------------|---------|----------|----------|------------|------------------|--------|
| Boulder Creek | Granite | RM 0.2 | 46.47597 | -113.23605 | 2010 | Low |
| Gold Creek | Powell | RM 6.1 | 46.52590 | -112.97927 | 2010 | Medium |
| Gold Creek | Powell | RM 5.2 | 46.53365 | -112.96513 | 2010 | Medium |
| Gold Creek | Powell | RM 3.9 | 46.53932 | -112.94329 | 2010 | Medium |
| Gold Creek | Powell | RM 3.8 | 46.54031 | -112.94154 | 2010 | Medium |
| Gold Creek | Powell | RM 3.7 | 46.54130 | -112.93914 | 2010 | High |
| Gold Creek | Powell | RM 3.6 | 46.54125 | -112.94197 | 2007, 2009, 2010 | High |
| Gold Creek | Powell | RM 1.1 | 46.57025 | -112.91977 | 2010 | Low |
| Clark Fork | Powell | RM 314.0 | 46.40120 | -112.74256 | 2010 | Medium |
| Cottonwood Creek | Powell | RM 9.0 | 46.36539 | -112.59667 | 2011 | Medium |
| Cottonwood Creek | Powell | RM 6.1 | 46.39970 | -112.64194 | 2009 | Medium |
| Cottonwood Creek | Powell | RM 5.6 | 46.39917 | -112.63923 | 2011 | Medium |
| Cottonwood Creek | Powell | RM 4.0 | 46.39288 | -112.67608 | 2009 | Low |
| Cottonwood Creek | Powell | RM 3.5 | 46.39505 | -112.68085 | 2009 | Low |
| Cottonwood Creek | Powell | RM 3.0 | 46.39981 | -112.69475 | 2009, 2011 | Low |
| Baggs Creek | Powell | RM 0.4 | 46.39676 | -112.63071 | 2011 | Medium |
| Dempsey Creek | Powell | RM 5.1 | 46.29003 | -112.82460 | 2010 | Low |
| Dempsey Creek | Powell | RM 4.8 | 46.28774 | -112.82007 | 2010 | Medium |
| Dempsey Creek | Powell | RM 4.7 | 46.28779 | -112.81874 | 2010 | Low |
| Dempsey Creek | Powell | RM 4.3 | 46.28645 | -112.81055 | 2010 | Low |

Table 27 (continued). Streams, locations, and entrainment ratings for ditches surveyed in the Upper Clark Fork River Basin from 2007-2011.

| Water Body | County | Location | Latitude | Longitude | Year(s) Surveyed | Rating |
|--------------------|------------|----------|----------|------------|------------------|--------|
| Dempsey Creek | Powell | RM 2.0 | 46.30569 | -112.76631 | 2011 | Medium |
| Dempsey Creek | Powell | RM 1.0 | 46.31170 | -112.74957 | 2011 | Medium |
| Racetrack Creek | Powell | RM 7.5 | 46.26722 | -112.86643 | 2010 | Medium |
| Racetrack Creek | Powell | RM 7.4 | 46.26676 | -112.86544 | 2010 | Medium |
| Racetrack Creek | Powell | RM 3.5 | 46.27562 | -112.80367 | 2011 | Medium |
| Racetrack Creek | Powell | RM 3.0* | 46.27701 | -112.79404 | 2010, 2011 | Medium |
| Racetrack Creek | Powell | RM 2.6 | 46.27617 | -112.78625 | 2011 | Low |
| Racetrack Creek | Powell | RM 2.0 | 46.27969 | -112.77071 | 2011 | Low |
| Lost Creek | Deer Lodge | RM 9.2 | 46.17057 | -112.97449 | 2010 | Low |
| Lost Creek | Deer Lodge | RM 7.6 | 46.18234 | -112.85185 | 2010 | Low |
| Lost Creek | Deer Lodge | RM 4.5 | 46.20286 | -112.80575 | 2011 | Medium |
| Lost Creek | Deer Lodge | RM 4.0 | 46.20605 | -112.79917 | 2011 | Medium |
| Lost Creek | Deer Lodge | RM 1.0 | 46.22041 | -112.76888 | 2011 | Medium |
| Warm Springs Creek | Deer Lodge | RM 10.0 | 46.13800 | -112.89300 | 2010 | High |
| Warm Springs Creek | Deer Lodge | RM 1.8 | 46.17200 | -112.80465 | 2011 | Medium |
| Mill Creek | Deer Lodge | RM 5.8 | 46.08991 | -112.90302 | 2011 | Medium |
| Mill Creek | Deer Lodge | RM 5.1 | 46.09534 | -112.89566 | 2011 | Medium |
| Browns Gulch | Deer Lodge | RM 4.3 | 46.03184 | -112.64339 | 2010 | Low |
| Browns Gulch | Deer Lodge | RM 4.1 | 46.02905 | -112.64420 | 2010 | Low |

* River mile designation changed from RM 3.5 (2010 Annual Report) to RM 3.0.

Mitigation Efforts

East Fork Rock Creek

Recent discussion between canal owners/operators (DNRC/Flint Creek Water Users) and fishery biologists may result in actions to mitigate the impacts of East Fork Reservoir operations on fishes in mainstem East Fork Rock Creek below the dam and populations within East Fork Reservoir. The absence of spring scouring flows and the lack of adequate instream flows appears to be altering fish habitat and affecting species composition, as well as abundance of salmonids in East Fork Rock Creek. Also, the entrainment of fish into the canal represents direct losses to fish populations in East Fork Rock Creek below the reservoir. To mitigate these issues, additional instream flows and a spring flushing flow may be provided to the mainstem and a fish screen installed at the headgate of the canal. In 2011, an instream flow study (PHABSIM) was completed by DNRC and other involved agencies to assess the instream flows necessary to provide quality habitat for bull trout and westslope cutthroat trout in East Fork Rock Creek below the dam. The results will likely assist with determining future instream flow requirements for this section of East Fork Rock Creek.

Cottonwood Creek

In the fall of 2011, the privately owned concrete diversion at RM 5.5 was replaced with a passage friendly rock v-weir structure. The project was a coordinated effort among Montana TU and the WRC of the Upper Clark Fork, with assistance also provided by Montana FWP. The project was permitted by the Deer Lodge Valley Conservation District, with input by FWP. Montana TU provided technical assistance and construction oversight, and the WRC contributed funds for material.

Storm Lake Creek


The water control and bypass structure diverting Storm Lake Creek to Silver Lake has recently been shown to impede bull trout migration up Storm Lake Creek. Electrofishing surveys in 2010 and 2011 have shown bull trout congregate below the structure. In 2011, a PIT reader station was installed in Storm Lake Creek approximately 0.8 miles upstream of the diversion structure. The objective of this effort was to determine if fish collected below the structure and passed above were moving upstream. On eleven occasions, between mid-June and early October, a total of eight bull trout ranging from 283-540 mm were collected below the structure and moved over. Of these fish, four were recaptured below the diversion one or more times, suggesting fish frequently go back over the drop structure after being released. Nevertheless, at least three of the eight bull trout tagged with a PIT tag passed through the PIT reader station 0.8 miles upstream. This work has shown that there is a definite need to provide fish passage at this structure to allow Silver Lake bull trout to return to Storm Lake Creek to spawn. The installation of a selective fish passage structure will be pursued at the site, and efforts to collect bull trout below the structure and move them over will continue until this occurs.

Literature Cited


- Bahn, L. 2007. An assessment of losses of native fish to irrigation diversions on selected tributaries of the Bitterroot River, Montana. Montana State University Thesis. Bozeman, MT.
- Behnke, R. 2002. Trout and salmon of North America. The Free Press, New York, NY. 359 pages.
- Christensen, T. 2005. Saving Farmland a Challenge. Western Montana In Business Monthly. Available: www.mtinbusiness.com/inbiz-0508/bus07.php. (2009)
- DeHaan, P., L. Godfrey, D. Peterson, and D. Brewer. 2010. Bull trout population genetic structure and entrainment in Warm Springs Creek, Montana. Abernathy Fish Technology Center, U.S. Fish and Wildlife Service, Abernathy, WA.
- Liermann, B., J. Lindstrom and R. Kreiner. 2009. An assessment of Fish Populations and Riparian Habitat in Tributaries of the Upper Clark Fork River Basin: Phase II. Montana Fish, Wildlife & Parks. Missoula, MT.
- Lindstrom, J., 2011. An Assessment of Fish Populations and Riparian Habitat in Tributaries of the Upper Clark Fork River Basin- 2009 Report. Montana Fish, Wildlife & Parks. Missoula, MT.
- Lindstrom, J., B. Liermann, and R. Kreiner. 2008. An Assessment of Fish Populations and Riparian Habitat in Tributaries of the Upper Clark Fork River Basin. Montana Fish, Wildlife & Parks. Missoula, MT.
- Mayfield, M.P. and T.E. McMahon. 2010. Fisheries restoration potential of the Clark Fork superfund site: mainstem radio telemetry project 2009 annual report. Montana State University. Bozeman, MT.
- Mayfield, M.P. and T.E. McMahon. 2011. Fisheries restoration potential of the Clark Fork superfund site: mainstem radio telemetry project 2010 annual report. Montana State University. Bozeman, MT.
- Montana Department of Natural Resources and Conservation (MDNRC). 2001. Montana stream permitting: a guide for conservation district supervisors and others. Helena, MT.
- TU (Montana Trout Unlimited). 2011. Upper Clark Fork Irrigation Inventory. TU, Missoula, MT.

Appendix A

Ranch Creek




Length distribution of westslope cutthroat trout collected in the ditch at RM 1.1 on Ranch Creek below the headgate in 2009 (n=8).




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.1 on Ranch Creek below the headgate in 2009 (n=5).

Ranch Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 1.1 on Ranch Creek below the headgate in 2010 (n=14).




Length distribution of brown trout collected in the ditch at RM 0.1 on Ranch Creek below the headgate in 2010 (n=10).

Ranch Creek continued...




Length distribution of brook trout collected in the ditch at RM 0.1 on Ranch Creek below the headgate in 2010 (n=8).

Stony Creek




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.3 on Stony Creek below the headgate in 2009 (n=53).

Stony Creek continued...




Length distribution of brown trout collected in the ditch at RM 0.3 on Stony Creek below the headgate in 2009 (n=53).

Beaver Creek




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.9 on Beaver Creek below the headgate in 2009 (n=16).

Beaver Creek continued...




Length distribution of eastern brook trout collected in the ditch at RM 0.9 on Beaver Creek below the headgate in 2009 (n=23).

Rock Creek




Length distribution of brown trout collected in the ditch at RM 50.6 on Rock Creek below the headgate in 2010 (n=5).

Rock Creek continued...




Length distribution of brown trout collected in the ditch at RM 50.5 on Rock Creek below the headgate in 2010 (n=14).




Length distribution of brown trout collected in the ditch at RM 50.5 on Rock Creek below the headgate in 2011 (n=9).

Rock Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 50.5 on Rock Creek below the headgate in 2011 (n=6).




Length distribution of brown trout collected in the ditch at RM 49.5 on Rock Creek below the headgate in 2011 (n=8).

West Fork Rock Creek




Length distribution of unidentifiable *Oncorhynchus* species collected in the ditch at RM 4.2 on West Fork Rock Creek below the headgate in 2010 (n=26).




Length distribution of westslope cutthroat trout collected in the ditch at RM 4.2 on West Fork Rock Creek below the headgate in 2010 (n=11).

Ross Fork Rock Creek




Length distribution of westslope cutthroat trout collected in the ditch at RM 9.4 on Ross Fork Rock Creek below the headgate in 2009 (n=4).




Length distribution of westslope cutthroat trout collected in the ditch at RM 6.5 on Ross Fork Rock Creek below the headgate in 2009 (n=24).

Ross Fork Rock Creek continued...




Length distribution of brown trout collected in the ditch at RM 6.5 on Ross Fork Rock Creek below the headgate in 2009 (n=9).




Length distribution of westslope cutthroat trout collected in the ditch at RM 6.2 on the Ross Fork Rock Creek below the headgate in 2009 (n=22).

Ross Fork Rock Creek continued...




Length distribution of brown trout collected in the ditch at RM 6.2 on the Ross Fork Rock Creek below the headgate in 2009 (n=31).




Length distribution of westslope cutthroat trout collected in the ditch at RM 6.0 on the Ross Fork Rock Creek below the headgate in 2009 (n=16).

Ross Fork Rock Creek continued...




Length distribution of brown trout collected in the ditch at RM 6.0 on the Ross Fork Rock Creek below the headgate in 2009 (n=17).




Length distribution of westslope cutthroat trout collected in the ditch at RM 5.5 on the Ross Fork Rock Creek below the headgate in 2009 (n=7).

Ross Fork Rock Creek continued...




Length distribution of brown trout collected in the ditch at RM 5.5 on the Ross Fork Rock Creek below the headgate in 2009 (n=8).

Middle Fork Rock Creek




Length distribution of brown trout collected in the ditch at RM 0.1 on Middle Fork Rock Creek 0.2 mi below the headgate in 2010 (n=5).

Middle Fork Rock Creek continued...




Length distribution of mountain whitefish collected in the ditch at RM 0.1 on Middle Fork Rock Creek 0.5 mi below the headgate in 2010 (n=6).

East Fork Rock Creek




Length distribution of westslope cutthroat trout collected in Flint Creek Main Canal below the headgate in 2008 (n=22).

East Fork Rock Creek continued...




Length distribution of eastern brook trout collected in Flint Creek Main Canal below the headgate in 2008 (n=176).




Length distribution of westslope cutthroat trout collected in Flint Creek Main Canal below the headgate in 2009 (n=13).

East Fork Rock Creek continued...




Length distribution of eastern brook trout collected in Flint Creek Main Canal below the headgate in 2009 (n=37).




Length distribution of westslope cutthroat trout collected in Flint Creek Main Canal below the headgate in 2010 (n=32).

East Fork Rock Creek continued...




Length distribution of brown trout collected in Flint Creek Main Canal below the headgate in 2010 (n=40).




Length distribution of unidentified *Oncorhynchus* species collected in Flint Creek Main Canal below the headgate in 2010 (n=49).

East Fork Rock Creek continued...




Length distribution of eastern brook trout collected in Flint Creek Main Canal below the headgate in 2010 (n=213).

Harvey Creek




Length distribution of brown trout collected in the ditch at RM 0.01 on Harvey Creek below the headgate in 2010 (n=29).

Harvey Creek continued...




Length distribution of unidentifiable *Oncorhynchus* species collected in the ditch at RM 0.01 on Harvey Creek below the headgate in 2010 (n=29).




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.01 on Harvey Creek below the headgate in 2011 (n=28).

South Fork Lower Willow Creek




Length distribution of westslope cutthroat trout collected in the ditch at RM 2.9 on South Fork Lower Willow Creek below the headgate in 2009 (n=18).

Boulder Creek




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.9 on Boulder Creek below the headgate in 2009 (n=5).

Boulder Creek continued...




Length distribution of brown trout collected in the ditch at RM 0.9 on Boulder Creek below the headgate in 2009 (n=5).




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.8 on Boulder Creek below the headgate in 2009 (n=7).

Boulder Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.9 on Boulder Creek below the headgate in 2010 (n=6).




Length distribution of brown trout collected in the ditch at RM 0.9 on Boulder Creek below the headgate in 2010 (n=9).

Boulder Creek continued...




Length distribution of brown trout collected in the ditch at RM 0.8 on Boulder Creek below the headgate in 2010 (n=39).




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.8 on Boulder Creek below the headgate in 2010 (n=6).

Boulder Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 0.2 on Boulder Creek below the headgate in 2010 (n=6).

Gold Creek




Length distribution of westslope cutthroat trout collected in the ditch at RM 6.1 on Gold Creek below the diversion in 2010 (n=15).

Gold Creek continued...




Length distribution of brown trout collected in the ditch at RM 6.1 on Gold Creek below the diversion in 2010 (n=8).




Length distribution of brown trout collected in the ditch at RM 5.2 on Gold Creek below the diversion in 2010 (n=11).

Gold Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 5.2 on Gold Creek below the diversion in 2010 (n=8).




Length distribution of brown trout collected in the ditch at RM 5.2 on Gold Creek below the diversion in 2010 (n=11).

Gold Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 5.2 on Gold Creek below the diversion in 2010 (n=8).




Length distribution of brown trout collected in the ditch at RM 3.9 on Gold Creek below the diversion in 2010 (n=18). An additional 102 brown trout were collected but not measured individually (ranged 50-70 mm).

Gold Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 3.9 on Gold Creek below the diversion in 2010 (n=16).




Length distribution of unidentified *Oncorhynchus* species collected in the ditch at RM 3.9 on Gold Creek below the diversion in 2010 (n=6).

Gold Creek continued...




Length distribution of brown trout collected in the ditch at RM 3.8 on Gold Creek below the diversion in 2010 (n=24). An additional 134 brown trout were collected but not measured individually (ranged 50-70 mm).




Length distribution of westslope cutthroat trout collected in the ditch at RM 3.8 on Gold Creek below the diversion in 2010 (n=19).

Gold Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 3.7 on Gold Creek below the diversion in 2010 (n=28).




Length distribution of brown trout collected in the ditch at RM 3.7 on Gold Creek below the diversion in 2010 (n=21). An additional 20 brown trout were collected but not measured individually (ranged 50-70 mm).

Gold Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 3.6 on Gold Creek below the headgate in 2007 (n=11).




Length distribution of brown trout collected in the ditch at RM 3.6 on Gold Creek below the headgate in 2007 (n=55).

Gold Creek continued...




Length distribution of brown trout collected in the ditch at RM 3.6 on Gold Creek below the headgate in 2009 (n=30).




Length distribution of brown trout collected in the ditch at RM 3.6 on Gold Creek below the diversion in 2010 (n=67).

Gold Creek continued...




Length distribution of westslope cutthroat trout collected in the ditch at RM 3.6 on Gold Creek below the diversion in 2010 (n=32).




Length distribution of unidentified *Oncorhynchus* species collected in the ditch at RM 3.6 on Gold Creek below the headgate in 2010 (n=7).

Gold Creek continued...




Length distribution of brown trout collected in the ditch at RM 1.1 on Gold Creek below the headgate in 2010 (n=24). An additional 137 brown trout were collected but not measured individually (ranged 40-70 mm).

Clark Fork River




Length distribution of brown trout collected in the Kohrs-Manning ditch below the headgate in 2010 (n=84).

Clark Fork River continued...




Length distribution of brown trout collected in Kohrs-Manning ditch below the intersection with Cottonwood Creek in 2010 (n=20).

Cottonwood Creek




Length distribution of eastern brook trout collected in the ditch at RM 5.6 on Cottonwood Creek below the headgate in 2011 (n=34).

Cottonwood Creek continued...




Length distribution of eastern brook trout collected in the ditch at RM 4.0 on Cottonwood Creek below the headgate in 2009 (n=8).




Length distribution of eastern brook trout collected in the ditch at RM 3.5 on Cottonwood Creek below the headgate in 2009 (n=11).

Cottonwood Creek continued...




Length distribution of eastern brook trout collected in the ditch at RM 3.0 on Cottonwood Creek below the headgate in 2009 (n=8).




Length distribution of eastern brook trout collected in the ditch at RM 2.7 on Cottonwood Creek below the headgate in 2011 (n=6).

Baggs Creek




Length distribution of eastern brook trout collected in the ditch at RM 0.4 on Baggs Creek below the headgate in 2011 (n=12).

Dempsey Creek




Length distribution of brown trout collected in the ditch at RM 4.8 on Dempsey Creek below the headgate in 2010 (n=85).

Racetrack Creek




Length distribution of brown trout collected in the ditch at RM 7.5 on Racetrack Creek below the headgate in 2010 (n=15).




Length distribution of brown trout collected in the ditch at RM 7.4 on Racetrack Creek below the headgate in 2010 (n=20).

Racetrack Creek continued...




Length distribution of brown trout collected in the ditch at RM 3.5 on Racetrack Creek below the headgate in 2011 (n=10).




Length distribution of brown trout collected in the ditch at RM 3.0 on Racetrack Creek below the headgate in 2010 (n=39).

Racetrack Creek continued...




Length distribution of brown trout collected in the ditch at RM 2.6 on Racetrack Creek below the headgate in 2011 (n=24).




Length distribution of brown trout collected in the ditch at RM 2.0 on Racetrack Creek below the headgate in 2011 (n=5).

Lost Creek




Length distribution of eastern brook trout collected in the ditch at RM 9.2 on Lost Creek below the headgate in 2010 (n=9).




Length distribution of brown trout collected in the ditch at RM 9.2 on Lost Creek below the headgate in 2010 (n=7).

Lost Creek continued...




Length distribution of brown trout collected in the ditch at RM 4.5 on Lost Creek below the headgate in 2011 (n=21).




Length distribution of brown trout collected in the ditch at RM 4.0 on Lost Creek below the headgate in 2011 (n=40).

Lost Creek continued...




Length distribution of brown trout collected in the ditch at RM 1.0 on Lost Creek below the headgate in 2011 (n=7).

Warm Springs Creek




Length distribution of brown trout collected in the ditch at RM 10.0 on Warm Springs Creek 2.0 miles below the headgate in 2010 (n=27). Section located below an intersection with Lost Creek, and thus many of the fish collected may have originated from Lost Creek.

Warm Springs Creek continued...




Length distribution of brown trout collected in the ditch at RM 10.0 on Warm Springs Creek 4.9 miles below the headgate in 2010 (n=5). This section was also located below the intersection with Lost Creek




Length distribution of brown trout collected in the ditch at RM 1.8 on Warm Springs Creek below the headgate in 2011 (n=48).

Storm Lake Creek



Length distribution of bull trout collected below water control structure barrier on Storm Lake Creek in 2010 (n=17).


Mill Creek



Length distribution of brown trout collected in the ditch at RM 5.1 on Mill Creek below the headgate in 2011 (n=6).

Appendix B

Ranch Creek



Headgate for ditch at RM 1.1 on Ranch Creek, surveyed in 2009 and 2010.

Stony Creek



Headgate and diversion for ditch at RM 0.3 on Stony Creek, surveyed in 2009.

Beaver Creek



Check dam and headgate for ditch at RM 0.9 on Beaver Creek, surveyed in 2009.

Rock Creek



View of ditch at RM 50.6 on Rock Creek (facing upstream), surveyed in 2010. Headgate similar to headgate for ditch at RM 50.5 (pictured below).

Rock Creek continued...



Diversion (left) for ditch at RM 50.5 on Rock Creek and headgate (right), surveyed in 2010 and 2011.

West Fork Rock Creek



Headgate for ditch at RM 4.2 on West Fork Rock Creek, surveyed in 2010.

Ross Fork Rock Creek



Headgate for ditch at RM 9.4 on Ross Fork Rock Creek, surveyed in 2007.



Diversion and headgate for ditch at RM 6.5 on Ross Fork Rock Creek, surveyed in 2007.

Ross Fork Rock Creek continued...




Diversion for ditch at RM 6.1 on Ross Fork Rock Creek, surveyed in 2009.



Diversion and headgate for ditch at RM 5.5 on Ross Fork Rock Creek, surveyed in 2007.

Middle Fork Rock Creek



Headgate for ditch at RM 0.1 on Middle Fork Rock Creek, surveyed in 2010.

East Fork Rock Creek



Diversion for Flint Creek Main Canal on East Fork Rock Creek, surveyed from 2007-2011.

East Fork Rock Creek



Headgate for Flint Creek Main Canal.



Flint Creek Main Canal directly below headgate (facing downstream).

Harvey Creek



Diversion and headgate for ditch at RM 0.01 on Harvey Creek, surveyed in 2010 and 2011. Clark Fork River Water enters from left above the diversion.

South Fork Lower Willow Creek



Diversion for ditch at RM 2.9 on South Fork Lower Willow Creek, surveyed in 2009.

Gird Creek



Headgate and check-dam (right) for ditch at RM 1.1 ditch on Gird Creek.

Boulder Creek



Headgate for ditch at RM 0.9 ditch on Boulder Creek, surveyed in 2009 and 2010.

Boulder Creek continued...



Headgate for ditch at RM 0.8 on Boulder Creek, surveyed in 2009 and 2010.



Diversion and headgate for ditch at RM 0.2 on Boulder Creek, surveyed in 2010.

Gold Creek



Ditch (right) at RM 6.1 on Gold Creek (left), surveyed in 2010.



Headgate for ditch at RM 5.2 on Gold Creek, surveyed in 2010.

Gold Creek continued...



Parshall flume in ditch at RM 3.9 on Gold Creek, surveyed in 2010.



Headgate for ditch at RM 3.8 on Gold Creek, surveyed in 2010.

Gold Creek continued...



Headgate for ditch at RM 3.7 on Gold Creek, surveyed in 2010.



Headgate for ditch at RM 3.6 on Gold Creek, surveyed in 2010.

Gold Creek continued...



Headgate for ditch at RM 1.1 on Gold Creek (facing upstream),
surveyed in 2010.



View of the fish screen in place on ditch at RM 1.1, surveyed in 2010.

Clark Fork/Cottonwood



Headgate for Kohrs-Manning ditch (RM 314.0) on the Clark Fork, surveyed in 2010.



Cottonwood Creek diversion and secondary headgate (right) on Kohrs-Manning ditch (approximately 0.25 mi below headgate).

Cottonwood Creek



Diversion for ditch at RM 6.0 on Cottonwood Creek, surveyed in 2009.



Headgate for ditch at RM 5.6 on Cottonwood Creek at intersection with Baggs Creek (channel on left), surveyed in 2011.

Cottonwood Creek continued...



Concrete diversion for ditch at RM 5.5 on Cottonwood Creek, surveyed 2009.
This structure was replaced with a rock v-weir structure in fall 2011.



Diversion for ditch at RM 5.4 on Cottonwood Creek, surveyed in 2009.

Cottonwood Creek continued...



Diversion for ditch at RM 3.5 on Cottonwood Creek, surveyed 2009.



Cottonwood Creek dewatered below the diversion at RM 3.5 on June 30, 2009.

Cottonwood Creek continued...



Diversion for ditch at RM 3.0 on Cottonwood Creek in 2009.




Diversion and headgate for ditch at RM 3.0 on Cottonwood Creek in 2011.

Baggs Creek



Headgate for ditch at RM 0.4 on Baggs Creek, surveyed in 2011.

Dempsey Creek



Headgate and diversion for ditch at RM 5.1 on Dempsey Creek, surveyed in 2010.

Dempsey Creek continued...



Headgate for ditch at RM 4.8 on Dempsey Creek, surveyed in 2010.



Check-dam (left) and headgate (right) for ditch at RM 4.7 on Dempsey Creek, surveyed in 2010.

Dempsey Creek continued...



Temporary diversion for ditch (left) at RM 2.0 on Dempsey Creek.



Diversion for ditch (background) at RM 1.0 on Dempsey Creek.

Racetrack Creek



Headgate (left) and diversion (right) for ditch at RM 7.5 on Racetrack Creek, surveyed in 2010.



Headgate for ditch at RM 7.4 on Racetrack Creek, surveyed in 2010.

Racetrack Creek continued...



Headgate for ditch at RM 3.5 on Racetrack Creek, surveyed in 2011.



Headgate for ditch at RM 3.5 on Racetrack Creek, surveyed in 2010 and 2011.

Racetrack Creek continued...



Headgate for ditch at RM 2.6 on Racetrack Creek, surveyed in 2011.




Head (foreground) and diversion (left) for ditch at RM 2.0 on Racetrack Creek, surveyed in 2011.

Lost Creek



Headgate for ditch below Dutchman Pond on Lost Creek (RM 9.2), surveyed in 2010.



Ditch at RM 7.6 on Lost Creek, surveyed in 2010.

Lost Creek continued...



Headgate for ditch at RM 4.5 on Lost Creek, surveyed in 2011.



Head (foreground) and diversion (upper left) for ditch at RM 4.0 on Lost Creek, surveyed in 2011.

Lost Creek continued...



Headgate of ditch at RM 1.0 on Lost Creek, surveyed in 2011.
Photo courtesy of C. Hackathorn.

Warm Springs Creek



Headgate for ditch at RM 10.0 on Warm Springs Creek, surveyed in 2010.

Warm Springs Creek continued...



Culvert entering ditch (FWP WMA) at RM 1.8 on Warm Springs Creek, surveyed in 2011.



View below culvert on ditch at RM 1.8.

Storm Lake Creek




View of the water control structure on Storm Lake Creek. Pipe in the foreground leads to Warm Springs Creek and spill on the right leads to Silver Lake.



View of the vertical barrier on the Storm Lake Creek structure.

Mill Creek



Headgate for ditch at RM 5.8 on Mill Creek, surveyed in 2011.



Headgate and diversion (left) for ditch at RM 5.1 on Mill Creek, surveyed in 2011.

Browns Gulch



Headgate (left) and diversion (right) for ditch at RM 4.3 on Brown's Gulch, surveyed in 2010.



View of ditch at RM 4.1 on Brown's Gulch below headgate, surveyed in 2010.

Flint Creek




Diversion located on Flint Creek near Hall, MT inventoried in 2010. Dimensions of this diversion are presented in the Telemetry Section.




Diversion (left) and headgate (right) of the diversion near Hall, MT.

Appendix C Ross Fork Rock Creek




Maximum daily temperatures recorded by thermographs located at RM 6.7 and RM 0.1 on Ross Fork Rock Creek in 2007.

South Fork Lower Willow Creek




Mean and maximum daily temperatures recorded by a thermograph located in South Fork Lower Willow Creek above the Lower Willow creek reservoir in 2007.

Gold Creek



Maximum daily temperatures recorded by thermographs located at RM 5.7 and RM 0.1 on Gold Creek in 2007.

Cottonwood Creek



Maximum daily temperatures recorded by thermographs located at RM 7.0 and RM 0.5 on Cottonwood Creek in 2007.

Appendix D

Reference table for the Montana Trout Unlimited report (TU 2011).

| Water Body | Survey Year(s) | FWP Report | TU Report | GPS N | GPS W |
|-------------------------|----------------|------------|------------------------|----------|------------|
| Ranch Creek | 2009, 2010 | RM 1.1 | n/a | 46.58778 | -113.66974 |
| Ranch Creek | 2009, 2010 | RM 0.1 | n/a | 46.58982 | -113.67128 |
| Stony Creek | 2009 | RM 0.3 | n/a | 46.34804 | -113.61127 |
| Beaver Creek | 2009 | RM 0.9 | n/a | 46.46953 | -113.51035 |
| Rock Creek | 2010 | RM 50.6 | n/a | 46.24004 | -113.51767 |
| Rock Creek | 2010, 2011 | RM 50.5 | n/a | 46.24091 | -113.51863 |
| Rock Creek | 2011 | RM 49.5 | n/a | 46.25475 | -113.51905 |
| West Fork Rock Creek | 2010 | RM 4.2 | n/a | 46.24443 | -113.59107 |
| Ross Fork Rock Creek | 2007 | RM 9.4 | n/a | 46.14201 | -113.59182 |
| Ross Fork Rock Creek | 2007 | RM 6.5 | n/a | 46.15901 | -113.56498 |
| Ross Fork Rock Creek | 2007 | RM 6.2 | n/a | 46.16042 | -113.56088 |
| Ross Fork Rock Creek | 2007 | RM 6.1 | n/a | 46.16096 | -113.56088 |
| Ross Fork Rock Creek | 2007 | RM 6.0 | n/a | 46.16124 | -113.56154 |
| Ross Fork Rock Creek | 2007 | RM 5.5 | n/a | 46.16489 | -113.56498 |
| Middle Fork Rock Creek | 2010 | RM 0.1 | n/a | 46.22779 | -113.51920 |
| East Fork Rock Creek | 2007-2011 | RM 8.6 | n/a | 46.13346 | -113.38535 |
| Harvey Creek | 2010, 2011 | RM 0.01 | Weaver Ranch Diversion | 46.70633 | -113.37312 |
| South Fork Lower Willow | 2007, 2009 | RM 2.9 | n/a | 46.52176 | -113.35271 |
| Gird Creek | 2009 | RM 1.5 | n/a | 46.49423 | -113.21259 |
| Boulder Creek | 2009, 2010 | RM 0.9 | n/a | 46.46613 | -113.23254 |
| Boulder Creek | 2009, 2010 | RM 0.8 | n/a | 46.46690 | -113.23292 |
| Boulder Creek | 2010 | RM 0.2 | n/a | 46.47597 | -113.23605 |

Reference table (continued) for the Montana Trout Unlimited report (TU 2011).

| Water Body | Survey Year(s) | FWP Report | TU Report | GPS N | GPS W |
|------------------|-----------------------|------------|--------------------------|----------|------------|
| Gold Creek | 2010 | RM 6.1 | Beck 1 | 46.52590 | -112.97927 |
| Gold Creek | 2010 | RM 5.2 | Beck 2 | 46.53365 | -112.96513 |
| Gold Creek | 2010 | RM 3.9 | Menonite 2 Diversion | 46.53932 | -112.94329 |
| Gold Creek | 2010 | RM 3.8 | Menonite 1 Diversion | 46.54031 | -112.94154 |
| Gold Creek | 2010 | RM 3.7 | Wall City 2 Diversion | 46.54130 | -112.93914 |
| Gold Creek | 2007, 2009, 2010 | RM 3.6 | Wall City 1 Diversion | 46.54125 | -112.94197 |
| Gold Creek | 2010 | RM 1.1 | Thomas Diversion | 46.57025 | -112.91977 |
| Clark Fork | 2010 | RM 314.0 | Kohrs Manning Diversion | 46.40120 | -112.74256 |
| Cottonwood Creek | 2011 | RM 9.0 | Applegate 4 Diversion | 46.36539 | -112.59667 |
| Cottonwood Creek | 2009 | RM 6.1 | n/a | 46.39970 | -112.64194 |
| Cottonwood Creek | 2009 (Fish Passage) | RM 6.0 | Dippold/Pryun Diversion | 46.39956 | -112.64265 |
| Cottonwood Creek | 2011 | RM 5.6 | McQueary 1 Diversion | 46.39917 | -112.63923 |
| Cottonwood Creek | 2009 (Fish Passage) | RM 5.5 | Olsen Diversion | 46.39662 | -112.64849 |
| Cottonwood Creek | 2009 (Fish Passage) | RM 5.4 | Smith Diversion | 46.39640 | -112.64888 |
| Cottonwood Creek | 2009 | RM 4.0 | Pryun Diversion | 46.39288 | -112.67608 |
| Cottonwood Creek | 2009 (Fish Passage) | RM 3.5 | Burt Diversion | 46.39511 | -112.68086 |
| Cottonwood Creek | 09 (Fish Passage), 11 | RM 3.0 | Applegate 1 Diversion | 46.39653 | -112.69200 |
| Baggs Creek | 2011 | RM 0.4 | McQueary 3 Diversion | 46.39676 | -112.63071 |
| Dempsey Creek | 2010 | RM 5.1 | Prison Ditch 1 Diversion | 46.29003 | -112.82460 |
| Dempsey Creek | 2010 | RM 4.8 | Prison Ditch 2 Diversion | 46.28774 | -112.82007 |
| Dempsey Creek | 2010 | RM 4.7 | Prison Ditch 3 Diversion | 46.28779 | -112.81874 |
| Dempsey Creek | 2010 | RM 4.3 | Prison Ditch 4 Diversion | 46.28645 | -112.81055 |

Reference table (continued) for the Montana Trout Unlimited report (TU 2011).

| Water Body | Survey Year(s) | FWP Report | TU Report | GPS N | GPS W |
|--------------------|----------------|------------|------------------------------------|----------|------------|
| Dempsey Creek | 2011 | RM 2.0 | Kramer 1 (Barn Box) Diversion | 46.30569 | -112.76631 |
| Dempsey Creek | 2011 | RM 1.0 | Berg Diversion | 46.31170 | -112.74957 |
| Racetrack Creek | 2010 | RM 7.5 | Morrison Ditch Diversion | 46.26722 | -112.86643 |
| Racetrack Creek | 2010 | RM 7.4 | Nicholes Diversion | 46.26676 | -112.86544 |
| Racetrack Creek | 2011 | RM 3.5 | Kelley Diversion | 46.27562 | -112.80367 |
| Racetrack Creek | 2010, 2011 | RM 3.0 | Berg (Box 13) Diversion | 46.27701 | -112.79404 |
| Racetrack Creek | 2011 | RM 2.6 | J-L (Branch) Diversion | 46.27617 | -112.78625 |
| Racetrack Creek | 2011 | RM 2.0 | Johnson (Yellowstone Trail) | 46.27959 | -112.77071 |
| Lost Creek | 2010 | RM 9.2 | Ueland Diversion | 46.17057 | -112.97449 |
| Lost Creek | 2010 | RM 7.6 | Dutchmen Dike Diversion | 46.18234 | -112.85185 |
| Lost Creek | 2011 | RM 4.5 | Beck 2 Diversion | 46.20286 | -112.80575 |
| Lost Creek | 2011 | RM 4.0 | Beck 1 Diversion | 46.20605 | -112.79917 |
| Lost Creek | 2011 | RM 1.0 | Lampert-Hadley Diversion | 46.22041 | -112.76888 |
| Warm Springs Creek | 2010 | RM 10.0 | Gardiner Ditch Diversion | 46.13800 | -112.89300 |
| Warm Springs Creek | 2011 | RM 1.8 | Warm Springs WMA Diversion | 46.17200 | -112.80465 |
| Mill Creek | 2011 | RM 5.8 | Opportunity 4 Diversion | 46.08991 | -112.90302 |
| Mill Creek | 2011 | RM 5.1 | Dakolias 2 Diversion | 46.09534 | -112.89566 |
| Browns Gulch | 2010 | RM 4.3 | Ueland's Reservoir Ditch Diversion | 46.03184 | -112.64339 |
| Browns Gulch | 2010 | RM 4.1 | Ueland Alfalfa Diversion | 46.02905 | -112.64420 |