

MONTANA DEPARTMENT OF FISH, WILDLIFE AND PARKS

FISHERIES DIVISION
JOB PROGRESS REPORT

STATE: MONTANA PROJECT TITLE: STATEWIDE FISHERIES INVESTIGATIONS
PROJECT NO.: F-46-R-7 STUDY TITLE: SURVEY AND INVENTORY OF COLDWATER
AND WARMWATER ECOSYSTEMS
JOB NO.: V-d JOB TITLE: NORTHEAST MONTANA COLDWATER
ECOSYSTEM INVESTIGATIONS
PROJECT PERIOD: JULY 1, 1993 THROUGH JUNE 30, 1994

ABSTRACT

Trout populations in Beaver Creek were not estimated due to poor sampling conditions. Trout gill net catch-per-unit-effort (CPUE) declined dramatically in Beaver Creek Reservoir, primarily due to failure of fall stocked Arlee to recruit. However, average size of trout has increased to an all-time high. A large year-class of northern pike was produced in 1993. Trout gill net CPUE also decreased in Bear Paw Lake due to reduced cutthroat stocking rates and failure of fall stocked Arlee rainbow to recruit. A total of 84,679 suckers, comprising a total biomass of 12.9 tons, have been removed since 1989. Despite this effort, white sucker standing crop is higher now than in pre-removal years. Crayfish standing crop has also been monitored since 1992. Biomass of crayfish decreased from 29 lbs./acre in 1992 to 21 lbs./acre in 1993, but total numbers increased from 21,577 to 25,906. Smallmouth bass were planted in Bear Paw Lake for the second consecutive year in July. Young-of-the-year (YOY) bass began feeding immediately on YOY suckers. By early August, however, YOY suckers had moved to deeper water and were not preyed upon by bass. Yearling bass preferred rocky areas away from sucker concentrations and did not eat suckers. A strong preference for crayfish was observed. Trout and sucker food competition is discussed. Grasshopper Reservoir netting revealed a few large trout are present following chemical rehabilitation. White suckers were found in Faber Reservoir for the first time. Management recommendations for all waters are presented.

OBJECTIVES AND DEGREE OF ATTAINMENT

Streams

1. To ensure within hydrologic constraints that stream flows supporting trout fisheries do not fall below 1975-85 averages. Objective accomplished utilizing state funding.
2. To maintain all of the region's streambanks and channels in their present or improved condition. Objective accomplished utilizing state funding.

3. To maintain water quality at or above 1975-85 average levels. Objective accomplished; monitored compliance with water quality standards and adjusted streamflows in Beaver Creek for temperature moderation.
4. To maintain fish populations and habitat in streams at present levels. Objective accomplished and data presented.
5. To maintain at least 6,000 angler days per year and a trout catch of 0.5 fish per hour. Objective accomplished and data presented.
6. To develop fishing access site acquisition and development for the region. Objective accomplished utilizing state funds.
7. To establish cooperative watershed management plans with federal agencies. Objective accomplished utilizing state funds.
8. To obtain greater public involvement by attending approximately 20 public/sportsmen's club meeting and initiating 2 news releases per year. Objective accomplished utilizing state funding.

Lakes

1. To maintain 70,000 angler days per year and provide catch rates of 0.5 fish per hour or greater. Objective accomplished and data presented.
2. To maintain acceptable trout fishing in waters with nongame and/or predator species. Objective accomplished and data presented.
3. To increase the number and distribution of public fishing waters by acquiring 2 reservoirs every 5 years. Objective accomplished.
4. To obtain public input for management decisions by attending 20 sportsmen's club meetings and providing 3 news releases per year. Objective accomplished utilizing state funding.
5. To develop fishing access site acquisition and development plan for the region. Objective accomplished utilizing state funding.

PROCEDURES

Streams were sampled with a direct current (DC) backpack electrofishing unit. Lakes were sampled with floating and/or sinking experimental gill nets. The gill nets measured 6-feet deep and 125-feet in length and consisted of 25-foot panels of 3/4-, 1-, 1 1/4-, 1 1/2-, and 2-inch square mesh. All fish were measured for total length (TL) and weighed to the nearest .01 pound. Crayfish were captured in 1/4-inch hardware cloth traps of varying size, with throat entrance holes measuring 1.5-2.0 inches in diameter. The traps were set around the entire reservoir in water less than 10 feet deep. Traps were baited with dead fish and canned cat food. Traps were fished for 48 hours. Crayfish were marked with a partial clip of the extreme edge of the right uropod and scattered within several hundred yards of the trapping location. A recapture sample was secured eight

days later using the same technique. Body length measurements for crayfish were taken from the tip of the rostrum to the posterior margin of the telson. Suckers were trapped, for population estimation, in 1/4-inch mesh frame traps. They were weighed, measured and marked with a paper punch hole in their tail.

RESULTS AND DISCUSSION

Beaver Creek

A two-pass electrofishing sample was attempted for Sections 02 and 03 of Beaver Creek to estimate trout numbers. Unusually high stream flows made sampling too inefficient to estimate trout numbers. Despite high flows throughout the year, reproductive success of rainbow and brook trout appeared low in both sections.

Electrofishing conducted in the creek above Bear Paw Lake captured no smallmouth bass which had been planted in the lake in 1992 and 1993. However, two yearling bass were found in the stilling basin below Bear Paw Lake Dam. No adult northern pike or walleye have been found in Beaver Creek above Beaver Creek Reservoir. However, several YOY northern pike were found in a beaver dam about 1/4 mile above Beaver Creek Reservoir in 1982. A YOY walleye was found below Beaver Creek Reservoir in 1990 indicating some downstream escapement occasionally occurs.

Beaver Creek Reservoir

This 200 surface-acre reservoir contains a variety of cold, cool, and warmwater species including rainbow trout, northern pike, walleye, yellow perch, white and longnose suckers. The reservoir has been managed primarily as a trout fishery since its filling in 1974.

Stocking of Eagle Lake and DeSmet rainbow trout commenced in 1985 to provide a longer-lived trout capable of utilizing Beaver Creek for natural reproduction. The domestic Arlee rainbow stocked previously had failed to provide significant natural reproduction, exhibited poor growth and was short-lived. Studies conducted over several years at Beaver Creek Reservoir indicated the Eagle Lake and DeSmet rainbow had increased longevity over Arlee rainbow. Studies also indicated Eagle Lake rainbow were easier to catch than the DeSmet rainbow (Needham and Gilge, 1987). Though Eagle Lake rainbow continue to utilize portions of Beaver Creek for spawning, recruitment to the lake fishery is minimal. Predation on YOY rainbow is suspected; however, recruitment and predation of naturally produced YOY rainbow is as yet undocumented. The stocking of DeSmet rainbow ceased in 1988 and the reservoir was planted exclusively with Eagle Lake rainbow until 1990 when Arlee rainbow were again introduced. Arlee rainbow have been restocked, along with Eagle Lake rainbow, at varying rates and at different times since 1990.

Gill-net surveys were conducted in 1974, 1977, and annually since 1980. These surveys were conducted to monitor growth and survival of hatchery trout and to determine relative abundance of other fishes. Results of these netting efforts are summarized in Table 1. The gill net CPUE of trout decline 77% from 1992, but average size improved considerably and condition remained good. Trout condition is believed to be associated with white sucker numbers, which have declined to all time lows, and should remain low, due primarily to northern pike predation.

Table 1. Summary of gill net catches and relative abundance of fishes in Beaver Creek Reservoir, 1974-93.

Year	Rainbow Trout			Northern Pike			Walleye			Yellow Perch			Sucker Sp. CPUE
	CPUE ¹	Ave. Lgth. (in.)	Ave. Wt. (lbs.)	CPUE	Ave. Lgth. (in.)	Ave. Wt. (lbs.)	CPUE	Ave. Lgth. (in.)	Ave. Wt. (lbs.)	CPUE	Ave. Lgth. (in.)	Ave. Wt. (lbs.)	
1974	24.0	10.7	0.60	48.98	---	---	---	---	---	---	---	---	89.7
1977	35.0	10.1	0.39	37.85	---	---	---	---	---	---	---	---	115.7
1980	23.3	10.1	0.35	33.97	---	---	---	---	---	---	---	---	83.3
1981	7.0	10.4	0.35	31.11	---	---	---	---	---	---	---	---	171.7
1982	8.3	11.2	0.55	37.15	2.3	15.8	0.99	---	---	---	---	---	112.3
1983	3.3	11.8	0.62	37.74	3.7	25.1	4.78	---	---	---	---	---	99.7
1984	3.0	11.3	0.59	40.89	3.7	26.6	5.49	---	---	---	---	---	58.7
1985	3.0	11.9	0.77	45.82	4.3	26.0	5.72	---	---	---	---	---	68.3
1986	13.0	11.9	0.66	39.16	4.2	16.7	2.13	---	---	---	---	---	42.0
1987	11.3	13.6	0.92	36.57	5.2	22.0	2.81	---	---	---	---	---	18.0
1988	9.7	14.7	1.17	36.83	3.0	27.6	7.30	0.7	10.6	0.36	8.2	5.9	18.0
1989	10.7	13.1	0.80	35.59	1.2	30.3	8.31	0.0	---	---	9.2	7.6	16.8
1990	18.5	12.0	0.61	35.30	0.7	21.0	2.90	1.8	13.2	0.86	13.0	8.5	9.8
1991	15.5	12.8	0.77	36.72	2.3	16.6	1.20	5.7	14.0	0.97	12.0	7.4	11.0
1992	13.7	13.7	0.98	38.11	3.3	25.6	5.32	2.3	17.8	2.15	6.0	6.4	7.7
1993	3.2	16.4	1.67	37.86	2.0	27.5	6.37	3.3	16.8	1.73	12.3	7.2	8.5

¹Number of fish caught per gill net.

²Condition factor = $\frac{W \times 10^5}{L^3}$

Arlee rainbow have exhibited consistently higher condition factors than Eagle Lake in this reservoir and the addition of the Arlee strain is suspected as the largest contributing factor to the increase in trout condition. The last time Arlee rainbow were present in the reservoir, the large sucker population presented significant competition. Sucker numbers have since declined to the lowest levels since the reservoir was first filled. Arlee rainbow were reintroduced to determine if they could achieve better growth than Eagle Lake rainbow under less competitive circumstances. It appears that growth performance is much better now than under the competitive circumstances of the past.

Adult northern pike numbers peaked in 1987, but are low at present due to several years of poor reproduction and heavy fishing pressure. The large year classes produced in 1990 and 1991 did not show up in the gill-net catch as expected. The depressed adult pike population is at least partially responsible for the increase in trout numbers in recent years. Northern pike, at present, are rather large, though not particularly numerous. They are still capable of causing significant predation on catchable trout and suckers. Pike predation is expected to increase in the next few years if the recently produced year-classes are as strong as earlier sampling indicated.

Walleye were introduced in 1987 due to local demand and are not considered at this time to be major predators on hatchery trout though the population is increasing in size. Gill net catch-rates increased over 1992, while average size decreased slightly. Gill net catch-rates for yellow perch increased over 1992, as did average size.

Bear Paw Lake

Bear Paw Lake is a 45 surface-acre reservoir on Beaver Creek in the Bear Paw Mountains. It is maintained with annual plants of McBride strain cutthroat and Arlee rainbow trout. In recent years, summer fishing pressure has exceeded 140 angler-days per surface acre. Fishing pressure becomes excessive when acceptable-size fish are readily available. A creel reduction from 10 to 5 fish was imposed in 1987 to distribute the catch under such conditions. Suckers have historically overpopulated this reservoir. The reservoir was chemically rehabilitated in 1983 to reduce sucker numbers. Post-rehabilitation trout growth in 1984 and 1985 was excellent. However, sucker gill net catches increased from a catch per net of 9 in 1985 to 278 per net in 1990 (Figure 1). Despite sucker removal efforts, the population remains high. Competition with large numbers of suckers has reduced growth rates of trout significantly. Fishing pressure has declined dramatically since 1988 due to the small size and poor condition of trout. Respondents to an angler survey indicated a desire to catch larger fish even at the expense of catching fewer fish. The survey results also indicated a preference for cutthroat trout over rainbow trout.

Drawdown and chemical rehabilitation has been attempted several times. This is not a preferred option, however, for sucker control in this reservoir, primarily due to the heavy recreational use, detoxification time, invertebrate loss, bank sloughing, and subsequent trout mortality.

Gill-netting trend data has been collected for a number of years in early September. Utilizing 1988 as a base year, numbers of mature (>10 inches) and juvenile suckers (< 10 inches), as they appeared in the catch, were compared with

trout CPUE and condition (Table 2). The trout gill-net catch decreased 90% from 1992. This appears to be due to reduced stocking rates of Yellowstone cutthroat trout and the inability of fall stocked Arlee rainbow to recruit to the fishery. No rainbow trout were captured in gill nets in 1993. Eagle Lake strain rainbow trout had been utilized experimentally since 1986, but were replaced in 1991 with spring planted Arlee rainbow. The Arlee rainbow were responsible for the large increase in relative plumpness exhibited in 1992. Fall stocking of Arlee rainbow began in 1992 to facilitate hatchery demands.

Bear Paw Lake Gillnet Results




Figure 1. Comparison of white sucker increases, after chemical rehabilitation in 1983, with trout growth in Bear Paw Lake.

A sucker control program was initiated in May of 1989. Frame traps were utilized to capture suckers in the spring as they frequented shorelines and the mouth of Beaver Creek prior to spawning. Trapping effort was increased in 1990, but was reduced in 1992. To date a total of 84,679 suckers with a total biomass of 12.9 tons have been removed from the reservoir (Table 3). Fall gill netting in 1989 indicated the effort had reduced the number of mature suckers significantly. However, by the fall of 1990, juvenile sucker numbers had increased 107% from 1988 levels. Figure 2 illustrates how juvenile suckers increased after removal of adult suckers, and how quickly the sucker population returned to pre-removal levels even while some control pressure remained. It appears that the biomass of suckers remained relatively stable as vast numbers of smaller suckers replaced

the larger suckers that were removed. Based on gill net CPUE, the current sucker population is higher than the preremoval level present in 1988.

Table 2. Relative abundance of suckers and trout as indicated by fall gill netting following sucker removal efforts (1988-93) in Bear Paw Lake.

Date	Suckers				Cutthroat Trout			Rainbow Trout		
	No. <10"		No. >10"		No.		C ²	No.		C ²
	Per	%	Per	%	Per	Avg.		Per	Avg.	
	Net	Chg ¹	Net	Chg	Net	Lgth.		Net	Lgth.	
Fall 1988	122	----	89	---	8.7	7.9	36.50	9.0	10.3	41.18
Fall 1989	152	+25	21	-76	19.0	8.1	30.11	15.3	8.4	33.74
Fall 1990	253	+107	25	-72	22.3	8.7	34.68	9.0	10.0	32.00
Fall 1991	198	+62	57	-36	15.0	9.1	34.50	4.0	10.2	34.87
Fall 1992	127	+4	84	-6	58.6	9.6	24.87	17.0	9.1	45.12
Fall 1993	200	+64	174	+96	6.0	9.1	33.18	0	----	-----

¹1988 used as base year determining percent change in relative abundance.

²Condition factor - $\frac{W \times 10^5}{L^3}$

Table 3. Number and poundage of white suckers removed from Bear Paw Lake from 1989 to 1993.

Year	Number	Pounds
1989	12,545	8,986
1990	44,622	10,206
1991	18,140	4,733
1992	4,133	828
1993	5,239	1,050
TOTALS	84,679	25,803

Bear Paw Lake sucker removal project




Figure 2. Number of juvenile and adult suckers captured per gill net in each year of the study compared with the pounds of suckers removed each year with decreasing effort.

Mechanical/physical removal of juvenile suckers is extremely difficult. Juvenile suckers are not as susceptible to capture as they do not congregate in the creek to spawn. The topography of the reservoir with its steep sides and submerged woody vegetation does not facilitate frame trapping or seining.

Crayfish have become abundant in recent years and may have an effect on food availability for trout. A study from Newcastle Reservoir in Utah by Hepworth and Duffield concluded that crayfish changed the reservoir ecosystem by altering the food web, thereby reducing energy transfer to rainbow trout. Momot (1978) described the ability of crayfish to dominate the benthic community and prey on amphipods, isopods, chironomids, cladocerans, ostracods, and odonate naiads. Crayfish have also been reported to eat gastropods (Dean, 1969). Crayfish are utilized, to some degree, as food by trout. However, only trout greater than 12 inches appear able to consume them on a regular basis. A 12-inch trout in Bear Paw Lake is a rarity, so utilization of crayfish by trout is negligible.

A proposal to introduce smallmouth bass for the purpose of inflicting significant predation on juvenile suckers was submitted in 1991. Juvenile bass were expected to feed primarily on YOY suckers, while adult bass were expected to prey heavily on crayfish. Adult smallmouth bass were not expected to consume a significant number of the catchable-size trout stocked annually. Though the primary objective is to improve trout growth, smallmouth bass should make an outstanding addition to the fishery.

A concern that arose during the lengthy environmental review process was that a "no action" alternative or a delay in implementing an action might cause frustrated fishermen to attempt their own predator introduction. A similar situation (sucker over-population) occurred downstream in Beaver Creek Reservoir in 1982. Unknown individual(s) introduced northern pike into this existing trout fishery apparently to control sucker numbers. Gill netting in Bear Paw Lake in 1992 proved that this was indeed the case as a single adult walleye was captured. It is believed that only a few fish were illegally introduced and the probability of successful reproduction in this reservoir is extremely low.

Following an environmental review, 25,000 smallmouth bass fingerlings, averaging 1.3 inches, were introduced into Bear Paw Lake in August of 1992. It was hoped that YOY suckers would suffer immediate depredation losses; however, the bass were too small to effectively feed on YOY suckers. A cool August and September undoubtedly reduced growth rates of bass and over-winter survival of this introductory plant was questionable. Fifty-six trap-days of effort in the spring of 1993 captured no bass. However, electrofishing the shoreline in mid-July captured 23 age I smallmouth bass ranging in length from 2.2-5.1 inches. All but two of the bass were taken from the dam face riprap. Bass preferred rocky substrate over emergent/submerged vegetation, which was plentiful. A total plant of 40,000 1.7-inch bass was made in July and August of 1993.

In order to evaluate changes in the forage base due to bass predation, baseline data on sucker and crayfish populations was gathered prior to the smallmouth bass introduction. Population data will be gathered annually. A mark-recapture estimate was made of both sucker and crayfish populations in both 1992 and 1993. The estimate of age II and older suckers was 72,737 in 1992. Total biomass was 13,992 pounds or 311 pounds per surface acre. The estimate of age II and older suckers in 1993 was 83,328 with a biomass of 21,324 pounds. Sucker biomass increased 163 pounds per acre from 1992 to 1993.

Adult crayfish, ranging in size from 2.6-3.8 inches were estimated at 21,577 with a total biomass of 1,292 pounds or 29 lbs./acre in 1992. Numbers of crayfish increased in 1993 to 25,906 but average size decreased (range 2.5-3.6 in.). A decrease in average size was reflected in a biomass reduction to 21 pounds per acre.

Dietary Studies

Dietary analysis of most fish species in the reservoir was undertaken in 1993. Samples were taken for stomach analysis in February, May, June, July, August, September, and October for suckers and trout. Smallmouth bass were sampled twice in July, once prior to the second year stocking and again 4 days after stocking. From then on, bass were sampled monthly through October.

The object of the studies was to delineate dietary overlap among suckers, trout, and smallmouth bass and determine the extent to which smallmouth bass utilized suckers and crayfish for food.

-Trout and suckers:

Mid-winter gut examination of suckers, rainbow, and cutthroat trout showed that zooplankton was the most common item ingested by all three. Trout had ingested no aquatic insects and selected heavily for cladocerans, while suckers selected mostly for copepods. Larger suckers also utilized chironomid larvae.

Through the summer, trout fed on a variety of zooplankters and insects while suckers primarily fed on zooplankters.

It would appear that most competition occurring between trout and suckers occurs within the zooplankton community. Though few insect larvae appeared in the sucker guts sampled, the sheer numbers of suckers present could consume significant numbers of larvae even at reduced feeding rates.

-Smallmouth bass young-of-the-year (1993 year-class):

Bass were stocked into the upper reservoir in mid-July. This stocking coincided with the presence of large schools of YOY white suckers in shallow water. Stomach content analysis, from fish four days post stocking, showed heavy utilization of YOY suckers by YOY bass. Seventy-eight percent of the bass sampled had ingested suckers. The mean number of suckers ingested per bass was 4.5.

By early August YOY suckers had moved off-shore into deeper water and stomach content analysis of bass indicated no utilization of suckers at that time. Throughout the remainder of the summer and fall, zooplankton (primarily copepods) and mayfly larvae became the principal food items of YOY bass. It appears there is a "window of opportunity" comprising less than a month, when bass of suitable size (>1.5") can utilize YOY white suckers before they move off shore and become unavailable. Bass did not follow the food source to deeper water, but switched to other available items.

-Smallmouth bass - yearling (1992 year-class):

Yearling bass were only found on the riprap of the dam face in 1993. It appeared that they did not venture to other areas of the reservoir to feed. Though YOY white suckers were plentiful in midsummer, they were found primarily in the weedy areas away from the dam and did not seem to overlap territory with yearling bass. With the exception of several YOY bass cannibalized in September, yearling bass did not prey on fish at any time of the year. Crayfish and chironomid larvae were prevalent in yearling bass stomachs throughout the summer until fall when feeding activity declined and mayfly larvae became the item of choice. Bass were expected to utilize the abundant crayfish as the bass matured, but it is apparent that even bass less than 2.5 inches select for crayfish as small as 3mm in length.

Both YOY and yearling bass took advantage of feeding opportunities for terrestrial insects as evidenced by large numbers of flying ants found periodically in stomachs.

Continued monitoring of these populations and food habit analysis of all fish species should assist in evaluating the affect of this experimental introduction and better define dietary overlap. It is uncertain whether continued removal of adult suckers would be of value at this time.

Grasshopper Reservoir

The most recent stocking strategy for this reservoir includes alternate year plants of Arlee and Eagle Lake rainbow trout. This is done to utilize the longevity of Eagle Lake rainbow, along with the growth and catchability characteristics of the Arlee.

Poor trout growth in the past has been associated with high numbers of white suckers in the reservoir. Spring trapping and removal of suckers was conducted from 1988 to 1991. Though the sucker catch rate decreased from 128 lbs./trap-day (TD) in 1988 to 11 lbs./TD in 1990, juvenile sucker numbers continued to increase at an alarming rate. Gill netting in the fall of 1991 revealed a low trout population accompanied by very low water levels. A decision was made to chemically rehabilitate the reservoir with rotenone. The reservoir and the immediate upstream drainage was successfully treated and the reservoir was fallowed over winter. Catchable-size (7 inch) and fingerling (3 inch) Arlee rainbow trout were planted in the spring of 1992.

A floating and a sinking experimental gill net were fished overnight in mid-September of 1992 and 1993. A total of 122 trout were captured in 1992 averaging 10.3 inches and 0.47 pounds. Due to the high trout density, no fish were planted in spring of 1993. Gill netting in the fall of 1993 captured only one trout which was 18.6 inches long and weighed 3.44 pounds. Fingerling Eagle Lake rainbow trout were planted immediately following fall gill netting. Condition factors of trout are the highest ever recorded in this reservoir. No suckers were netted indicating successful rehabilitation with no recent recontamination.

Faber Reservoir

This reservoir is one of the most popular fishing access sites in northcentral Montana. It has been a consistent producer of quality rainbow trout for three decades.

Adult largemouth bass were found in the reservoir in 1990, the result of an illegal introduction. Shoreline seining and electrofishing was conducted in August of 1993 to determine if bass had successfully reproduced. No YOY bass were found; however, numerous YOY white suckers and several adult suckers were captured. This is the first observation of suckers in the reservoir. Two gill nets set the following week captured 25 rainbow trout and 81 adult suckers. Similar gill net sets in 1987 produced 51 trout and no suckers. The suckers ranged in length from 10.6-13.3 inches. Though numerous, the presence of suckers appears to have had little affect on trout growth rates or condition to date. Overall condition of trout remains excellent.

RECOMMENDATIONS

Beaver Creek: Continue to monitor trout populations in all sections. Stock brown trout periodically.

Beaver Creek Reservoir: Continue stocking of Eagle Lake and Arlee rainbow catchable-size trout. Evaluate the Erwin domestic strain as a possible replacement for Arlee due to projected future shortfalls of Arlee rainbow in the hatchery system. Increase stocking rate as northern pike population increases.

Bear Paw Lake: Continue McBride strain cutthroat stocking at reduced rates. Continue Arlee catchable stocking at current rates. Annual population estimates of suckers and crayfish should be made to monitor effects of smallmouth bass introduction. Determine food competition overlay between species and evaluate extent of bass predation on suckers.

Increase number and size of smallmouth bass planted annually and coordinate stocking with YOY sucker schooling for maximum predation effect. Investigate rock placement around reservoir to help redistribute bass.

Grasshopper Reservoir: Continue with annual plants of Arlee fingerlings and alternate year plants of Eagle Lake rainbow. Monitor with annual gill net survey.

Faber Reservoir: Continue to monitor for bass reproduction and monitor growth and condition of trout annually in light of recent sucker infestation.

LITERATURE CITED

- Dean, J. L. 1969. Biology of the crayfish *Orconectes causeyi* and its use for control of aquatic weeds in trout lakes. U.S. Bur. of Sport Fisheries and Wildlife Technical Paper 24.
- Hepworth, Dale K. and Daniel J. Duffield, 1987. Interactions between an exotic crayfish and stocked rainbow trout in Newcastle Reservoir, Utah. No. Amer. Jour. of Fisheries Mgt. 7:554-561.
- Momot, W.T., H. Gowing, and P.D. Jones. 1978. The dynamics of crayfish and their role in ecosystems. American Midland Naturalist 99:10-35.
- Needham, Robert G. and Kent Gilge, 1987. Inventory and Survey of the Project Area, Job Prog. Rept. for D-J Project F-11-4-34, Job. No. I-a. 40pp. (mimeo).

Waters referred to:

15-4570-03 Beaver Creek Reservoir
15-4560-05 Bear Paw Lake
15-5380-07 Grasshopper Reservoir
15-0320-01 Beaver Creek
15-5140-01 Faber Reservoir

Key Words or Fish Species:

Arlee, Eagle Lake rainbow trout, sucker removal, crayfish, population estimates, smallmouth bass, dietary studies.

Prepared by: Kent W. Gilge
Date: June 30, 1994

