

## **Montana Conservation Genetics Laboratory**

Division of Biological Sciences \* University of Montana \* Missoula, MT 59812 (406)243-5503/6749 Fax (406)243-4184

July 20, 2006

Lee Nelson Genetics Contact, Region 3 Mt. Dept. of Fish, Wildlife, and Parks 415 South Front Street Townsend, Montana 59644

Lee:

The paired interspersed nuclear DNA elements (PINE) technique has been used to analyze DNA from the following trout samples:

Summary of	f results.							,
Sample #	Water Name/Location/Collection Date/ Collector	a N	# markers	Species ID	Power (%)	<b>% WCT</b>	e Individuals	t
3300	Muskrat Creek	24	R7Y4	WCT	R99Y99	100	24	

6/21/2006 Lee Nelson

<sup>a</sup>Number of fish successfully analyzed. If combined with a previous sample, the number in parentheses indicates the combined sample size <sup>b</sup>Number of markers analyzed that are diagnostic for the non-native species (R=rainbow trout, W=westslope cutthroat trout, Y=Yellowstone cutthroat trout).

<sup>c</sup>Codes: WCT = westslope cutthroat trout (*Oncorhynchus clarki lewisi*); RBT = rainbow trout (*O. mykiss*); YCT = Yellowstone cutthroat trout (*O. clarki bouvieri*). Only one species code is listed when the entire sample possessed alleles from that species only. However, it must be noted that we cannot definitively rule out the possibility that some or all of the individuals are hybrids. We may not have detected any non-native alleles at the loci examined because of sampling error (see Power %). Species codes separated by "x" indicate hybridization between those species. <sup>d</sup>Number corresponds to the percent chance we have to detect 1% hybridization given the number of individuals successfully analyzed and the number of diagnostic markers used. For example, 25 individuals are required to yield a 95% chance to detect as little as 1% hybridization with Yellowstone cutthroat trout into what once was a westslope cutthroat trout population. Not reported when hybridization is detected.

<sup>•</sup>Indicates the genetic contribution of the hybridizing taxa in the order listed under c to the sample assuming Hardy-Weinburg proportions. This number is reported if the sample appears to have come from a hybrid swarm. That is, a random mating population in which species markers are randomly distributed among individuals.

<sup>f</sup>Indicates number of individuals with genetic characteristics corresponding to the species code column when the sample can be analyzed on the individual level. This occurs when marker alleles are not randomly distributed among individuals and hybridization appears to be recent and/or if the sample appears to consist of a mixture of populations and hybrids and non-hybrids can be reliably distinguished.

## **Methods and Data Analysis**

The PINE technique uses short synthetically made segments of DNA called primers, in pairs, to search for relatively small segments of organismal DNA flanked by particular, often viral, DNA inserts. During the

polymerase chain reaction (PCR), the primers bind to the ends of the inserts and many copies of the organismal DNA between the primers are made. While the DNA from some organisms may have two appropriately spaced inserts to which the primers can attach, the DNA from other organisms may have only one or none of the appropriately spaced inserts in particular regions. During PCR we will fail to copy DNA in the latter two cases. Thus, the PINE technique coupled with PCR is used to search for evidence of genetic variation based on the presence or absence of particular DNA fragments. The fragments are labeled by the primers used to produce them and their length in terms of the number of nucleotides in the fragment.

The fragments are made using dye labeled nucleotides and after PCR are separated from each other via electrophoresis in polyacrylamide gels. Smaller fragments move through the gels at a faster rate than larger fragments. The use of dye labeled nucleotides allows one to visualize the position of the fragments in the gels after electrophoresis using a spectrophotometer and the size of the fragments is determined by comparison to the position of synthetic fragments of known size that were also migrated into the gel.

When DNA from westslope cutthroat trout, *Oncorhynchus clarki lewisi*, and rainbow trout, *O. mykiss*, is compared with PINE analysis and three different pairs of primers seven fragments are usually characteristic of westslope cutthroat trout and seven fragments are usually characteristic of rainbow trout (Table 1). Likewise, when DNA from westslope and Yellowstone cutthroat trout, *O. c. bouvieri*, is compared using the same procedure two fragments are usually characteristic of westslope cutthroat trout and four fragments are usually characteristic of Yellowstone cutthroat trout (Table 1).

Fragments produced from the DNA of one taxon and not another are commonly termed diagnostic or marker loci because they can be used to help determine whether a sample came from a non-hybridized population of one of the taxa or a population in which hybridization between them has or is occurring. Individuals from a non-hybridized population will possess fragments characteristic of only that taxon. In contrast, since half the DNA of first generation hybrids comes from each of the parental taxa the DNA from such individuals will yield all the fragments characteristic of the two parental taxa. In later generation hybrids, the amount and particular regions of DNA acquired from the parental taxa will vary among individuals. Thus, DNA from later generation hybrid individuals will yield only a subset of the parental fragments and the particular subset will vary among individuals.

In a sample from a random mating hybrid swarm, that is a population in which the genetic material (i.e. fragments) of the parental taxa is randomly distributed among individuals such that essentially all of them are of hybrid origin, the frequency of the fragment producing allele from the non-native taxon is expected to be nearly equal among the diagnostic loci since their presence can all be traced to a common origin or origins. Thus, if a sample contains substantial variation at only a single marker locus where the presence of the fragment is usually characteristic of a non-native taxon and lacks such fragments at all other markers this is probably not indicative of hybridization. Rather, it much more likely represents the existence of genetic variation for the presence or absence of the fragment within this particular population of the native taxon.

An important aspect of PINE marker loci is that individuals homozygous for the presence allele (pp) or heterozygous (pa) will both yield the fragment. That is, p is dominant to a. Thus, in order to estimate the genetic contribution of the native taxon to a hybrid swarm we concentrate on the marker loci at which the p allele is characteristic of the non-native taxon. Furthermore, we must assume that genotypic distributions in the population reasonably conform to expected random mating proportions. Under this assumption the frequency of the native a allele is approximately the square root of the frequency of individuals in the population lacking the fragment (aa). The frequency of the non-native allele then is one minus this value. We focus on the p alleles characteristic of the non-native taxon because with low levels of hybridization it is the presence of these alleles that are likely to provide evidence of hybridization. With low levels of hybridization, it is likely all individuals in

the sample will genotypically be *pp* or *pa* where the *p* allele is characteristic of the native taxon. Thus, like in non-hybridized populations all individuals in the sample will yield the fragment providing no evidence of hybridization.

Failure to detect evidence of hybridization in a sample does not necessarily mean the population is nonhybridized because there is always the possibility that we would not detect evidence of hybridization because of sampling error. In order to assess the likelihood the population is non-hybridized, we determine the chances of not detecting as little as a one percent genetic contribution of a non-native taxon to a hybrid swarm. This is simply  $0.99^{2NX}$  where N is the number of fish in the sample and X is the number of marker loci where the *p* allele is characteristic of the non-native taxon.

In samples showing evidence of hybridization, that is; fragments characteristic of a non-native taxon were detected at two or more marker loci, we used two approaches to determine if the population appeared to be a hybrid swarm. First, contingency table chi-square analysis was used to test for heterogeneity of allele frequencies among the marker loci. Next, we computed a hybrid index for each individual in the sample. Each diagnostic locus at which an individual possessed a PINE fragment characteristic of the non-native taxon was given a value of one. Each diagnostic locus at which an individual did not possess a PINE fragment characteristic of the non-native taxon was given a value of zero. These values summed over all diagnostic loci represent an individual's hybrid index. The observed distribution of hybrid index scores was then statistically compared to the expected random binomial distribution based on the estimated native and non-native genetic contributions to the sample. If the allele frequencies were statistically homogeneous among the diagnostic loci and the observed distribution of hybrid indices statistically conformed to the expected random binomial distribution, then the sample was considered to have come from a hybrid swarm.

Heterogeneity of allele frequencies among marker loci can arise in very old hybrid swarms as the frequencies over time diverge from each other due to genetic drift. In this case, however, the non-native fragments will still be randomly distributed among individuals. Thus, samples with these characteristics were also considered to have come from hybrid swarms.

There are two likely reasons why a non-random distribution of non-native fragments may be observed among individuals in a sample. It may contain individuals from genetically divergent populations with different amounts of hybridization or hybridization may have only recently occurred in the population. Based on PINE data alone, these two situations will generally be difficult to distinguish from each other. Regardless of the explanation, when the non-native fragments are not randomly distributed among individuals in a sample estimating a mean level of hybridization has little, if any, biological meaning and, therefore, is often not estimated.

## **Results and Discussion:**

## Muskrat Creek 3300

Diagnostic PINE fragments characteristic of only westslope cutthroat trout were detected in the sample. Previous allozyme (#459, October 1, 1990, N=10) and PINE (#2872, August 27, 2003, N=50; #3271, May 1, 2006, N=22) analyses also found no evidence of hybridization in Muskrat Creek. With the combined sample size of 106, we have greater than a 99% probability of detecting as little as a one percent rainbow or Yellowstone cutthroat trout genetic contribution to a hybrid swarm. The Muskrat Creek population, therefore, is almost certainly non-hybridized westslope cutthroat trout.

Robb Leary John Powell **Table 1:** Diagnostic PINE markers for westslopecutthroat, Yellowstone cutthroat, and rainbow trout. Xindicates the fragment is present in the particulartaxon.

Markers	Yellowstone	Westslope	Rainbow
Hpa1 5'/Hpa1 3'			
232	Х		
153		Х	
110.5			Х
72	Х	Х	
70			Х
69	Х	Х	
66			Х
Fok1 5'/Tc1			
369			X
366	Х	Х	
230			Х
159	Х		
138	Х		
110		Х	
Hpa1 5'/33.6+2			
395			Х
388	Х	Х	
266			Х
248	Х		
148	Х	Х	