

7-12-04
22200 23605
Region 3
Also
F-02 25
RefID: 86691

**INTERIM DATA COMPILATION AND UPDATE FOR STUDY SECTIONS OF THE
BIG HOLE RIVER OF SOUTHWEST MONTANA
2004**

By:

**Richard A. Oswald
and
Scott A. Lula**

June 2004
Montana Department of Fish, Wildlife & Parks
Region Three
1400 S. 19th Avenue
Bozeman, Montana

Executive Summary

Fisheries data from the Big Hole River of southwest Montana were last compiled, presented, and discussed by Oswald (2002a) in a written report submitted in compliance with the Federal Aid in Fish and Wildlife Restoration Acts. The report covered the sampling period from 2000 through 2001 and Project Numbers F-78-R-6 through F-113-R-1. Fisheries data collected during the 2000 - 2002 period for streams within the Beaverhead River Drainage were analyzed and reported in detail by Oswald (2003) and data for area reservoirs and lakes were reported by Oswald (2004).

Data presented in graphic form in this report represent sampling conducted within the Big Hole River study sections in 2003 and 2004 and will be fully discussed in an edited written report in 2005. At this time, the data should be considered as a preliminary addition to existing data sets and will not be fully interpreted until additional data has been gathered and presented in an edited written format.

Stream flow data were extracted from USGS records from the Melrose Gage Station which has been operated on the Big Hole River since 1923. There are many ways to analyze flow data in terms of its affect on aquatic habitats. Figures 1 - 3 in Appendix A utilize several different means of comparison. Annual water yield in acre - feet is compared to the averages for the period of record and for the 1982 - 2002 period presented in Figure 1. Summer mean base flow, as exemplified by the month of August, is compared with the upper and lower minimum flow inflection points in Figure 2 and minimum August flows are compared in Figure 3. The most recent study period has been characterized by low flow regimes in association with exceptional drought conditions which have been prevalent in southwest Montana.

Rainbow trout population data are presented in Appendix B, Figures 4 through 11, for the Jerry Creek and Melrose Study Sections. Population parameters analyzed include density, standing crop, recruitment, numbers of 13 inch and 16 inch and larger fish. Rainbow trout populations, under the current sampling period, have begun to evidence symptoms of decline under low flow regimes associated with drought. In addition to drought related stress, whirling disease has begun to affect recruitment of rainbow trout in the Melrose Study Section. Test cages of juvenile rainbow trout were placed in strategic locations within the Melrose Section reach in 2003 and 2004 to ascertain the frequency and intensity of the infection. It is also hoped that the test cages will be usable as an indicator of the source of the whirling disease infection. Low flow regimes and concern for compounding stress to trout populations at low flow have resulted in limited fall sampling efforts which have excluded the Maiden Rock Section and limited sampling effort in the Jerry Creek and Melrose Study Sections.

Brown trout population data are presented in Appendix C, Figures 12 - 20, for the Maiden Rock, Melrose, and Hog Back Study Sections. Brown trout population parameters presented include density, standing crop, numbers of 13 inch and larger and 18 inch and larger fish. Data, particularly that of the Hog Back Study Section, are symptomatic of the affects of persistent low flow and elevated water temperatures on brown trout populations.

LITERATURE CITED

- Oswald, R.A. 2002a. Inventory and survey of the salmonid populations of the Big Hole River of southwest Montana. Job Prog. Rpt., Fed. Aid in Fish and Wildlife Restoration Acts. Mont. Proj. Nos. F-78-R-6 and F-113-R-1 27pp.
- _____. 2003. Inventory and survey of selected stream fisheries of the Red Rock, Ruby, and Beaverhead Rivers of southwest Montana; 2000 - 2002. Job Prog. Rpt., Fed. Aid in Fish and Wildlife Restoration Acts. Mont. Proj. Nos. F-78-R-6, F-113-R-1, and F-113-R-2 71pp.
- _____. 2004. Inventory and survey of fisheries in lowland lakes and reservoirs of the Red Rock, Ruby, Beaverhead, and Big Hole River drainages of southwest Montana, 2002 - 2004. Job Prog. Rpt., Fed. Aid in Fish and Wildlife Restoration Acts. Mont. Proj. Nos. F-113-R-2, R-3, and R-4 47pp

Report Prepared By: Richard A. Oswald and Scott A. Lula, MFWP, Region 3, Bozeman, June 2004

All Work Included in this Report in Conjunction with Federal Aid in Fish and Wildlife Restoration Acts:
Montana Project Numbers: F-113-R-2, R-3, and R-4

Montana Fish, Wildlife & Parks Project Numbers 3320S and 3320T

APPENDIX A

Figure 1. Annual Water Yield (Acre Feet) for the Big Hole River at the USGS Melrose Gage 1982 - 2003.




Figure 2. Mean August flow as compared with the FWP recommended optimum and minimum instream flow at the USGS Melrose Gage on the Big Hole River; 1982 - 2003.





Figure 3. Minimum August flows (ADF) measured at the USGS Melrose Gage for the 1982 - 2003 period of record on the Big Hole River.



APPENDIX B

Figure 4. Estimated fall density and standing crop of Rainbow Trout in the Jerry Creek Section of the Big Hole River, 1986 - 2003.




Figure 5. Estimated fall density of Age I Rainbow Trout in the Jerry Creek Section of the Big Hole River, 1986 - 2003.




Figure 6. Estimated fall density of 13 inch and larger Rainbow Trout in the Jerry Creek Section of the Big Hole River 1986 - 2003.




Figure 7. Estimated fall density of 16 inch and larger Rainbow Trout in the Jerry Creek Section of the Big Hole River, 1986 - 2003.




Figure 8. Estimated fall density and standing crop of Age I and older rainbow trout in the Melrose Section of the Big Hole River, 1981 - 2003.




Figure 9. Estimated fall density of Age I Rainbow Trout in the Melrose Section of the Big Hole River, 1981 - 2003.




Figure 10. Estimated fall density of 13 inch and larger Rainbow Trout in the Melrose Section of the Big Hole River, 1981 - 2003.





Figure 11. Estimated fall density of 16 inch and larger Rainbow Trout in the Melrose Section of the Big Hole River, 1981 - 2003.



APPENDIX C

Figure 12. Estimated spring density and standing crop of Brown Trout in the Maiden Rock Section of the Big Hole River 1981 - 2002.




Figure 13. Estimated spring density of 13 inch and larger Brown Trout in the Maiden Rock Section of the Big Hole River, 1981 - 2002.




Figure 14. Estimated spring density of 18 inch and larger Brown Trout in the Maiden Rock Section of the Big Hole River, 1981 - 2002.




Figure 15. Estimated spring density and standing crop of Brown Trout in the Melrose Section of the Big Hole River, 1981 - 2003.




Figure 16. Estimated spring density of 13 inch and larger Brown Trout in the Melrose Section of the Big Hole River 1981 - 2003.




Figure 17. Estimated spring density of 18 inch and larger Brown Trout in the Melrose Section of the Big Hole River, 1981 - 2003.




Figure 18. Estimated spring density and standing crop of Brown Trout in the Hog Back Section of the Big Hole River, 1987 - 2003.





Figure 19. Estimated spring density of 13 inch and larger Brown Trout in the Hog Back Section of the Big Hole River, 1987 - 2003.



**Figure 20. Estimated spring density of 18 inch and larger
Brown Trout in the Hog Back Section of the Big Hole
River, 1987 - 2003.**

