

MONTANA DEPARTMENT OF FISH, WILDLIFE AND PARKS
FISHERIES DIVISION

JOB PROGRESS REPORT

STATE: MONTANA PROJECT TITLE: STATEWIDE FISHERIES INVESTIGATIONS
PROJECT NO.: F-46-R-1 STUDY TITLE: SURVEY AND INVENTORY OF COLDWATER
STREAMS
JOB NUMBER: I-d
JOB TITLE: LOWER CLARK FORK RIVER FISHERY INVESTIGATION
PROJECT PERIOD: JULY 1, 1987 THROUGH JUNE 30, 1988

ABSTRACT

A fishery inventory and planning study was continued on the middle Clark Fork River system. Rainbow trout comprise the bulk of the sport fishery along with a few brown, westslope cutthroat and bull trout.

Preliminary estimates in five study sections on the Clark Fork River indicate the river supports from 175 to 402 catchable rainbow trout per mile. Catchable brown, westslope cutthroat and bull trout were present in all study sections, but their numbers were usually too low to estimate. This density of catchable trout is less than expected for comparable trout streams the size of the Clark Fork.

Visual surveys were made in the Milltown, Missoula, Huson and Superior study sections in an attempt to locate trout redds. Brown trout redds were located in the Milltown and Missoula sections. Young-of-the-year trout were relatively scarce in all four study areas. Saturation plants of 10,000 hatchery reared young-of-the-year brown trout were made in the Huson section to aid in evaluating whether recruitment is a limiting factor for trout populations in the Clark Fork River.

OBJECTIVES AND DEGREE OF ATTAINMENT

The long range objective of the study is to follow inventory procedures developed in earlier studies (Wiperman 1973, Berg 1975, 1981 and 1983) and use the resulting data to prepare recommendations for aquatic resource management on this section of the Clark Fork River. Specific objectives during this report period were:

1. Determine species distribution and abundance and relative condition of fish populations in the Clark Fork River and its tributaries.
2. Measure physical trout habitat parameters in the Clark Fork River and its tributaries and evaluate correlations with trout population characteristics.

3. Maintain trout populations and habitat conditions in the lower Clark Fork River and its major tributaries at levels at least as good as present status.
4. Monitor spawning migrations of rainbow, cutthroat, brown and bull trout in tributaries of the Clark Fork River.
5. Monitor outmigrations of juvenile trout from tributaries to the main stem of the Clark Fork River and determine the relative importance of various tributaries in providing recruitment to the trout population in the main river.
6. Evaluate whether recruitment is a limiting factor for trout populations in the Clark Fork River and identify factors which may contribute to the scarcity of a brown trout fishery in the Clark Fork River below Missoula.
7. Correlate parameters identified in water quality studies conducted by DFWP and other agencies with relative abundance of the fishery in the Clark Fork River.
8. Maintain water quality at or above 1984-86 average levels as measured at Montana Department of Health and Environmental Sciences water quality monitoring stations.
9. Determine and maintain adequate instream flow levels in the Clark Fork River and its major tributaries.
10. Define fish movement patterns and relative angler harvest and maintain a trout fishery on the lower Clark Fork River of at least 40,000 man-days per year with an average catch rate of 0.2 fish per hour.

Objectives 3, 7, 8 and 9 are state-funded. Progress was accomplished on all federally funded objectives. Findings are presented in the appropriate sections of this report.

PROCEDURES

Water Temperature

Thirty-day continuous recording thermographs were used to monitor water temperature on the Clark Fork River stations at Milltown Dam and Petty Creek. The recorder box was positioned on the stream bank as far above the high water mark as possible. A thermocouple lead, varying in length from 8 to 23 m, was extended into the water through flexible, plastic sewer pipe. Water temperature data for the St. Regis River, Fish Creek and the Clark Fork River stations at Superior and below St. Regis were supplied by the U.S. Geological Survey (USGS).

Stream Flow and Water Velocity

Stream flow and water velocity were measured with Marsh-McBirney instantaneous or Price AA current meters, except on the main stem of the Clark Fork River where stream flow was monitored by continuous recording USGS gage stations.

Juvenile and Adult Fish Populations

Fry Nets

Timing and abundance of fry outmigration from tributaries were evaluated using square framed 0.68 m² drift nets with graduated mesh ranging from 6.4 mm (1/4 in.) immediately inside the net opening to 1.6 mm (1/16 in.) in the conical shaped collecting bag. The drift nets were fished in a stationary position in the water column overnight at each site. The volume of water filtered was measured with a current meter positioned at the center of the net orifice.

After the net was retrieved from the stream, trout fry and other fish species were identified and counted. Trout fry were measured to the nearest millimeter in total length and released at the capture site. The fry drift nets were primarily effective for sampling age 0 and I outmigrants.

"Idaho Weir" Fish Traps

Idaho weir fish traps set in the lower reaches of tributaries were used to monitor trout spawning migrations from the river into tributaries. The traps were developed from specifications provided by the Region I Office, Idaho Fish and Game, 2320 Government Way, Coeur d'Alene, ID 83814 (Greg Mouser, personal communication). A detailed description of construction of these traps and procedures for their installation and use will be provided in the completion report for this project. The Idaho weir fish traps were primarily effective for monitoring upstream and downstream movements of adult trout.

Boom-suspended Electrofishing

A boom-suspended electrofishing system was used to sample fish populations on the main stem of the Clark Fork River and in the lower reach of the Bitterroot River. The electrofishing system was adapted from Novotny and Priegel (1974) and is described by Berg (1981). The electrofishing apparatus were mounted on a 4.5 m (14.6 foot) aluminum drift boat powered by a 9.9 horsepower outboard and a 6.1 m (20 foot) aluminum jet boat powered by a 215 horsepower inboard.

The boom-suspended electrofishing apparatus was the most effective technique for sampling fish in the Clark Fork main stem and lower Bitterroot rivers. Much of the boom-suspended electrofishing was accomplished at night due to increased efficiency.

Mobile Electrofishing

A mobile electrofishing system was used to sample fish in tributaries larger than about 10 cfs. The system was also used to sample juvenile and forage fish along shoreline areas of the Clark Fork River.

The mobile electrofishing system consisted of a hand-held mobile positive electrode, a stationary negative electrode mounted on a 1.0 m² float attached to the boat and a portable 1350-watt, 115 volt (60 Hz single phase) alternating current generator. A Coffelt model VVP-2C rectifying unit was used to change the alternating current to pulsed direct current. Output from the rectifying unit was adjustable from 0 to 300 volts half-wave 60 Hz in 25 to 50 volt increments. The electrofishing system was carried in a 5.8 m (19 foot) aluminum freight canoe. In tributaries where the freight canoe could not be floated, electrofishing with this system was accomplished by bank shocking with 76.2 m (250 feet) of 16/2 electrical cord.

Backpack Electrofishing

A backpack electrofishing system was used to sample fish in tributaries smaller than about 10 cfs. Coffelt model BP-6 and Smith-Root Type V A backpack electrofishers were utilized. The backpack electrofishing system consisted of a hand-held mobile positive electrode, a negative electrode consisting of braided copper wire and the portable backpack rectifying and battery or generator unit.

Fish Sample Processing and Tagging

Fish captured by various methods were measured to the nearest mm in total length and weighed to the nearest 10g. Sex and spawning condition (gravid, ripe or spawned) were recorded for fish captured during their spawning season. Several thousand catchable game fish were marked with individually numbered Floy t-tags to evaluate growth rate, movement and angler harvest. All fish were released near the capture site.

Fish Population Estimates

Population estimates were made using the Peterson mark-recapture formula as modified by Chapman (1951):

$$N = \frac{(M+1)(C+1)}{(R+1)} - 1$$

where: N = population estimate
 M = the number of marked fish
 C = the number of fish in the recapture sample
 R = the number of marked fish in the recapture sample (C)

Multiple marking and recapture runs were often needed to collect an adequate sample size. A partial fin clip or fin punch was used to mark the fish. A minimum of two weeks was allowed before recapture runs were made. Additional methods used for population and standing crop estimates are described by Vincent (1971 and 1974).

Fish Aging

Scales were collected from some fish for age determination. The scale samples were imprinted on an acetate slide, and the imprints were projected at 44X on a Norwest nmi 90 microfiche reader. Annuli were identified and ages assigned following procedures described by Jearld (1983) and Tesch (1971).

FINDINGS, RESULTS AND DISCUSSION

Description of Area and Location of Study Sections

This study area lies in west central Montana and includes a 192.1-kilometer (km) (119.4 - mile) reach of the main stem of the Clark Fork River from Milltown Dam to the confluence of the Flathead River. Five study sections, Milltown Dam, Missoula, Huson, Superior and St. Regis, were established in this reach (Figure 1). In addition, perennial tributaries to the Clark Fork River in this reach were studied. The principal tributaries include the Bitterroot and St. Regis rivers and Rattlesnake, Ninemile, Sixmile, Petty, Fish, Trout, Cedar and Tamarack creeks.

The Clark Fork River forms at the confluence of Silver Bow and Warm Springs creeks near Anaconda, Montana, and flows northwestward approximately 560 km (350 river miles) to Lake Pend Oreille in northern Idaho. The 192 km reach of the Clark Fork covered by this study is entirely free-flowing. The drainage area in this reach is mountainous and is covered with large forested tracts, the continuity of which is broken by grazing and cropland areas which are situated in valleys at lower elevations.

The Clark Fork Basin has been widely known for its mining and smelting industries. The copper mines at Butte and smelters at Anaconda, located in the headwaters of this drainage, are internationally famous. The smelters at Anaconda are presently shut down, while mining operations at Butte were resumed in July, 1986, after being shut down for several years. Logging, lumbering and paper manufacturing industries are supported by forests of the basin. Tourist trade is a large contributor to the economy. The basin is nationally known for its scenic beauty, fishing, hunting and other recreational features. Agriculture is also an important industry in the basin.

Four hydropower dams are located on the main stem of the Clark fork River upstream from Lake Pend Oreille. Milltown Dam, the upstream boundary of the present study area, is located 362 km upstream from Lake Pend Oreille. Thompson Falls, Noxon Rapids and Cabinet Gorge dams are situated on the lower Clark Fork River 113, 50 and 18 km upstream from Lake Pend Oreille. Thompson Falls Dam is located 57 km downstream from the lower boundary of the present study area. The four main stem dams contain little storage capacity and have little influence on seasonal discharge patterns.




Figure 1. Map of Clark Fork River showing location of study sections.

Drainage Area and Stream Discharge

The drainage area of the middle Clark Fork River increases from 15,537 km² to 27,736 km², or by about 79 percent, between Milltown Dam and the confluence of the Flathead River (USGS 1983). Average stream discharge increases from 86.38 m³/sec (3050 ft³/sec) to 214.75 m³/sec (7,583 ft³/sec), or by about 149 percent between these boundaries. The drainage area and stream discharge statistics do not include the Flathead River drainage.

Stream flow is monitored by the USGS at gages located 4.5 km downstream from Milltown Dam (Milltown Dam gage), 1.6 km downstream from the confluence of the Bitterroot River (Missoula gage), and 0.6 km downstream from the confluence of the St. Regis River (St. Regis gage). Mean annual discharges for 54-year periods of record are 2.72 km³/year (2,210,000 acre-feet/yr) at Milltown Dam and 4.95 km³/year (4,014,000 acre-feet/yr) at Missoula compared to 6.77 km³/year (5,494,000 acre-feet/yr) at St. Regis for a 73-year period of record.

Stream Gradient

The Clark Fork River enters the study area immediately below Milltown Dam at an elevation of 987.6 m (3,240 ft) msl, dropping 231.6 m (760 ft) to an elevation of 755.9 m (2,480 ft) msl near the confluence of the Flathead River (Table 1). Stream gradient averages 1.23 m/km (6.48 ft/mi) and varies from 0.81 m/km (4.26 ft/mi) between Cedar and Dry creeks to 2.81 m/km (14.81 ft/mi) between Milltown Dam and Marshall Creek. Stream gradients were determined by measurements taken from USGS topographic maps.

Table 1. Stream gradients of the middle Clark Fork River from Milltown Dam to confluence of the Flathead River.

Kilometer	Approximate Location	Elevation (meters, msl)	Gradient (m/km)	Gradient (ft/mi)
586.3	Milltown Dam	987.6	-	-
582.0	Marshall Creek	975.4	2.81	14.81
574.4	Rattlesnake Creek	963.2	1.61	8.51
564.1	Bitterroot River	944.9	1.78	9.38
549.8	Harper's Bridge	929.6	1.06	5.62
540.6	Mill Creek	920.5	1.00	5.26
508.3	Petty Creek	890.0	0.94	4.98
491.7	Fish Creek	853.4	2.21	11.65
462.3	Cedar Creek	816.9	1.24	6.56
447.1	Dry Creek	804.7	0.81	4.26
422.8	Tamarack Creek	780.3	1.00	5.30
397.6	Flathead River	755.9	0.97	5.10

Water Temperature

Water temperatures were monitored on the Clark Fork River near Milltown Dam, Petty Creek, Superior and St. Regis and in the lower reaches of Fish Creek and the St. Regis River during the report period. The data are on file and will be presented in the completion report for this project.

Fish Species Composition

Fifteen species representing six families of fish occur in the middle Clark Fork River between Milltown Dam and the confluence of the Flathead River (Table 2). The bulk of the sport fishery in this 192.1-kilometer (119.4-mile) reach of the river is provided by rainbow trout along with a few brown, bull and westslope cutthroat trout. Mountain whitefish provide an important winter sport fishery. Common nongame fish species found in this reach include squawfish, redbside shiners, longnose dace, largescale suckers and slimy sculpins.

Trout Population Estimates

Trout populations have been estimated by electrofishing and mark/recapture procedures in five study sections on the Clark Fork River. The study sections are located in the vicinities of Milltown Dam, Missoula, Huson, Superior, and St. Regis (Table 3). Estimates in the five study sections indicate the river supports from 175 to 402 catchable rainbow trout per mile (Table 4). Rainbow comprised more than 90 percent of the catchable trout population in all of the study sections. Catchable brown, westslope cutthroat and bull trout were present in the river, but their numbers were usually too low to estimate. In September, 1986, estimates of 16 catchable brown and 22 catchable westslope cutthroat trout per mile were obtained in the Missoula study section.

This density of catchable trout is less than expected for comparable trout streams the size of the Clark Fork. While the Clark Fork River supports an average of two to four hundred catchable trout per mile, other large trout rivers in Montana often support two to three thousand or more catchable trout per mile (Berg 1984).

Major tributaries to the Clark Fork River support larger populations of catchable trout than the main stem of the river. The mean number of catchable rainbow trout per mile in the Blackfoot River over a three-year period from 1983 to 1985 was 445 percent larger than the mean number of catchable rainbow per mile in the Clark Fork River during a three-year period from 1984 to 1986 (Tables 4 and 5). The comparison of the Blackfoot River with the Clark Fork is appropriate since both rivers have similar physical habitat characteristics. Higher water quality in the Blackfoot River appears to be the major difference between the two rivers.

Table 2. Fish species found in the Clark Fork River in Montana between Milltown Dam and the confluence of the Flathead River.

SALMONIDAE (Trout Family)	
<u>Prosopium williamsoni</u> - Mountain whitefish	A ^{1/}
<u>Salmo clarki lewisi</u> - Westslope cutthroat trout	R*
<u>Salmo gairdneri</u> - Rainbow trout	C
<u>Salmo trutta</u> - Brown trout	R*
<u>Salvelinus fontinalis</u> - Brook trout	R*
<u>Salvelinus confluentus</u> - Bull trout	R
ESOCIDAE (Pike Family)	
<u>Esox lucius</u> - Northern pike	R
CYPRINIDAE (Minnow Family)	
<u>Mylocheilus caurinus</u> - Peamouth	R
<u>Ptychocheilus oregonensis</u> - Squawfish	A
<u>Rhinichthys cataractae</u> - Longnose dace	C
<u>Richardsonius balteatus</u> - Redside shiner	A
CATOSTOMIDAE (Sucker Family)	
<u>Catostomus catostomus</u> - Longnose sucker	R
<u>Catostomus macrocheilus</u> - Largescale sucker	A
CENTRARCHIDAE (Sunfish family)	
<u>Micropterus salmoides</u> - Largemouth bass	R
COTTIDAE (Sculpin Family)	
<u>Cottus cognatus</u> - Slimy Sculpin	C

^{1/} Relative Abundance - A = Abundant, C = Common, R = Rare.

* Common in some tributaries of the Clark Fork in the study area.

Table 3. Location, length and river mile index boundaries of fish population study sections on the Clark Fork River.

Section Name	Description of Location	Section Length (mi)	River Mile Index Boundaries
Milltown	Milltown Dam to 2.8 miles upstream from confluence of Rattlesnake Cr.	3.4	364.4 to 361.0
Missoula	Confluence of Bitterroot R. to 0.5 mile upstream from Harper Bridge	8.6	350.5 to 341.9
Huson	Confluence of Sixmile Cr. to 4.0 miles upstream from confluence of Petty Cr.	4.5	328.2 to 323.7
Superior	Confluence of Cedar Cr. to confluence of Dry Cr.	6.3	286.6 to 280.3
St. Regis	Confluence of St. Regis R. to 1.6 miles downstream	1.6	270.7 to 269.1

Table 4. Trout population estimates in five study sections of the Clark Fork River.

Study Section	Date of Estimate	Fish Species	Section Length(mi)	Catchable <u>1</u> / Trout/Section	Catchable <u>1</u> / Trout/Mile
Missoula	Sept. 1984	Rainbow	8.6	1506	175
Missoula	June 1985	Rainbow	8.6	1804	210
Milltown	June 1985	Rainbow	3.6	1035	288
Superior	July 1985	Rainbow	6.3	1382	219
Huson	Sept. 1985	Rainbow	4.5	1749	389
Missoula	Sept. 1986	Rainbow	8.6	3461	402
		Brown	8.6	137	16
		W.S. Cutthroat	8.6	187	22
Huson	Sept. 1986	Rainbow	4.5	1504	334
St. Regis	Sept. 1987	Rainbow	1.6	345	<u>216</u>
				All sections-Rainbow Mean (<u>x</u>)	279

1/ Catchable trout 7-inches total length and larger.

Table 5. Trout population estimates in the Johnsrud section of the Blackfoot River, approximately 13 miles upstream from Bonner.

Date of Estimate	Fish Species	Section Length (mi)	Catchable <u>1</u> / Trout/Section	Catchable <u>1</u> / Trout/Mile
June 1985	Rainbow	3.6	5,225	1,451
June 1984	Rainbow	3.6	3,186	885
June 1983	Rainbow	3.6	<u>5,445</u>	<u>1,512</u>
		Mean (<u>x</u>)	4,618	1,282

1/ Catchable trout 7-inches total length and larger.

Scales were collected from trout during population estimates to determine growth rates and age structure of the trout populations. Preliminary findings indicate growth rates of trout in the Clark Fork are relatively high when compared to trout streams of similar size. This indicates that food supply is probably not a limiting factor for trout populations in the Clark Fork River. Furthermore, it suggests that the Clark Fork River may be "under seeded" and that recruitment may be a limiting factor. Estimates of size and age composition, growth rates, biomass, and condition factors of trout populations in the middle Clark Fork River will be presented in the completion report for this project.

Trout Spawning and Rearing Habitat in the Clark Fork River

Visual surveys have been made in the Milltown, Missoula, Huson and Superior study sections during the rainbow and brown trout spawning periods in an attempt to locate trout redds. To date, only brown trout redds have been located in the river. They were found in the Milltown and Missoula sections. Since a very limited amount of time has been spent on the visual surveys, additional observations must be made to evaluate the extent of trout spawning in the river.

The search for trout redds in the Clark Fork River is hindered during both rainbow and brown trout spawning periods by poor visibility in deep water areas where spawning could occur. Visibility is sometimes precluded even in shallow water during the rainbow trout spawning period due to highly turbid spring runoff conditions. For this reason, use of the Clark Fork River for trout spawning is also being evaluated by electrofishing during the spawning periods in an attempt to locate concentrations of mature fish in spawning condition.

Suitable rainbow, cutthroat and brown trout rearing habitat is found primarily along the edge of the Clark Fork River's channel. Limited electrofishing surveys of this habitat indicated young-of-the-year trout were relatively more abundant in the Milltown and Superior study sections than in the Missoula and Huson sections (Table 6).

Young-of-the-year trout were relatively scarce in all four study areas. In the Missouri River below Holter Dam, a stream comparable in size to the Clark Fork below Milltown Dam, young-of-the-year trout were at least ten times more abundant than in the Clark Fork (Berg 1983).

Table 6. Average size and relative abundance of young-of-the-year trout sampled by electrofishing in four study sections of the Clark Fork River during late summer 1985.

Study Section	Date	Trout Species	Average Length (mm)	Juvenile Trout Electrofished/Hour
Milltown	8/26/85	Rainbow	57	7.1
		Brown	90	10.7
Missoula (side channel)	8/28/85	Rainbow	76	1.7
		Brown	94	10.0
Missoula (main river)	8/28/85	Rainbow	63	1.4
		Brown	-	0.0
Huson	8/30 & 9/4/85	Rainbow	60	3.6
		Brown	77	0.3
Superior	9/5/85	Rainbow	58	14.6
		Brown	81	1.1

Tributary Trout Spawning Migrations

In an effort to evaluate spawning periodicity and sources of trout recruitment in the middle Clark Fork River, the lower reaches of several tributaries were electrofished or trapped during trout spawning periods to locate spawning migrants from the Clark Fork River.

Most members of the trout family migrate during the spawning season in search of suitable spawning sites (Hubbs and Lagler 1970). Spawning movements of lake dwelling salmonid populations into inlet or outlet streams have been extensively documented for rainbow (Rayner 1942, Hartman et al. 1962, Calhoun 1966, Scott and Crossman 1973) and brown trout (Fenderson 1958, Stuart 1957) and mountain whitefish (Snyder 1918, Calhoun 1966).

Less information is available on spawning movements of river dwelling salmonid populations into feeder streams. Calhoun (1966) reports resident rainbow trout populations in streams tend to move upstream, and if possible into tributaries to spawn. River dwelling brown trout in Ontario normally seek tributary streams for spawning purposes (MacKay 1963). Spawning movements of mountain whitefish from larger streams into some tributaries have been observed in Montana (Liebelt 1970, Brown 1971).

Electrofishing and "Idaho weir" fish trapping surveys indicate rainbow, brown and westslope cutthroat trout migrate from the Clark Fork River into tributaries to spawn (Berg 1986). Trout fry outmigrations from several tributaries, monitored with fry traps, indicated tributaries provide recruitment of juvenile trout to the Clark Fork River (Table 7). Monitoring of adult trout spawning migrations into tributaries and of fry outmigrations from tributaries in the vicinities of Superior and St. Regis was accomplished during this report period. Analysis of this data is in progress and findings will be presented in the next progress report.

Table 7. Trout fry outmigration rates monitored in five tributaries of the Clark Fork River during 1985 (Berg 1986).

Stream	Total Trap Nights	Rainbow Trout		W.S. Cutthroat Trout		Brown Trout		Bull Trout	
		Total Number Captured	Average Catch/ Trap Night						
Fish Cr	57	626	11.00	25	0.44	3	0.05	1	0.02
Ninemile Cr	46	493	10.72	0	0.00	14	0.30	0	0.00
Petty Cr	49	346	7.06	7	0.14	0	0.00	0	0.00
Rattlesnake Cr	31	65	2.10	1	0.03	1	0.03	0	0.00
Sixmile Cr	5	4	0.80	0	0.00	0	0.00	0	0.00

Angler Harvest Rates and Fish Movement Patterns

A total of about 7,000 trout have been marked with individually numbered Floy T-tags since the inception of this study. Tags recovered during our surveys will be used to evaluate trout movement patterns in the middle Clark Fork River drainage.

An indication of angler harvest of trout in the Clark Fork River and its tributaries is being provided by angler-returned fish tags. Preliminary estimates suggested westslope cutthroat and bull trout were relatively more vulnerable to harvest than rainbow and brown trout (Berg 1986).

Since large numbers of tagged trout are still at large in the study area, tag returns are being updated on a daily basis. A computer program is being developed to analyze trout movement patterns and angler harvest rates. An updated summary of these findings will be presented in the next progress report.

CONCLUSIONS

The middle Clark Fork River and its tributaries support a fishery with substantial recreational value. The sport fishery is provided mainly by rainbow trout and a few brown, bull and westslope cutthroat trout. However, catchable trout population numbers are considerably lower than expected for a river of its size.

A variety of factors probably contribute toward suppressing the fishery in this reach of river. Water quality degradation factors which may be influencing the fishery include the Frenchtown pulp mill and Missoula sewage treatment plant effluents, potentially toxic heavy metals originating from mine tailings in the upper Clark Fork drainage and fine sediments originating from various human related activities which could impair trout food production or trout reproductive success. Stream dewatering and water temperature affects from irrigation water withdrawals also influence the river fishery particularly through indirect effects on tributary streams which typically are more severely dewatered than the main river.

Trout population estimates presently can not be used to differentiate the effects of the various factors on the sport fishery. However, the estimates do indicate that trout populations are depressed in the Clark Fork River from Milltown Dam to St. Regis despite the inflow of major tributaries with relatively high water quality. Findings from studies conducted to date suggest that if water quality is improved in the middle Clark Fork River, it should be capable of supporting larger populations of catchable trout.

Saturation plants of 10,000 hatchery reared young-of-the-year brown trout were made in the Huson study section during the early and late summers of 1986 and 1987, respectively, to aid in evaluating whether recruitment is a limiting factor for trout populations in the Clark Fork

River. Spawn were taken from a wild stock of brown trout at Harrison Lake, Montana, for the 1986 plant and from a wild stock of brown trout from Warm Springs Creek, Montana, for the 1987 plant. The eggs were fertilized and incubated at the Washoe Park State Fish Hatchery. The brown trout were reared in the hatchery until they were 2 to 4 inches in total length before being planted in the Huson section. Electrofishing surveys and population estimates will be continued in the Huson section to determine whether these fish eventually recruit into the adult population.

Prepared by: Rodney K. Berg

Date: August 10, 1988

Waters Referred to:

Clark Fork River, Sec. 02	05-1456
Clark Fork River, Sec. 03	06-1121
Fish Creek	05-2768
Ninemile Creek	05-5168
Petty Creek	05-5552
Rattlesnake Creek	06-5149
Sixmile Creek	05-6368

Key Words:

Trout spawning - Tributaries/mainstem
 Westslope cutthroat trout
 Bull trout
 Trout fry outmigrations
 Trout population estimates

LITERATURE CITED

- Berg, R. K. 1975. Fish and game planning, Upper Yellowstone and Shields River drainages. Job Comp. Rept., Fed. Aid to Fish and Wildl. Rest. Proj. No. FW-3-R. Job 1-A. 92 pp.
- _____. 1981. Fish populations of the Wild and Scenic Missouri River, Montana. Job Comp. Rept., Fed. Aid to Fish and Wildl. Rest. Proj. No. FW-3-R. Job 1-A. 242 pp.
- _____. 1983. Middle Missouri River Planning project. Job Prog. Rept., Fed. Aid to Fish and Wildl. Rest. Proj. No. FW-3-R-11. Job 1-A 30 pp.
- _____. 1984. Trout heaven. Montana Outdoors, Sept./Oct. 1984, 27-30 pp.
- _____. 1986. Lower Clark Fork basin investigations. Job Prog. Rept., Fed. Aid to Fish and Wildl. Rest. Proj. No. F-37-R-1. 39 pp.
- Brown, C. J. D. 1971. Fishes of Montana. Endowment and Res. Found., Mont. St. Univ., Bozeman. 207 pp.
- Calhoun, A. J. 1966. Inland Fisheries management. Calif. Dept. of Fish & Game, Sacramento. 546 pp.
- Chapman, D. G. 1951. Some properties of the hypergeometric distribution with applications to zoological sample censuses. Univ. of Calif. Pub. in Stat. 1 (7): 131-160.
- Fenderson, C. N. 1958. Brown trout, Salmo trutta Linnaeus. Fishes of Maine, 2nd Ed. Ed. W. H. Everhart. 34-37 pp.
- Hartman, G. F., T. G. Northcote and C. C. Lindsey. 1962. Comparison of inlet and outlet spawning runs of rainbow trout in Loon Lake, British Columbia. J. Fish. Res. Bd. Can. 19(2): 173-200.
- Hubbs, C. L. and K. F. Lagler. 1970. Fishes of the Great Lakes region. Univ. of Mich. Press, Ann Arbor. 213 pp.
- Jearld, R. A. 1983. In Fisheries Techniques. American Fisheries Society, Bethesda, Maryland.
- Liebelt, J. 1970. Studies on the behavior and life history of the mountain whitefish (Prosopium williamsoni - Girard). PhD. Thesis, Mont. St. Univ. 45 pp.
- MacKay, H. H. 1963. Fishes of Ontario. The Bryant Press Ltd. Toronto, On. 300 pp.
- Novotny, D. W. and G. R. Priegel. 1974. Electrofishing boats - improved designs and operational guidelines to increase the effectiveness of boom shockers. Wisc. Dept. Nat. Resc. Tech. Bull. No. 73. 48 pp.

- Rayner, H. J. 1942. The spawning migration of rainbow trout at Skaneateles Lake, New York. *Trans. Am. Fish. Soc.* 71:180-183.
- Scott, W. B. and E. J. Crossman. 1973. *Freshwater fishes of Canada.* Fish. Res. Bd. of Canada, Ottawa. 966 pp.
- Snyder, J. O. 1918. The fishes of the Lahontan system of Nevada and northeastern California. *Bull. US Bur. Fish.* 35:31-86.
- Stuart, T. A. 1957. The migration and homing behavior of brown trout. *Freshw. Salm. Fish. Res. Scot.* 18:3-27.
- Tesch, F. W. 1971. Age and growth. In *Methods for assessment of fish production in fresh waters.* IBP handbook No. 3 Blackwell Scientific Pub., Oxford and Edinburgh, England. 348 pp.
- USGS. 1983. *Water resources data for Montana.* US Dept. of Interior.
- Vincent, E. R. 1971. River electrofishing and fish population estimates. *Prog. Fish. Cult.* 33(3):163-167.
- _____. 1974. Addendum to river electrofishing and fish population estimates. *Prog. Fish. Cult.* 36(3): 182.
- Wipperman, A. H. 1973. *Smith River drainage inventory and planning investigation.* Job Comp. Rept., Fed. Aid to Fish and Wildl. Rest. Proj. No. FW-1-R, Job 1a.