

IMPACTS OF STOCKING HERBIVOROUS FISHES
FOR AQUATIC MACROPHYTE REMOVAL UPON
SOUTH DAKOTA PONDS

BY

JOHN CHRISTOPHER YOUNG

A thesis submitted
in partial fulfillment of the requirements
for the degree, Master of Science, Major in
Wildlife and Fisheries Sciences
(Fisheries Option)
South Dakota State University

1987

IMPACTS OF STOCKING HERBIVOROUS FISHES
FOR AQUATIC MACROPHYTE REMOVAL UPON
SOUTH DAKOTA PONDS

This thesis is approved as a creditable and independent investigation by a candidate for the degree, Master of Science, and is acceptable for meeting the thesis requirements for this degree. Acceptance of this thesis does not imply that the conclusions reached by the candidate are necessarily the conclusions of the major department.

Timothy M. Moe 4/1/87

Dr. Timothy Moe
Major Advisor

Date

Charles G. Scalet 4/1/87

Dr. Charles G. Scalet
Head, Wildlife and
Fisheries Sciences
Department

Date

ACKNOWLEDGEMENTS

I would like to thank Dr. Timothy Modde for his guidance and encouragement during this study and in the drafting of this manuscript. I would also like to thank David Gilbraith for his cooperation and partnership for the three years of this study.

I would like to thank Drs. Charles G. Scalet and Gary E. Larson for their recommendations and critical reviews of the manuscript.

I would also like to thank Keith Wintersteen, Dave Cernik, and Steve Morrow for their assistance in the field portion of this study.

Funding and equipment were provided by the South Dakota Department of Game, Fish and Parks, the South Dakota State University Agricultural Experiment Station, and the Department of Wildlife and Fisheries Sciences.

IMPACTS OF STOCKING HERBIVOROUS FISHES
FOR AQUATIC MACROPHYTE REMOVAL UPON
SOUTH DAKOTA PONDS

Abstract

Aquatic macrophyte removal by herbivorous fishes was investigated to determine changes in the predator-prey relationship of largemouth bass, Micropterus salmoides, and bluegills, Lepomis macrochirus, stocked into 15 prairie ponds. Monosex grass carp, Ctenopharyngodon idella, and hybrid grass carp, Ctenopharyngodon idella x Hypophthalmichthys nobilis, were each stocked at a rate of 247 fish/hectare into five study ponds. Largemouth bass and bluegills were each stocked into 12 study ponds at 247 fish/hectare in July, and 1235 fish/hectare in September, 1984, respectively. Additional largemouth bass were stocked at a rate of 247 fish/hectare into nine study ponds in July, 1985. Conductivity was the only water quality variable which differed significantly ($P \leq 0.05$) due to treatments. Substantial, but not significant ($P > 0.05$), differences in vegetation removal were observed among ponds. The failure to detect statistical differences likely resulted because of greater variability among ponds within treatments than between treatments. Vegetation removal appeared to be greatest in the monosex grass carp treatment with slight differences between the control and hybrid grass carp treatment ponds. Water transparency declines, as measured by secchi disc visibility and turbidity, were greatest in the monosex grass carp treatment. The evaluation of vegetation removal on largemouth bass-

bluegill interactions was not possible because of the failure of bluegills to reproduce. Survival of stocked bluegills was low as a result of small stocking length. Mean survival rate for age-I largemouth bass was 38% for eight ponds. Mean growth and mean W_r for age-I largemouth bass was 160 mm and 119 g, respectively. Abundance of age-0 largemouth bass appeared to be influenced greater by the survival rate of age-I largemouth bass than by vegetation abundance.

TABLE OF CONTENTS

	Page
INTRODUCTION	1
METHODS	5
Study Area	5
Design	5
Field and Laboratory Methods	7
Analysis	12
RESULTS	18
Pond Selection Evaluation	18
Water Quality	18
Vegetation Removal	20
Fish Populations	30
DISCUSSION	37
RECOMMENDATIONS	43
LITERATURE CITED	44
APPENDIX TABLES	52

LIST OF FIGURES

Figure	Page
1. Map of South Dakota illustrating locations of control ponds (1-5), hybrid grass carp ponds (6-10), and monosex grass carp ponds (11-15). Pond code numbers underscored identify ponds eliminated from fish data analysis due to fish contamination. * Indicates one hybrid grass carp pond which winterkilled during 1984	6
2. Mouth widths of age-I largemouth bass, <u>Micropterus salmoides</u> , in study ponds with (■) and without (□) fathead minnows, <u>Pimephales promelas</u> compared to body depths of age-0 large-mouth bass with (●) and without (○) fathead minnows, and initial body depth of age-0 largemouth bass (△) at pond stocking 10 July 1985	17
3. Control and treatment means (horizontal bars), and standard error (rectangles) for secchi disk visibility sampled from 14 South Dakota study ponds between June, 1984, and August, 1985	26
4. Control and treatment means (horizontal bars) and standard error (rectangles) for turbidity sampled in 14 South Dakota study ponds between June, 1984, and August, 1985	27
5. Control and treatment raw means (horizontal bars) and standard deviations (rectangles) of aquatic vegetation heights sampled from 1.5 - 2.1 m transect water depths for control ponds (white), hybrid grass carp ponds (striped), and monosex grass carp ponds (black) for 13 South Dakota ponds between July, 1984 and August, 1985	29
6. Control and treatment raw means (horizontal bars) and standard deviations (rectangles) of aquatic vegetation heights sampled from 2.4 - 3.0 m transect water depths for control ponds (white), hybrid grass carp ponds (striped), and monosex grass carp ponds (black) for 13 South Dakota study ponds between July, 1984, and August, 1985	31
7. Second order regression of the abundance of age-0 largemouth bass, <u>Micropterus salmoides</u> , and survival rate of age-I largemouth bass from three control ponds (C), two hybrid grass carp ponds (H), and four monosex grass carp ponds (M) sampled between 17 and 30 September, 1985	35

LIST OF FIGURES

Figure	Page
1. State of South Dakota illustrating locations of control ponds (1-5), hybrid grass carp ponds (6-10), and monosex grass carp ponds (11-15). Pond code numbers underscored identify ponds eliminated from fish data analysis due to fish contamination. * Indicates one hybrid grass carp pond which winterkilled during 1984	6
2. Mouth widths of age-I largemouth bass, <u>Micropterus salmoides</u> , in study ponds with (■) and without (□) fathead minnows, <u>Pimephales promelas</u> compared to body depths of age-0 large-mouth bass with (●) and without (○) fathead minnows, and initial body depth of age-0 largemouth bass (△) at pond stocking 10 July 1985	17
3. Control and treatment means (horizontal bars), and standard error (rectangles) for secchi disk visibility sampled from 14 South Dakota study ponds between June, 1984, and August, 1985	25
4. Control and treatment means (horizontal bars) and standard error (rectangles) for turbidity sampled in 14 South Dakota study ponds between June, 1984, and August, 1985	26
5. Control and treatment raw means (horizontal bars) and standard deviations (rectangles) of aquatic vegetation heights sampled from 1.5 - 2.1 m transect water depths for control ponds (white), hybrid grass carp ponds (striped), and monosex grass carp ponds (black) for 13 South Dakota ponds between July, 1984 and August, 1985	29
6. Control and treatment raw means (horizontal bars) and standard deviations (rectangles) of aquatic vegetation heights sampled from 2.4 - 3.0 m transect water depths for control ponds (white), hybrid grass carp ponds (striped), and monosex grass carp ponds (black) for 13 South Dakota study ponds between July, 1984, and August, 1985	30
7. Second order regression of the abundance of age-0 largemouth bass, <u>Micropterus salmoides</u> , and survival rate of age-I largemouth bass from three control ponds (C), two hybrid grass carp ponds (H), and four monosex grass carp ponds (M) sampled between 17 and 30 September, 1985	36

LIST OF TABLES

Table	Page
1. Pond name, group, date detected, and fish species contaminating nine South Dakota study ponds between May, 1984, and May, 1985	8
2. Pond name, group, code number from Figure 1, and comparison of surface areas (hectares) estimated with Missouri field chart and standard survey methods for 15 South Dakota study ponds between August, 1983, and August, 1984	10
3. Pond name, group, population estimate (N), sampling variance V(N) and 95% confidence interval around N for age-I largemouth bass, <u>Micropterus salmoides</u> , from eight South Dakota study ponds sampled between April and October, 1985	13
4. Pond Name, group, population estimate, number of fish stocked in study ponds in 1984, and survival rate (percent) for age-I largemouth bass, <u>Micropterus salmoides</u> , from eight South Dakota study ponds sampled between 7 and 26 September 1985	14
5. Pond name, group, number of fish electrofished per hour for age-I bluegills, <u>Lepomis macrochirus</u> , and age-0 and age-I largemouth bass, <u>Micropterus salmoides</u> , sampled from nine South Dakota study ponds between August and September, 1985	16
6. Nested factorial analysis of variance for differences in conductivity (micromhos/cm) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	19
7. Nested factorial analysis of variance for differences in salinity (parts per thousand) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	21
8. Nested factorial analysis of variance for differences in total hardness (mg/l as CaCO ₃) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	22

LIST OF TABLES (Continued)

Table	Page
9. Nested factorial analysis of variance for differences in sulfate (mg/l) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	23
10. Nested factorial analysis of variance for differences in potassium (mg/l) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	24
11. Nested factorial analysis of variance for differences in pond surface levels (cm) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	25
12. Nested factorial analysis of variance for differences in aquatic vegetation height in (cm) due to treatments, sessions, years, and transect water depths sampled from 13 South Dakota study ponds between July, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	28
13. Waller-Duncan K-Ratio t-test to determine differences in first year growth of largemouth bass, <u>Micropterus salmoides</u> sampled in four study ponds without fathead minnows between April and June, 1985	32
14. Waller-Duncan K-Ratio t-test to determine differences in W_r of age-I largemouth bass, <u>Micropterus salmoides</u> sampled in four study ponds without fathead minnows between April and June, 1985	33
15. Pond name, sampling date, number of fish electrofished per hour, and TL (mm) for age-I bluegills, <u>Lepomis macrochirus</u> from four study ponds without fathead minnows between May and September, 1985	34

LIST OF APPENDIX TABLES

Appendix	Page
Table 1. Nested factorial analysis of variance for differences in secchi disk visibility (m) and maximum pond depth (cm) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt)	52
Table 2. Nested factorial analysis of variance for differences in conductivity (micromhos/cm ²) and salinity (parts/thousand) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt)	53
Table 3. Nested factorial analysis of variance for differences in turbidity (formazin turbidity units) and pH (standard units) due to treatments and sessions sampled from 15 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for treatment F-tests indicated by (Trt)	54
Table 4. Nested factorial analysis of variance for differences in phenol and total alkalinity (mg/l as CaCO ₃) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatments F-tests indicated by (Trt)	55
Table 5. Nested factorial analysis of variance for differences in total hardness (mg/l as CaCO ₃) and nitrate-nitrogen (mg/l) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt)	56
Table 6. Nested factorial analysis of variance for differences in phosphate (mg/l) and sulfate (mg/l) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt)	57
Table 7. Nested factorial analysis of variance for differences in potassium (mg/l) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt)	58

LIST OF APPENDIX TABLES (Continued)

Appendix	Page
Table 8. Nested factorial analysis of variance for differences in pond surface levels (cm) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-test indicated by (Trt)	59
Table 9. Nested factorial analysis of variance for differences in secchi disk visibility (m) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	60
Table 10. Nested factorial analysis of variance for differences in turbidity (formazin turbidity units) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	61
Table 11. Nested factorial analysis of variance for differences in pH (standard units) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	62
Table 12. Nested factorial analysis of variance for differences in phenol alkalinity (mg/l as CaCO_3) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	63
Table 13. Nested factorial analysis of variance for differences in total alkalinity (mg/l as CaCO_3) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984 and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	64
Table 14. Nested factorial analysis of variance for differences in nitrate-nitrogen (mg/l) due to treatments, sessions, and years sampled from 14 study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	65

LIST OF APPENDIX TABLES (Continued)

Appendix	Page
Table 15. Nested factorial analysis of variance for differences in phosphate (mg/l) due to treatments, sessions, and years sampled from 14 study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	66
Table 16. Nested factorial analysis of variance for differences in maximum pond depth (cm) due to treatments, sessions, and years sampled from 14 South DAKota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt)	67

INTRODUCTION

Submergent vegetation frequently reaches nuisance levels in prairie ponds because of shallow water depths and high nutrient loads. Excessive macrophyte densities hamper recreational activities and disrupt predator-prey relationships by reducing predator efficiency (Crowder and Cooper 1979). Investigations of different macrophyte densities demonstrate that increased cover reduces predator efficiency (Glass 1971) by decreasing prey vulnerability and subsequent prey capture rates (Crowder and Cooper 1979). Savino and Stein (1982) reported reduced predation movements of largemouth bass, Micropterus salmoides, upon bluegills, Lepomis macrochirus, with increasing vegetation stem density. Thus bluegills avoid predation by largemouth bass by seeking refuge in aquatic vegetation (Mittlebach 1981).

The traditional largemouth bass-bluegill stocking combination developed in Alabama by Swingle and Smith (1941) must be modified in order to be successful in northern ponds (Modde and Scalet 1986). Excessive bluegill recruitment may occur in northern ponds because bluegills spawn one year before largemouth bass (Dillard and Novinger 1975). Differential growth rates between bluegills and largemouth bass at northern latitudes also decreases the ability of largemouth bass to control bluegill recruitment (Modde and Scalet 1985). Strategies to correct bluegill imbalances in ponds includes stocking more vulnerable prey species such as the fathead minnow, Pimephales promelas, as forage for largemouth bass. Also, split-stocking bluegills in the fall and largemouth bass the subsequent summer should insure adequate bluegill recruitment as forage for largemouth bass fingerlings (Hill 1980).

Aquatic macrophyte removal should decrease searching time for largemouth bass and presumably increase bluegill vulnerability.

Methods to control nuisance aquatic macrophytes includes surface drawdowns, herbicides, pond fertilization, and mechanical harvesting. These options are expensive and often impractical in small northern ponds. However, biological control of aquatic macrophytes utilizing grass carp, Ctenopharyngodon idella, represents an economical and longer lasting alternative (Shireman 1982). These fish feed on a broad range of aquatic plant species (Cross 1969) and are capable of consuming large quantities of vegetation. Hickling (1966) stated that mechanical rupturing of aquatic plant cells is necessary for digestion because these fish lack cellulase. Furthermore, to compensate for poor assimilation and rapid food passage in the gut, grass carp are required to consume voluminous quantities of aquatic vegetation (Hickling 1966). Cross (1969) noted that under optimum conditions grass carp consumed their body weight daily.

Grass carp stockings by state agencies and private hatcheries in the 1960's enabled these fish to gain access to several major river drainages in the central United States (Guillory and Gasaway 1978). Concerns for negative impacts on aquatic systems by feral grass carp populations were heightened with the discovery of grass carp reproduction in the lower Mississippi River (Conner et al. 1980). Several states banned grass carp from their waters (Young et al. 1983) fearing waterfowl habitat destruction (Gasaway and Drda 1977) or negative impacts on native fish faunas such as those reported by Vinogradov and Zolotova (1974) in Russia. In response to this concern

sex reversal, gynogenesis, triploidization, and hybridization have been attempted to produce sterile fish (Jensen et al. 1983).

Hybrid grass carp, an intergeneric cross between female grass carp and male bighead carp, Hypophthalmichthys nobilis (Lazareva et al. 1977), were first produced in Hungary in 1968 and were presumed to be sterile and triploid (Marian and Krasznai 1978). More recently it was determined that both diploid and triploid fish resulted from this cross (Magee and Philipp 1982). Hybrid grass carp were anticipated to provide aquatic macrophyte control while eliminating the chance for natural reproduction. Subsequent research has shown that the hybrids are inferior to grass carp for aquatic vegetation removal (Young et al. 1983; Harberg and Modde 1985; Osborne 1985). Monosex grass carp, an exclusively female population, represent an alternative for aquatic macrophyte control (Boney et al. 1984). Their use should obviously be limited to watersheds where male grass carp are absent.

Documentation of the effects of grass carp introductions on predator-prey interactions are limited in the literature. Previous studies of grass carp introductions have been confined either to lakes (e.g. Ware and Gasaway 1976; Mitzner 1978; Bailey 1978) or hatchery ponds (e.g. Ball 1977; Lembi et al. 1978; Rottman and Anderson 1977; Baur et al. 1979). Savino and Stein (1982) noted that prior to their experimental observations, the relationship between vegetation density and bluegill vulnerability had never been explicitly demonstrated. Therefore, aquatic plant control in ponds using herbivorous fishes requires evaluation to ascertain whether aquatic macrophyte removal will increase prey vulnerability to predation.

The objectives of this study were to:

1. Measure the influence of herbivorous fishes upon aquatic macrophytes,
2. determine changes in water quality attributable to the presence or absence of herbivorous fishes , and
3. determine the impact of herbivorous fishes on predator-prey relationships.

METHODS

Study Area

Study ponds were located in eastern and central South Dakota (Figure 1). This area has a continental-type climate with warm summers and cold winters. Average annual precipitation is 64 cm in the southeast to 33 cm in the northwest of the state (Spuhler et al. 1971). South Dakota is characterized by rolling grasslands intermixed with agricultural crops and pasture.

Design

Criteria used to select study ponds were:

1. maximum pond depth of at least 3.6 m,
2. absence of fish life,
3. surface area of 0.4 - 2.0 hectares, and
4. abundance of aquatic macrophytes

Study ponds were surveyed initially by telephone inquiries utilizing South Dakota Game, Fish and Parks pond stocking requests. Subsequent pond visits evaluated the suitability of each pond site to study criteria.

The study evaluated 15 ponds divided evenly among one control and two treatment groups (monosex grass carp and hybrid grass carp ponds). Control and treatment ponds were distinguished by the absence or presence of herbivorous fishes, respectively. Limited availability of study ponds prevented pond randomization. Herbivorous fishes were stocked into 10 treatment ponds in fall, 1983. Five control ponds were chosen in spring, 1984.




Figure 1. Map of South Dakota illustrating locations of control ponds (1-5), hybrid grass carp ponds (6-10), and monosex grass carp ponds (11-15). Pond code numbers underscored identify ponds eliminated from fish data analysis due to fish contamination. *Indicates one hybrid grass carp pond which winterkilled during 1984.

Study ponds were stocked with fish in the following sequence.

1. Monosex grass carp (mean TL 63.2 mm) were stocked into five treatment ponds at 247 fish/hectare on 18 August, 1983.
2. Hybrid grass carp (mean TL 170.6 mm) were stocked into five treatment ponds at 247 fish/hectare on 7 - 8 September, 1983.
3. Largemouth bass (mean TL 31.7 mm) were stocked into 12 ponds at 247 fish/hectare on 12 - 13 July, 1984.
4. Bluegill (mean TL 31.7 mm) were stocked into 12 ponds at 1235 fish/hectare on 11 - 12 September, 1984.
5. Largemouth bass (mean TL 48.5 mm) were stocked into nine ponds at 247 fish/hectare on 10 July, 1985.

Largemouth bass were stocked in only 12 ponds in 1984 due to contamination of piscivorous fishes in three ponds (Table 1). In 1985 two additional ponds were contaminated and one pond was subject to winterkill.

Field and Laboratory Methods

Physical and chemical water quality variables were sampled during three periods in both 1984 and 1985. Sampling periods for water quality will be referred to as sessions in this study. Conductivity, salinity, and surface temperature were determined with a Yellow Springs Instrument S-C-T meter. Secchi disc visibility in the water column was measured with a standard secchi disc. A 1 liter water sample was collected 30 cm beneath the pond surface near the pond center for chemical analysis. An additional water sample from 1 m depth was returned to the South Dakota

Table 1. Pond name, group, date detected, and fish species contaminating nine South Dakota study ponds between May, 1984, and May, 1985.

Pond Name	Group	Date	Contaminating Fish Species
Kippes	monosex	May, 1984	<u>Pimephales promelas</u>
Fox	monosex	May, 1984	<u>Pimephales promelas</u>
Ortman	monosex	May, 1984	<u>Lepomis cyanellus</u>
Willis	hybrid	May, 1984	<u>Ictalurus nebulosus</u>
Heathon	control	Jul, 1984	<u>Lepomis macrochirus</u>
Ridings	hybrid	Aug, 1984	<u>Pimephales promelas</u>
Thompson	hybrid	Aug, 1984	<u>Pimephales promelas</u>
Dailey	control	May, 1985	<u>Perca flavescens</u>
Dailey	hybrid	May, 1985	<u>Perca flavescens</u>

State Water Quality Lab for free potassium determination. Lembi et al. (1978) recommended sampling potassium as a possible means to measure macrophyte removal by grass carp. Maximum pond depth was determined with a Lowrance X-15 chart recorder.

Nitrate-nitrogen, phosphate, sulfate, and turbidity were measured with a Hach DR-EL spectrophotometer. Phenol and total alkalinity and total hardness were determined by titration. A Hellige Lilliput portable meter was used to measure pH. Surface areas for all ponds were estimated initially utilizing a field chart¹ integrating pond perimeter length with pond shape and later re-estimated by standard survey methods in July, 1984 (Table 2). Wooden stakes calibrated in 10 cm increments were placed into ponds to document water level fluctuations.

Height of aquatic vegetation in the water column was traced with a Lowrance X-15 chart recorder along three transects at two different depths for two sessions in 1984 and three sessions in 1985. The X-15 transducer was mounted at pond surface level on the stern of an outboard motor driven boat. A 30 m rope with floats at 3 m intervals delineated transects in 1.5 - 2.1 m and 2.4 - 3.0 m water depths. Transects were obtained by running the boat parallel to the transect rope at a given depth and scoring a vertical line on the X-15 graph at each float thereby establishing a station for aquatic vegetation height and pond depth determinations. Field notes assisted interpretation of graph results in the lab. Macrophyte reductions were measured by changes in the vegetation height:pond depth ratio at each station. Common species

¹ Mimeo, Missouri Department of Conservation

Table 2. Pond name, group, code number from Figure 1, and comparison of surface areas (hectares) estimated with Missouri field chart and standard survey methods for 15 South Dakota study ponds between August, 1983, and August, 1984.

Pond Name	Group	code number	Surface Area	
			Estimated	Surveyed
Moore	control	1	0.73	0.56
Madsen	control	2	0.61	0.54
Frantz	control	3	0.81	0.63
Dailey	control	4	1.01	0.64
Heathon	control	5	1.42	0.84
Dailey	hybrid	6	1.21	1.70
Ridings	hybrid	7	1.42	1.80
Thompson	hybrid	8	1.42	1.32
Krull	hybrid	9	0.40	0.23
Willis	hybrid	10	1.01	no map
Pajl	monosex	11	1.21	1.36
Fox	monosex	12	1.26	1.03
Demray	monosex	13	0.73	0.60
Ortman	monosex	14	0.61	no map
Kippes	monosex	15	0.81	0.45

of aquatic macrophytes encountered in this study include Chara spp., Ceratophyllum demersum, Potamogeton spp., Ranunculus spp., and Utricularia vulgaris.

Fish populations were sampled in the spring, 1985, using a 39 m x 2.4 m seine with a mesh of 191 mm and a 240-volt electrofishing unit. Captured fish were anesthetized with MS-222 at a rate of 79 mg/liter in a small basin. Age-I largemouth bass were marked for population estimates by clipping the left pelvic fin. These fish were measured for total length (TL) to the nearest millimeter and total weight to the nearest gram. Largemouth bass scales were taken below the lateral line and just above the pectoral fin apex to determine growth to first annulus formation. Fish were dipped into a saturated salt solution and placed in a recovery tub prior to release into the pond.

Scale impressions were made on cellulose acetate strips with a roller press as outlined by Smith (1954). Impressions were viewed on a Bell and Howell microfiche projector. First annulus and scale radius were measured to the nearest millimeter from the scale focus.

Analysis

Population estimates were determined with the modified Peterson equation:

$$\hat{N} = \frac{(M + 1)(C + 1)}{(R + 1)} \quad (\text{Ricker 1975})$$

where,

$\hat{N}$ = estimated population number at time of marking

M = number of fish marked

C = catch or sample taken for census

R = number of recaptured marks in sample

The 95% confidence interval around the population estimate (Table 3) was calculated by the equation:

$$\hat{N} \pm 1.96 \sqrt{\hat{V}(\hat{N})} \quad (\text{Everhart and Young 1981})$$

where,

$\hat{V}(\hat{N})$ = sampling variance for N, calculated by the equation:

$$\hat{V}(\hat{N}) = \frac{N^2 (C - R)}{(C + 1)(R + 2)} \quad (\text{Everhart and Young 1981})$$

Survival rates were determined by dividing the population estimates by the number of fish stocked to study ponds in 1984 (Table 4).

Condition for age-I largemouth bass was determined using the index of relative weight (W_R):

$$W_R = \frac{W}{W_S} \times 100 \quad (\text{Wege and Anderson 1978})$$

where,

W_R = relative weight

W = actual weight of the fish

W_S = standard weight for a fish of a given length calculated by:

Table 3. Pond name, group, population estimate ($\hat{N}$), sampling variance $\hat{V}(\hat{N})$, and 95% confidence interval around $\hat{N}$ for age-I largemouth bass, Micropterus salmoides, from eight South Dakota study ponds sampled between April and October, 1985.

Pond Name	Group	$\hat{N}$	$\hat{V}(\hat{N})$	95% Confidence interval for N
Moore	control	19	14.4	11.6 - 26.4
Madsen	control	18	27.8	7.7 - 28.3
Frantz	control	141	4,418.0	10.7 - 271.3
Ridings	hybrid	85	2,890.0	-20.4 - 190.4
Thompson	hybrid	276	15,870.0	31.0 - 521.0
Pajl	monosex	77	174.0	51.1 - 102.9
Fox	monosex	12	36.0	0.2 - 23.8
Demray	monosex	141	1,729.0	59.0 - 223.0

Table 4. Pond name, group, population estimate, number of fish stocked in study ponds in 1984, and survival rate (percent) for age-I largemouth bass, Micropterus salmoides, from eight South Dakota study ponds sampled between 7 and 26 September, 1985.

Pond Name	Group	Population Estimate	Number Stocked	Survival (percent)
Moore	control	19	178	10.5
Madsen	control	18	150	11.7
Frantz	control	141	200	70.5
Ridings	hybrid	85	350	24.3
Thompson	hybrid	276	350	79.0
Pajl	monosex	77	300	26.0
Fox	monosex	12	310	3.9
Demray	monosex	141	180	78.5

$$\text{Log } W_s = -5.316 + 3.191 \text{ Log length}$$

First year growth for age-I largemouth bass was back calculated from scales using the corrected Lee formula:

$$L_n = \frac{a + S_n (L_c - a)}{S_c} \quad (\text{Carlander 1977})$$

where,

L_n = length of fish at time of annulus n formation

a = constant value of 22 for largemouth bass (Stone 1981)

S_n = Scale radius at annulus n

S_c = scale radius

L_c = length of fish at capture

Abundances of age-I bluegills, age-0 and age-I largemouth bass were determined by electrofishing shoreline perimeters and calculating the number of fish captured hourly (Table 5). Mouth widths for age-I largemouth bass and body depths for age-0 largemouth bass were determined from equations based on total length (Lawrence 1958) and were graphed to evaluate age-0 largemouth bass vulnerability to age-I largemouth bass predation in ponds with and without fathead minnows (Figure 2).

Nested factorial analysis of variance was used to determine differences in physical and chemical water quality variables sampled during 1984 between treatment and control ponds. Ponds within treatments designated as Pond (Trt) was the error term used to assess whether variability was greater within or between treatments. Nested factorial analysis of variance was also used to determine differences in water quality variables that may have been due to treatments, sessions,

Table 5. Pond name, group, number of fish electrofished per hour for age-I bluegills, Lepomis macrochirus, and age-0 and age-I largemouth bass, Micropterus salmoides, sampled from nine South Dakota study ponds between August and September, 1985.

Pond Name	Group	Bluegills Age-I	Largemouth Bass	
			Age-0	Age-I
Moore	control	36.92	26.54	4.62
Madsen	control	5.81	25.16	11.61
Frantz	control	5.45	0.00	6.82
Thompson	hybrid	5.81	3.87	16.45
Ridings	hybrid	15.29	12.94	4.71
Kippes	monosex	0.00	50.45	0.00
Pajl	monosex	8.94	17.87	20.43
Fox	monosex	0.94	0.94	10.31
Demray	monosex	6.38	7.66	28.09




Figure 2. Mouth widths of age-1 largemouth bass, Micropterus salmoides, in study ponds with (■) and without (□) fathead minnows, Pimephales promelas, compared to body depths of age-0 largemouth bass with (●) and without (○) fathead minnows, and initial body depth of age-0 largemouth bass (Δ) at pond stocking 10 July 1985.

and years. Differences in aquatic vegetation heights that may have been due to treatments, sessions, years, and transect water depth were assessed by nested factorial analysis of variance. All effects in the study were fixed. Waller-Duncan k-ratio t-tests were used to detect differences in first year growth and relative weight for largemouth bass in four ponds without fathead minnow contamination.

RESULTS

Pond Selection Evaluation

Water quality variables sampled during 1984 indicated no significant differences ($P > 0.05$) due to treatment effects for secchi disk visibility and maximum pond depth (Appendix Table 1), conductivity and salinity (Appendix Table 2), turbidity and pH (Appendix Table 3), phenol and total alkalinity (Appendix Table 4), total hardness and nitrate-nitrogen (Appendix Table 5), phosphate and sulfate (Appendix Table 6), potassium (Appendix Table 7), and pond surface levels (Appendix Table 8). Since no differences due to treatment effects were determined, study ponds were considered similar for purposes of data analyses.

Water Quality

Conductivity differed significantly ($P \leq 0.05$) due to treatments and to a treatment by year interaction (Table 6). Otherwise water quality variables did not differ significantly ($P > 0.05$) due to treatments likely because variability among ponds within treatments is large

Table 6. Nested factorial analysis of variance for differences in conductivity (micromhos/cm³) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	120,734.20	0.93
Trt*Year	2	561,661.05	4.32+
Year*Pond(Trt)	10	130,064.58	
Trt	2	7,612,223.00	4.04+
Pond(Trt)	11	1,886,441.50	
Session	2	72,052.88	0.97
Trt*Session	4	75,729.23	1.01
Session*Pond(Trt)	21	74,653.97	
Residual	20	99,130.33	

+Denotes significance ($P \leq 0.05$)

compared to variability between treatments. Salinity differed significantly ($P \leq 0.05$) between sessions (Table 7). Differences in total hardness were significant ($P \leq 0.01$) due to a treatment by year interaction (Table 8). Sulfates differed significantly ($P \leq 0.05$) for both years and a treatment by year interaction (Table 9). Potassium levels were significantly different ($P \leq 0.01$) between sessions (Table 10). Pond surface levels declined significantly ($P \leq 0.01$) for sessions with lowest levels recorded in the fall (Table 11). No differences in water transparency were significant ($P > 0.05$), although field observations and examination of graphed raw means of secchi disk visibility (Figure 3) and turbidity (Figure 4) suggested a trend despite the large variability. Remaining water quality variables indicated no significant differences ($P > 0.05$) due to treatments, sessions, and years for secchi disk visibility (Appendix Table 9), turbidity (Appendix Table 10), pH (Appendix Table 11), phenol alkalinity (Appendix Table 12), total alkalinity (Appendix Table 13), nitrate-nitrogen (Appendix Table 14), phosphate (Appendix Table 15), and maximum pond depth (Appendix Table 16).

Vegetation Removal

Aquatic vegetation heights were not significantly different ($P > 0.05$) because variability among ponds within groups exceeded differences between groups. Aquatic vegetation heights differed significantly ($P \leq 0.01$) between transect water depths (Table 12). Differences in aquatic vegetation heights were not significant ($P > 0.05$), although inspection of transect raw means sampled in 1.5–2.1 m (Figure 5) and

Table 7. Nested factorial analysis of variance for differences in salinity (parts per thousand) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	0.0004	0.00
Trt*Year	2	0.2279	1.88
Year*Pond(Trt)	10	0.1241	
Trt	2	3.2729	3.56
Pond(Trt)	11	0.9183	
Session	2	0.2159	4.27+
Trt*Session	4	0.0870	1.72
Session*Pond(Trt)	21	0.0506	
Residual	20	0.0436	

+ Denotes significance ($P \leq 0.05$)

Table 8. Nested factorial analysis of variance for differences in total hardness (mg/l as CaCO_3) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	110,022.45	3.82
Trt*Year	2	226,495.33	7.86++
Year*Pond(Trt)	10	28,826.10	
Trt	2	760,868.10	2.03
Pond(Trt)	11	375,325.31	
Session	2	23,529.71	2.11
Trt*Session	4	8,623.93	0.77
Session*Pond(Trt)	21	11,144.44	
Residual	20	9,820.25	

++Denotes significance ($P \leq 0.01$)

Table 9. Nested factorial analysis of variance for differences in sulfate (mg/l) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984 and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	170,246.39	5.25+
Trt*Year	2	176,041.01	5.43+
Year*Pond(Trt)	10	32,414.51	
Trt	2	1,461,597.50	2.66
Pond(Trt)	11	549,596.28	
Session	2	21,681.36	1.16
Trt*Session	4	29,648.06	1.58
Session*Pond(Trt)	21	18,718.50	
Residual	20	30,379.13	

+Denotes significance ($P \leq 0.05$)

Table 10. Nested factorial analysis of variance for differences in potassium (mg/l) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	96.15	1.63
Trt*Year	2	45.15	0.77
Year*Pond(Trt)	10	58.99	
Trt	2	16.14	0.03
Pond(Trt)	11	536.19	
Session	2	72.83	6.18++
Trt*Session	4	2.92	0.25
Session*Pond(Trt)	20	11.78	
Residual	19	5.22	

+Denotes significance ($P \leq 0.01$)

Table 11. Nested factorial analysis of variance for differences in pond surface levels (cm) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	1923.66	2.39
Trt*Year	2	77.19	0.10
Year*Pond(Trt)	7	803.90	
Trt	2	270.75	0.64
Pond(Trt)	11	423.10	
Session	2	1,559.37	10.32++
Trt*Session	4	262.21	1.73
Session*Pond(Trt)	18	151.14	
Residual	3	79.85	

++Denotes significance ($P \leq 0.01$)




Figure 3. Control and treatment means (horizontal bars) and standard error (rectangles) for secchi disk visibility sampled in 14 South Dakota study ponds between June, 1984, and August, 1985.




Figure 4. Control and treatment means (horizontal bars) and standard error (rectangles) for turbidity sampled in 14 South Dakota study ponds between June, 1984, and August, 1985.

Table 12. Nested factorial analysis of variance for differences in aquatic vegetation height (cm) due to treatments, sessions, years, and transect water depths sampled from 13 South Dakota study ponds between July, 1984 and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Years	1	7.0925	4.16
Trt*Years	2	0.4263	0.25
Years*Pond(Trt)	10	1.7033	
Trt	2	8.9349	0.56
Pond(Trt)	10	15.8427	
Session	2	0.2168	0.15
Trt*Session	4	0.5613	0.39
Session*Pond(Trt)	20	1.4450	
Years*Session	1	1.3418	5.38+
Trt*Years*Session	2	0.1178	0.47
Yrs*Sess*Pond(Trt)	8	0.2459	
TDepth	1	9.5231	13.98++
Trt*TDepth	2	0.9311	1.37
Pond*TDpth(Trt)	10	0.6812	
Years*TDepth	1	0.5838	1.15
Trt*Sess*TDepth	4	0.6405	0.01
Yrs*Pond*TDpth(Trt)	8	0.5068	
Session*TDepth	2	0.0439	0.12
Trt*Sess*TDepth	4	0.6405	1.74
Sess*Pond*TDpth(Trt)	17	0.3683	
Residual	3193	0.0474	

++Denotes significance ($P \leq 0.01$)

+ Denotes significance ($P \leq 0.05$)




Figure 5. Control and treatment raw means (horizontal bars) and standard deviations (rectangles) of aquatic vegetation heights sampled from 1.5 - 2.1 m transect water depths for control ponds (white), hybrid grass carp ponds (striped), and monosex grass carp ponds (black) for 13 South Dakota study ponds between July, 1984, and August, 1985.

2.4-3.0 m (Figure 6) water depths suggested trends corroborated by field observations that substantial macrophyte reductions occurred in the monosex grass carp treatment.

Fish Populations

The influence of macrophyte removal on bluegill vulnerability was undetected because of the failure of stocked bluegill to reproduce. Bluegill recruitment was never observed in any study pond in 1985. Abundance of age-I bluegills electrofished hourly was observed to be low (Table 5). Five study ponds were eliminated from analyses because of fish contamination. Additionally, one hybrid grass carp pond winterkilled in 1984 and, thus, was ignored in the analyses.

First year survival estimates for largemouth bass in eight ponds ranged from 0-79% with a mean of 38%. Survival did not appear to be associated with macrophyte removal (Table 4). Aquatic macrophytes retarded electrofishing efforts to recapture marked fish particularly in densely vegetated control ponds. Growth (Table 13) and W_T (Table 14) for age-I largemouth bass were compared from four ponds without fathead minnows. Abundance of age-I bluegills from these same four ponds was determined by electrofishing (Table 15). Abundance of age-0 largemouth bass in nine study ponds appeared to be influenced by the survival rate of age-I largemouth bass (Figure 7), however, no significant correlation was determined for this relationship. Growth and survival estimates for monosex grass carp and hybrid grass carp were undetermined because of low numbers of fishes captured in this study. Grass carp seined from two monosex grass carp ponds averaged 569 mm and 2,390 g for 14 fish




Figure 6. Control and treatment raw means (horizontal bars) and standard deviations (rectangles) of aquatic vegetation heights sampled from 2.4 - 3.0 m transect water depths for control ponds (white), hybrid grass carp ponds (striped), and monosex grass carp ponds (black) for 13 South Dakota study ponds between July, 1984, and August, 1985.

Table 13. Waller-Duncan K-ratio t-test to determine differences in first year growth of largemouth bass, Micropterus salmoides, sampled in four study ponds without fathead minnows between April and June, 1985.

FIRST YEAR GROWTH			
Pajl	Frantz	Moore	Madsen
monosex	control	control	control
148.58	158.29	164.44	68.59
	*		
*			

* Horizontal lines indicate similar growth for largemouth bass.

Table 14. Waller-Duncan K-ratio t-test to determine differences in W_t of age-I largemouth bass, Micropterus salmoides, sampled in four study ponds without fathead minnows between April and June, 1985.

RELATIVE WEIGHT			
Madsen	Moore	Frantz	Pajl
control	control	control	monosex
101.06	121.11	127.21	127.64
	* _____		

* Horizontal line indicates similar relative weight for largemouth bass.

Table 15. Pond name, sampling date, number of fish electrofished per hour, and (TL) (mm) for age-I bluegills, Lepomis macrochirus, from four study ponds without fathead minnows between May and September, 1985.

Pond Name	Date	number of bluegill electrofished per hour	TL (mm)
Madsen	6/15	0.80	
	8/02	0.98	
	9/21	5.81	107.3
Frantz	8/02	14.03	
	9/21	5.45	130.8
Moore	5/26	0.92	61.0
	6/14	4.80	74.0
	6/15	16.92	
	8/03	24.00	107.2
	9/20	36.92	130.7
Pajl	6/16	18.0	
	8/02	32.9	104.0
	9/21	8.94	120.9




Figure 7. Second order regression of the abundance of age-0 largemouth bass, *Micropterus salmoides*, and survival rate of age-1 largemouth bass from three control ponds (C), two hybrid grass carp ponds (H), and four monosex grass carp ponds (M) sampled between 17 and 30 September 1985.

from one pond, and 520 mm and 1,714 g for six fish in the other pond in the fall, 1985.

DISCUSSION

Water Quality

Specific influences of herbivorous fishes on water quality are difficult to summarize because of inconsistent responses reported in the literature (Bailey 1978; Mitzner 1978; Canfield et al. 1983; Mitchell et al. 1984). Hickling (1966) stated that grass carp digestion reduced plant matter into fragments of 3 mm² or less. Later Opuszynski (1972) speculated that pulverized grass carp excreta provides nutrients to phytoplankton, often resulting in algal blooms. Gasaway and Drda (1978) noted increases in chlorophyll concentrations and reductions in phytoplankton species diversity following macrophyte removal by grass carp in Florida ponds. Avault et al. (1968) observed the formation of a brown organic scum and a 300% increase in free potassium levels following macrophyte removal by grass carp in research pools. However, Terrell (1982) suggested that nutrients released from grass carp feces are quickly bound in the bottom sediments and are unavailable to phytoplankton in softwater acidic ponds in Georgia. Other studies observed no changes in algal densities (Lembi et al. 1978; Mitzner 1978), whereas some studies noted zooplankton increases following grass carp introductions (Ritenour 1976; Mitchell et al. 1984; Richard et al. 1985).

In the present study, water transparency appeared to decline greatest in the monosex grass carp treatment. Rapid macrophyte removal and subsequent increased pond mixing may explain the reduced secchi disk visibility and the increased turbidity observed in the present study.

Wiley and Gordon (1984) suggested that the influence of herbivorous fishes on aquatic systems depends on stocking density. Perhaps the removal rate of aquatic macrophytes ultimately dictates the impact of grass carp introductions on water quality.

Vegetation Control

Several studies have documented that grass carp are efficient in removing aquatic macrophytes (e.g. Ware and Gasaway 1976; Mitzner 1978; Bailey 1978; Lembi et al. 1978; Mitchell et al. 1984; Fowler 1985). However, no significant ($P > 0.05$) reductions in aquatic macrophyte heights between treatment and control ponds were detected in the present study. Most likely, this was because pond variability within groups was large compared to variability between pond groups. Nevertheless, observation suggested that aquatic macrophytes were greatly reduced in the monosex grass carp treatment with slight differences visible between control and hybrid grass carp ponds. Aquatic macrophyte removal by hybrid grass carp was not apparent in the present study. Higher stocking rates for hybrid grass carp are recommended for vegetation control (Shireman 1982; Osborn 1982; Young et al. 1983; Harberg and Modde 1985), although Osborne (1985) concluded that stocking hybrid grass carp to control hydrilla (Hydrilla verticillata) in Florida was not economically feasible.

Survival of both herbivorous fishes appeared to be low in the present study. Several factors could account for this. Mitzner (1978) cited hauling stress as a major cause of mortality for grass carp stocked into Red Haw Lake, Iowa. The monosex grass carp and hybrid

grass carp used in this study were obtained in Arkansas and endured hauling for 30-50 hours prior to pond stocking. Colle et al. (1978) implicated piscivorous birds as one factor to explain 95% mortality of 5000 grass carp with mean TL of 48 mm stocked to a vegetated Florida pond devoid of piscivorous fishes. This speculation warrants consideration for ponds in the present study since the mean stocking length of monosex grass carp was 63.2 mm, and piscivorous birds were often observed near study ponds. Harberg and Modde (1985) reported survival rates of 59.1% and 32.2% for hybrid grass carp stocked into two western South Dakota ponds after one growing season. Hybrid grass carp were observed to be less numerous in ponds than monosex grass carp in the present study. Hardin and Atterson (1984) observed 98% mortality for grass carp stocked into Lake Wales, Florida, attributing mortality initially to largemouth bass predation and later to other unknown causes. Therefore, stocking length, survival rate, and piscivorous predator abundance are important factors to consider in determining adequate stocking rates of herbivorous fishes for vegetation control.

Fish Populations

The influence of declining cover on the largemouth bass-bluegill relationship was not determined in the present study. Studies by Ball (1977) and Baur et al. (1979) demonstrated that grass carp stocked into shallow hatchery ponds decreased gamefish production only when complete macrophyte removal resulted. Ball (1977) stated that the production estimates for gamefish in ponds exhibited so much variation that it was impossible to isolate the effects of grass carp introductions into

Indiana hatchery ponds. Baur et al. (1979) reported that adverse effects on gamefish in hatchery ponds were less likely when grass carp stocking rates were less than 30 fish/hectare. Wiley and Gordon (1984) reported that declines in centrarchid young-of-the-year production were greatest in ponds overstocked with hybrid grass carp. Bailey (1978) stated that the impact of grass carp introductions on fish populations in 31 Arkansas lakes were negligible but vegetation removal did appear to improve the conditions of largemouth bass, bluegills, and redear sunfish, Lepomis microlophus. Colle and Shireman (1980) observed that the conditions for bluegills, redears, and largemouth bass decreased as hydrilla density increased in the water column in Lake Wales, Florida. Shireman and Maceina (1981) reported low condition factors for largemouth bass > 250 mm TL when hydrilla coverage exceeded 30%, whereas smaller largemouth bass were not affected until coverage surpassed 50% in laboratory studies.

Effects of vegetation removal on survival, growth, and relative weight of age-I largemouth bass were not determined in the present study. Mean survival of 38% for age-I largemouth bass was lower than the 50% value reported by Stone (1981) and the 59% value observed by Beck (1986) for South Dakota ponds stocked with various fish combinations. Mean first year TL of age-I largemouth bass from four ponds without fathead minnows was 160 mm, which is similar to the 153 mm TL recorded by Stone (1981) in 16 eastern South Dakota ponds. Mean W_T for age-I largemouth bass from the same four ponds was 119 g, which exceeded the value of 114 g noted by Stone (1981) in South Dakota ponds. Wege and Anderson (1978) demonstrated that W_T of largemouth bass of less

than 203 mm TL and between 203-302 mm TL were significantly correlated ($r=.60$ and $r=.58$, respectively) to the log density of fish prey smaller than 76 mm TL, suggesting the importance of small prey to small largemouth bass. Age-I largemouth bass in Madsen's pond had the highest mean growth of 168.6 mm but the lowest W_r of 101, possibly reflecting a reduced forage base since this pond in particular was noted to have the lowest bluegill abundance among these four ponds.

Bluegill survival was presumably affected by the mean stocking length of 31.7 mm in 1984 compared to the mean length of 50.0 mm stocked by Gilbraith (personal communication) in 1983. Reduced stocking length potentially increased bluegill vulnerability to the age-0 largemouth bass stocked into all study ponds in July, 1984. Werner et al. (1983) reported that bluegill juveniles remain in vegetation to avoid largemouth bass predation at the expense of optimal foraging in the water column and hence their growth is reduced. Small stocking size and constant vulnerability to largemouth bass predation may have accounted for low bluegill survival in the present study.

Reynolds and Babb (1978) found high survival of age-0 largemouth bass in ponds with 50% of the surface vegetated regardless of the adult largemouth density. However, Durocher et al. (1984) determined no relationship between percent cover of submergent vegetation and the abundance of small largemouth bass in 30 Texas reservoirs. The survival rate of age-I largemouth bass appeared to influence age-0 largemouth bass abundance in the present study. Age-I largemouth bass may have preyed upon age-0 largemouth bass in the absence of small bluegills. Comparison of age-0 largemouth bass body depths with age-I largemouth

bass mouth widths indicated that age-0 largemouth bass were vulnerable to age-I largemouth bass predation in the present study. Fathead minnow presence appeared to reduce the vulnerability of age-0 largemouth bass to age-I fish toward the end of the study.

Monosex grass carp were observed to remove aquatic macrophytes rapidly in the present study. Decreased water transparency may occur following grass carp introductions. Declines in water transparency in the present study likely resulted from overstocking monosex grass carp. The influence of declining cover on bluegill vulnerability was not evident in the present study. Further research is necessary to ascertain if cover manipulations are a feasible means to enhance bluegill vulnerability to largemouth bass predation.

RECOMMENDATIONS

1. Monosex grass carp appear to be an effective means to reduce aquatic macrophytes in prairie ponds. Subsequent changes in water quality will likely depend on the stocking density.
2. Grass carp stocking rates should be based on the following:
vegetated area to be controlled, stocking length, aquatic plant preference of grass carp, and the timetable to achieve a specific level of control. Neither complete macrophyte control nor a rapid removal rate are recommended in order to minimize undesirable changes in water transparency.
3. Grass carp probably should be stocked in the summer months to maximize growth and survival and to be of sufficient size to elude predation by piscivorous birds.

LITERATURE CITED

- Avault, J. W., Jr., R. W. Smitherman, and E. W. Shell. 1968.
Evaluation of eight species of fish for aquatic weed control. Food
and Agriculture Organization of the United Nations, Report 44, VII
E-3:109-122.
- Bailey, W. M. 1978. A comparison of fish populations before and after ✓
extensive grass carp stocking. Transactions of the American
Fisheries Society 107:181-206.
- Ball, R. L. 1977. Effects of the white amur on largemouth bass- ✓
bluegills-redear sunfish populations. Indiana Department of Natural
Resources, Progress Report for 1976, West Lafayette, Indiana, USA.
- Baur, R. J., D. H. Buck, and C. R. Rose. 1979. Production of age-0 ✓
largemouth bass, smallmouth bass, and bluegills in ponds stocked
with grass carp. Transactions of the American Fisheries Society
108:496-498.
- Beck, R. D. 1986. Growth, survival, and reproductive success of
largemouth bass stocked with selected forage fishes in South Dakota
ponds. Masters thesis. South Dakota State University, Brookings,
South Dakota, USA.
- Boney, S. E., W. L. Shelton, S. L. Yang, and L. O. Wilken. 1984. Sex ✓
reversal and breeding of grass carp. Transactions of the American
Fisheries Society 113:348-353.
- Canfield, D. E., Jr., M. J. Maciena, and J. V. Shireman. 1983. Effects
of hydrilla and grass carp on water quality in a Florida lake.
Water Resources Bulletin. 19:773-778.

- Carlander, K. D. 1977. Handbook of freshwater fishery biology, volume 2. The Iowa State University Press, Ames, Iowa, USA.
- Colle, D. E., J. V. Shireman, and R. W. Rottman. 1978. Food selection by grass carp fingerlings in a vegetated pond. Transactions of the American Fisheries Society 107:149-152.
- Colle, D. E., and J. V. Shireman. 1980. Coefficients of condition for largemouth bass, bluegills, and redear sunfish in hydrilla-infested lakes. Transactions of the American Fisheries Society 109:521-531.
- Conner, J. V., R. P. Gallagher, and M. F. Charty. 1980. Larval evidence for natural reproduction of the grass carp (Ctenopharyngodon idella) in the lower Mississippi River. Pages 1-19 in L. A. Fuiman, editor. Proceedings of the fourth annual larval fish conference. FWS/OBS-80-43. U.S. Fish and Wildlife Service, Washington, D.C., USA.
- Cross, D. G. 1969. Aquatic weed control using grass carp. Journal of Fish Biology 1:27-30.
- Crowder, L. B., and W. E. Cooper. 1979. Structural complexity and fish-prey interactions in ponds: a point of view. Pages 2-10 in D. L. Johnson and R. A. Stein, editors. Response of fish to habitat structure in standing water. North Central Division, American Fisheries Society, Special Publication 6, Bethesda, Maryland, USA.
- Dillard, J. G., and G. D. Novinger. 1975. Stocking largemouth bass in small impoundments. Pages 459-474 in H. Clepper, editor. Black bass biology and management. Sport Fishing Institute, Washington, D.C., USA.
- Durocher, P. P., W. C. Provine, and J. E. Kraai. 1984. Relationship

- between abundance of largemouth bass and submerged vegetation in Texas reservoirs. North American Journal of Fisheries Management 4:84-88.
- Everhart, W. H. and W. D. Young. 1981. Principles of fishery science, 2nd edition. Cornell University Press. Ithaca, New York, USA.
- Fowler, M. C. 1985. The results of introducing grass carp, Ctenopharyngodon idella Val., into small lakes. Aquaculture and Fisheries Management 16:189-201.
- Gasaway, R. D., and T. F. Drda. 1977. Effects of grass carp introduction on waterfowl habitat. Proceedings of the Forty-Second North American Wildlife Conference 42:73-85. ✓
- Gasaway, R. D., and T. F. Drda. 1978. Effects of grass carp introduction on macrophytic vegetation and chlorophyll content of phytoplankton in four Florida lakes. Florida Scientist 41:100-109.
- Glass, N. R. 1971. Computer analysis of predation energetics in the largemouth bass. Pages 325-363 in B. C. Patten, editor. Systems analysis and simulation ecology, volume 1. Academic Press, New York, New York, USA.
- Guillory, V., and R. D. Gasaway. 1978. Zoogeography of the grass carp in the United States. Transactions of the American Fisheries Society 107:105-112.
- Harberg, M. C., and T. Modde. 1985. Feeding behavior, food consumption, growth, and survival of hybrid grass carp in two South Dakota ponds. North American Journal of Fisheries Management 5:457-464
- Hardin, S., and J. Atterson. 1984. Ecological effects of grass carp

- introduction in Lake Wales, Florida. Game and Freshwater Fish Commission Final Report, Florida.
- Hickling, C. F. 1966. On the feeding process in the white amur, Ctenopharyngodon idella. Journal of Zoology 148:408-419.
- Hill, K. 1980. Evaluation of the split stocking method in Iowa farm ponds. Iowa Conservation Commission, Dingell-Johnson Project F-90-R, Study number 403.1, Ames, Iowa, USA.
- Jensen, G. L., W. L. Shelton, S. L. Yang, and L. O. Wilken. 1983. Sex reversal of gynogenetic grass carp by implantation of methyltestosterone. Transactions of the American Fisheries Society 112:79-85.
- Lawrence, J. M. 1958. Estimated sizes of various forage fishes largemouth bass can swallow. Proceedings of the Annual Conference, Southeastern Association of Game and Fish Commissioners 11:220-225.
- Lazareva, L. P., M. O. Omarov, and A. N. Leznia. 1977. Feeding and growth of the bighead, Aristichthys nobilis, in the waters of Dagestan. Journal of Ichthyology (USSR) 17:65-71.
- Lembi, C. A., B. G. Ritenour, E. M. Iverson, and E. C. Forss. 1978. The effects of vegetation removal by grass carp on water chemistry and phytoplankton in Indiana ponds. Transactions of the American Fisheries Society 107:161-171.
- Magee, S. M., and D. P. Philipp. 1982. Biochemical genetic analyses of the grass carp female x bighead male F-1 hybrid and the parental species. Transactions of the American Fisheries Society 111:593-602.
- Marian, T., and Z. Krasznai. 1978. Karyological investigations on

- Ctenopharyngodon idella and Hypophthalmichthys nobilis and their cross breeding. *Aquaculture Hungarica* (Szarvas) 1:44-50.
- Mitchell, C. P., G. R. Fish, and A. M. R. Burnet. 1984. Limnological changes in a small lake stocked with grass carp. *New Zealand Journal of Marine and Freshwater Research* 18:103-114.
- Mittlebach, G. G. 1981. Foraging efficiency and body size: a study of optimal diet and habitat use by bluegills. *Ecology* 62:1370-1386.
- Mitzner, L. 1978. Evaluation of biological control of nuisance aquatic vegetation by grass carp. *Transactions of the American Fisheries Society* 107:135-145.
- Modde, T., and C. G. Scalet. 1985. Latitudinal growth effects on predator-prey interactions between largemouth bass and bluegills in ponds. *North American Journal of Fisheries Management* 5:457-464.
- Modde, T., and C. G. Scalet. 1986. Latitudinal influences upon largemouth bass and bluegill interactions in small impoundments in R. H. Stroud, editor. *Role of Culture in Fishery Management*. American Fisheries Society, Special Publication.
- Opuszynski, K. 1972. Use of phytophagous fish to control aquatic plants. *Aquaculture* 1:61-74.
- Osborne, J. A. 1982. The potential of the hybrid grass carp as a weed control agent. *Journal of Freshwater Ecology* 1:353-360.
- Osborne, J. A. 1985. A preliminary study of the efficacy of hybrid grass carp for hydrilla control. *Journal of Aquatic Plant Management* 23:16-20.
- Reynolds, J. B., and L. R. Babb. 1978. Structure and dynamics of largemouth bass populations. Pages 50-61 in G. D. Novinger and J.

- G. Dillard, editors. New Approaches to the management of small impoundments. Special Publication Number 5, North Central Division, American Fisheries Society, Bethesda, Maryland, USA.
- Richard, D. I., J. W. Small, and J. A. Osborne. 1985. Response of zooplankton to the reduction and elimination of submerged vegetation by grass carp and herbicide in four Florida lakes. *Hydrobiologia* 123:97-108.
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. Bulletin 191, Fisheries Research Board of Canada, Ottawa, Quebec, Canada.
- Ritenour, B. G. 1976. Effects of the white amur, (Ctenopharyngodon idella Val.) on small impoundments in southern Indiana. Masters thesis, Purdue University, West Lafayette, Indiana, USA.
- Rottman, R. W., and R. O. Anderson. 1977. Limnological and ecological effects of grass carp in ponds. Proceedings of the Annual Conference, Southeastern Game and Fish Commissioners 30:24-39.
- Savino, J. F., and R. A. Stein. 1982. Predator-prey interaction between largemouth bass and bluegills as influenced by simulated submersed vegetation. Transactions of the American Fisheries Society 111:255-266.
- Shireman, J. V. 1982. Cost analysis of aquatic weed control: fish versus chemicals in a Florida lake. The Progressive Fish Culturist 44:199-200.
- Shireman, J. V., and M. J. Maceina. 1981. The utilization of grass carp, Ctenopharyngodon idella Val., for hydrilla control in Lake Baldwin, Florida. Journal of Fish Biology 19:629-636.

- Smith, S. H. 1954. A method of producing plastic impressions of fish scales without using heat. *The Progressive Fish Culturist* 16:75-78.
- Spuhler, W. E., W. F. Lytle, and D. Moe. 1971. The climate of South Dakota. Bulletin 582. South Dakota Agricultural Experiment Station, South Dakota State University, Brookings, South Dakota, USA.
- Stone, C. C. 1981. Growth and survival of largemouth bass in new stocked South Dakota impoundments. Masters thesis, South Dakota State University, Brookings, South Dakota, USA.
- Swingle, H. S., and E. V. Smith. 1941. Experiments on the stocking of fish ponds. *Transactions of the Fifth North American Wildlife Conference* 5:267-276.
- Terrell, T. T. 1982. Responses of plankton communities to the introduction of grass carp into some Georgia ponds. *Journal of Freshwater Ecology* 1:395-406.
- Vinogradov, V. K., and Z. K. Zolotova. 1974. The influence of grass carp on aquatic ecosystems. *Hydrobiological Journal* 10:72-78.
- Ware, F. J., and R. D. Gasaway. 1976. Effects of grass carp on fish populations in two Florida lakes. *Proceedings of the Annual Conference in the Southeastern Association of Game and Fish Commissioners* 30:324-335.
- Wege, G. J., and R. O. Anderson. 1978. Relative weight (W_r): a new index of condition for largemouth bass. Pages 79-91 in G. D. Novinger and J. G. Dillard, editors. *New Approaches to the Management of small impoundments*. Special Publication 5. North Central Division, American Fisheries Society, Bethesda, Maryland,

USA.

Werner, E. E., J. F. Gilliam, D. J. Hall, and G. G. Mittlebach. 1983.

An experimental test of the effects of predation risk on habitat use in fish. *Ecology* 64:1540-1548.

Wiley, M. J., and R. W. Gordon. 1984. Biological control of aquatic macrophytes by herbivorous carp. Aquatic Biology Technical Report (10), Illinois Natural History Survey, Federal Aid Project F-37R, Urbana, Illinois, USA.

Young, L. M., J. P. Monaghan, Jr., and R. C. Heidinger. 1983. Food preferences, food intake, and growth of the F-1 hybrid of grass carp female x bighead male. *Transactions of the American Fisheries Society* 112:661-664.

Appendix

Table 1. Nested factorial analysis of variance for differences in secchi disk visibility (m) and maximum pond depth (cm) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt).

Source	df	Secchi MS	MaxDepth MS
Session	2	0.1869	4,013.39
Treatment	2	1.8020	6,031.77
Sess*Trt	4	0.0862	3,296.55
Pond(Trt)	12	2.0547	11,155.16
Residual	21	0.4087	1,734.91

Appendix

Table 2. Nested factorial analysis of variance for differences in conductivity (micromhos/cm³) and salinity (parts/thousand) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt).

Source	df	Conduct MS	Salinity MS
Session	2	571,785.13	0.2451
Treatment	2	3,099,743.40	1.3806
Sess*Trt	4	167,388.81	0.1026
Pond(Trt)	12	1,045,278.00	0.6557
Residual	23	139,309.56	0.0737

Appendix

Table 3. Nested factorial analysis of variance for differences in turbidity (formazin turbidity units) and pH (standard units) due to treatments and sessions sampled from 15 South Dakota study ponds between June, 1984 and August, 1985. Error terms used for treatment F-tests indicated by (Trt).

Source	df	turbidity MS	pH MS
Session	2	303.20	1.3942
Treatment	2	1,674.94	0.3580
Session*Trt	4	110.49	0.0230
Pond(Trt)	12	1,166.97	1.1158
Residual	23	225.20	0.1512

Appendix

Table 4. Nested factorial analysis of variance for differences in phenol and total alkalinity (mg/l as CaCO_3) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated (Trt).

Source	df	Phenol Alka MS	Total Alka MS
Session	2	123.62	1,887.64
Treatment	2	307.09	1,272.51
Sess*Trt	4	190.69	1,255.43
Pond(Trt)	12	1,287.29	14,455.99
Residual	23	139.67	320.35

Appendix

Table 5. Nested factorial analysis of variance for differences in total hardness (mg/l as CaCO_3) and nitrate-nitrogen (mg/l) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt).

Source	df	Hardness MS	Nitrate MS
Session	2	36,787.16	0.2216
Treatment	2	185,761.77	0.6612
Sess*Trt	4	10,808.34	0.8125
Pond(Trt)	12	167,752.52	0.5552
Residual	23	17,593.03	0.3146

Appendix

Table 6. Nested factorial analysis of variance for differences in phosphate (mg/l) and sulfate (mg/l) due to treatments, and session sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt).

Source	df	Phosphate MS	Sulfate MS
Session	2	1.5350	397.08
Treatment	2	0.1518	659,189.40
Sess*Trt	4	0.1514	1,416.19
Pond(Trt)	12	0.2823	344,152.44
Residual	23	0.2150	1,123.87

Appendix

Table 7. Nested factorial analysis of variance for differences in potassium (mg/l) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt).

Source	df	Potassium MS
Session	2	54.56
Treatment	2	46.58
Sess*Trt	4	1.42
Pond(Trt)	12	406.35
Residual	22	11.06

Appendix

Table 8. Nested factorial analysis of variance for differences in pond surface levels (cm) due to treatments and sessions sampled from 15 South Dakota study ponds between June and August, 1984. Error terms used for treatment F-tests indicated by (Trt).

Source	df	Pond Surface level MS
Session	2	2,465.99
Treatment	2	220.74
Sess*Trt	4	81.91
Pond(Trt)	12	308.08
Residual	17	112.25

Appendix

Table 9. Nested factorial analysis of variance for differences in secchi disk visibility (m) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	1.1883	1.98
Trt*Year	2	0.0743	0.12
Year*Pond(Trt)	10	0.5994	
Treatment	2	2.7755	1.20
Pond(Trt)	11	2.3137	
Session	2	0.7577	1.59
Trt*Session	4	0.0542	0.11
Session*Pond(Trt)	21	0.4772	
Residual	18	0.2702	

Appendix

Table 10. Nested factorial analysis of variance for differences in turbidity (formazin turbidity units) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	593.85	3.65
Trt*Year	2	150.74	0.93
Year*Pond (Trt)	10	162.80	
Treatment	2	3,200.72	2.31
Pond (Trt)	11	1,383.69	
Session	2	338.09	1.36
Trt*Session	4	63.71	0.62
Session*Pond (Trt)	21	247.89	
Residual	20	85.58	

Appendix

Table 11. Nested factorial analysis of variance for differences in pH (standard units) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	0.2856	0.39
Trt*Year	2	0.3192	0.43
Year*Pond (Trt)	10	0.7370	
Treatment	2	1.0689	0.83
Pond (Trt)	11	1.2937	
Session	2	0.2111	0.57
Trt*Session	4	0.2066	0.56
Session*Pond (Trt)	21	0.3708	
Residual	20	0.2581	

Appendix

Table 12. Nested factorial analysis of variance for differences in phenol alkalinity (mg/l as CaCO_3) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	209.25	0.25
Trt*Year	2	877.50	1.03
Year*Pond (Trt)	10	847.83	
Treatment	2	5.63	0.00
Pond (Trt)	11	1,485.32	
Session	2	108.95	0.61
Trt*Session	4	202.98	1.14
Session*Pond (Trt)	21	177.82	
Residual	20	175.58	

Appendix

Table 13. Nested factorial analysis of variance for differences in total alkalinity (mg/l as CaCO_3) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	3,546.30	3.12
Trt*Year	2	1,036.24	0.91
Year*Pond (Trt)	10	1,135.96	
Treatment	2	3,389.35	0.14
Pond (Trt)	11	23,658.69	
Session	2	502.07	1.04
Trt*Session	4	1,323.82	2.73
Session*Pond (Trt)	21	484.14	
Residual	20	404.40	

Appendix

Table 14. Nested factorial analysis of variance for differences in nitrate-nitrogen (mg/l) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	0.0326	0.13
Trt*Year	2	0.3334	1.28
Year*Pond (Trt)	10	0.2605	
Treatment	2	1.3023	2.76
Pond (Trt)	11	0.4714	
Session	2	0.1065	0.44
Trt*Session	4	0.4669	1.95
Session*Pond (Trt)	21	0.2395	
Residual	20	0.1966	

Appendix

Table 15. Nested factorial analysis of variance for differences in phosphate (mg/l) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	0.6451	2.28
Trt*Year	2	0.0932	0.33
Year*Pond (Trt)	10	0.2831	
Treatment	2	0.0722	0.23
Pond (Trt)	11	0.3094	
Session	2	0.2406	1.38
Trt*Session	4	0.0736	0.42
Session*Pond (Trt)	21	0.1739	
Residual	20	0.1327	

Appendix

Table 16. Nested factorial analysis of variance for differences in maximum pond depth (cm) due to treatments, sessions, and years sampled from 14 South Dakota study ponds between June, 1984, and August, 1985. Error terms used for appropriate F-tests indicated by (Trt).

Source	df	MS	F
Year	1	1,980.27	0.58
Trt*Year	2	1,698.27	0.50
Year*Pond (Trt)	10	3,387.43	
Treatment	2	6,021.43	0.28
Pond (Trt)	11	21,810.69	
Session	2	970.58	0.70
Trt*Session	4	2,660.54	1.91
Session*Pond (Trt)	21	1,389.63	
Residual	18	684.01	

