# FISH COMMUNITIES AND HABITATS IN THIRTEEN MILE CREEK, DAWSON COUNTY, MONTANA

Mark D. Barnes Department of Natural Resources, Chinese Culture University Hwa Kang, Yang Ming Shan, Taipei, Taiwan 11192

> Steven D. Lee 215 3rd Street Glendive, Montana 59330

Ye Yan-Xi Department of Natural Resources, Chinese Culture University Hwa Kang, Yang Ming Shan, Taipei, Taiwan 11192

May 2009



Montana Department of Fish, Wildlife & Parks Miles City, Montana 59301

## FISH COMMUNITIES AND HABITATS IN THIRTEEN MILE CREEK, DAWSON COUNTY, MONTANA

## **INTRODUCTION**

Until recently little research has inquired into the distribution, habitat requirements, and ecological relationships of fishes in small prairie streams in eastern Montana. Clancey (1978) conducted a study of longitudinal distribution of fishes and aquatic habitats in Sarpy Creek, Big Horn and Treasure counties, to provide baseline information for the assessment of potential impacts of coal mining in the area. Barfoot (1993, 1999) studied longitudinal distribution of fishes and aquatic habitats in Little Beaver Creek, Carter and Fallon counties, Montana, and Bowman and Slope counties, North Dakota, in order to test the hypothesis that longitudinal zonation of fishe communities in streams is related primarily to changes in stream geomorphology.

Several wider surveys have sought to develop baseline data on fish communities and habitats in eastern Montana in order to assess the potential impacts of future energy development, particularly strip mining of coal and on-site power generation, on these communities. The results of these studies were collected and summarized by Elser et al. (1980).

Of these extensive surveys, two generated data on fish communities in smaller prairie streams. Elser et al. (1978) conducted an inventory of fishes and aquatic habitats in Beaver Creek, three of its tributaries, and seven north-flowing tributaries of the lower Yellowstone River. Morris et al. (1981) conducted a similar inventory of 45 tributaries of the lower Yellowstone River and assigned value ratings to each stream based on habitat and species value and recreational fishery potential.

More recently the Montana Department of Fish, Wildlife and Parks and the Montana Cooperative Fish and Wildlife Research Unit initiated extensive surveys of Montana prairie streams covering 305 sites in 240 drainages east of the Rocky Mountain region (Jones-Wuellner and Bramblett 2004, McDonald 2003). Bramblett et al. (2004) also developed a multimetric index of biological integrity for Montana prairie streams using fish assemblages.

Given the potential impacts of current land uses, including livestock grazing and irrigated agriculture, and of future energy development, including coal bed methane production, on stream fish communities in eastern Montana, it would be desirable to learn more about the distribution and habitat requirements of stream fishes in the region. Moreover, from the perspective of basic ecological research, it would be interesting to further explore factors that influence longitudinal zonation of fish communities in small prairie streams.

From 1997 to 2000, Barnes (1997, 1999), Barnes and Westlind (2000), and Barnes and Silbernagle (2001) conducted a quantitative study of fish population densities, longitudinal distribution, and habitat variables in Burns Creek, a perennial groundwater driven stream in Dawson and Richland counties, Montana. Results of this study suggested that several persistent habitat factors as well as random contingent phenomena both strongly influenced density and longitudinal distribution of fishes in that system. The most important persistent factors were stream width-depth ratio, riparian vegetation height and overhang, and instream cover, whereas the most important contingent phenomena were storm flow flushing and changing patterns in number

and location of beaver (Castor canadensis) dams.

Because it is perennial and groundwater driven, Burns Creek may not typify small prairie streams in eastern Montana, most of which are runoff driven and highly seasonal. Therefore, during the summers of 2001, 2002, and 2003, the sampling methodology developed for Burns Creek was applied to O'Fallon Creek, a largely runoff driven tributary of the lower Yellowstone River in Prairie, Custer, Fallon, and Carter counties, Montana (Barnes et al. 2002, Barnes and Siegle 2003, Barnes 2004). Results of the O'Fallon Creek studies suggested that contingent phenomena may more strongly influence density and longitudinal distribution of fishes in that system than more persistent habitat factors such as stream width-depth ratio and riparian vegetation. In this intermittent system, fish moving upstream or downstream may be trapped, often in high densities, in shrinking pools with low quality habitat (low dissolved oxygen concentrations, high total suspended solids concentrations, relatively shallow depths, and lack of riparian or instream vegetation) as riffles dry up and interrupt the hydrologic continuity of the stream. Similarly, drying riffles may exclude fish from preferred habitats. The overall "flashiness" of the system probably prevented fish communities from stabilizing and efficiently partitioning available habitats.

During the summers of 2004, 2005, and 2006, we applied the sampling methodology developed for Burns Creek and O'Fallon Creek to Upper Sevenmile Creek, an intermittent tributary of the Yellowstone River near Glendive, Dawson County, Montana (Barnes et al. 2005, Barnes et al. 2006, Barnes et al. 2007), and during the summers of 2007 and 2008 to Thirteen Mile Creek, an intermittent tributary of the Yellowstone River near Intake, Dawson County, Montana (Barnes et al. 2008). This report summarizes the results of the study in Thirteen Mile Creek during 2008.

#### **STUDY AREA**

Thirteen Mile Creek originates in northwestern Dawson County, Montana, and flows southeast 80.8 km to its confluence with the Yellowstone River at river kilometer 116 near Intake (Figures 1 and 2a). Channel elevation ranges from 945.1 m at the main stem headwaters to 610.0 m at the mouth, with a mean gradient of 4.1 m/km. Thirteen Mile Creek is a third order stream (Strahler 1952) with four major intermittent tributaries: (1) South Fork (Lower) Thirteen Mile Creek (Figure 2b) with confluence at 19.2 km; (2) North Fork Thirteen Mile with confluence at 37.2 km; (2) Center Fork Thirteen Mile Creek with confluence at kilometer 51.08; and South Fork (Upper) Thirteen Mile Creek with confluence at 58.8 km. Total channel length (all stream orders) of the system is 183.2 km. The whole system drains 414.8 km<sup>2</sup> and has a drainage density of 0.4 km/km<sup>2</sup> (MFIS 2009, Morris et al. 1981, NRIS 2009).

The Thirteen Mile Creek watershed occupies the edge of the glaciated Missouri Plateau, Short Grass Region, Great Plains Province. The climate is semiarid continental, with mean maximum temperature of 13.9°C and mean minimum temperature of -0.6°C (daily extremes from -47.2°C on 16 February 1936 to 43.8°C on 6 July 1934), mean annual precipitation of 34.8 cm (annual range from 15.1 cm in 1934 to 68.9 cm in 1906), and mean annual snowfall of 85.1 cm (maximum of 163.8 cm in 1975), with mean annual snow depth of 2.5 cm (as measured at Savage, Montana, 1905-2009) (WRCC 2009). Upland terrain consists of rolling hills that have been dissected into badlands by the stream system. Downstream of State Route 16, Thirteen Mile Creek cuts through alluvial terraces and emerges onto the floodplain of the Yellowstone River. Land use on the watershed is approximately 50% grass rangeland, 40% crop/pasture, 9% mixed rangeland, and 1% other (primarily mine/quarry and other agriculture) (NRIS 2009).

Non-crop upland vegetation of the Thirteen Mile Creek watershed consists predominantly of grasses (Poaceae) and sagebrush (*Artemesia* spp.) with scattered stands of ponderosa pine (*Pinus ponderosa*) and Rocky Mountain juniper (*Juniperus scopulorum*). Valley floors also support scattered stands of eastern cottonwood (*Populus deltoides*), green ash (*Fraxinus pennsylvanica*), Russian olive (*Eleagnus angustifolia*), and buffaloberry (*Shepherdia canadensis*). Riparian and littoral vegetation is dominated by sedges (*Carex* spp.), rushes (*Juncus* spp.), cattails (*Typha* spp.), snowberry (*Symphoricarpos albus*), milkweed (*Asclepias* spp.), and willow (*Salix* spp.).

The Thirteen Mile Creek watershed is underlain by Cretaceous and Paleocene sedimentary rocks consisting primarily of highly erodible sandstones and shales. Exposed rocks in the watershed consist of non-marine sediments of the Fort Union Formation, which contains economically extractable deposits of coal and discharges significant amount of groundwater (Alt and Hynd 1986, Morris et al. 1981). Stream discharge has not been systematically recorded in Thirteen Mile Creek (USGS 2009).

## **MATERIALS AND METHODS**

Fish populations and habitats were evaluated at 21 sites in Thirteen Mile Creek from 23 July to 10 August 2008 (Figure 1, Table 1). A site was defined as a contiguous unit of riffle and pool habitat as defined by Armantrout (1998). Only 40-m lengths of very long riffles and pools were included in fish and habitat sampling. Five sites lacked riffle development: site 1, a low gradient site at the Yellowstone River confluence (Figure 2c); site 2, a beaver impoundment constructed since the 2007 sampling period and inundating the previously existing riffle; site 15, an isolated pool at the County Road 542 crossing; and sites 20 and 21, long upland channels filled with terrestrial vegetation and isolated shallow pools near the headwaters of Thirteen Mile Creek (Figure 2d). All other sites sampled consisted of riffle-pool units.

On arrival at each site a metric Hip Chain distance measurer (No. 06220, Legend, Inc., Reno, NV) was used to measure total site length and the lengths of component riffles and pools. Orange DayGlo ribbon (DayGlo Color Corp., Cleveland, OH) was tied to conspicuous littoral vegetation every five meters to serve as a baseline for installing block seines, determining flow velocity, and spacing habitat sampling transects. A water sample for total suspended solids (TSS) was then collected, dissolved oxygen, conductivity, and pH were measured, and water and air temperatures were recorded as described below. Block seines were then installed for fish sampling. After fish sampling was completed, habitat variables were assessed as described below.

**Fish.** We used a simple DeLury (1947) type capture-removal approach to estimate fish community densities in adjacent riffle and pool habitats. Block seine installation and fish sampling were done before habitat evaluation in order to minimize movement of fish out of the disturbed site. Two 7.5-m x 1.2-m x 6.3-mm mesh straight seines, one 8.2-m x 1.5-m x 6.3-mm

mesh bag seine, and one 12.1-m x 1.8-m x 6.3-m straight seine were used as block seines. Block seines were installed at the upstream and downstream ends of each site and between adjacent riffles and pools (i.e., three block seines per site).

In pools we used a short, straight minnow seine (3.6 -m x 1.5 -m x 6.3 -mm mesh) to make successive removal passes through each isolated section. In wider pools a bag seine (8.2 -m x 1.5 -m x 6.3 -mm mesh) was used to make removal passes. In all pools two operators towed the working seine from the downstream end to the upstream end of each isolated section. The lead line was kept on the stream bottom and the seine landed at a convenient upstream location.

In riffles towing the seine was ineffective due to extremely shallow depths and the tendency of riffle species to hide under cobbles. Therefore, one operator held the working seine stationary at the downstream end of the isolated section, across the whole wetted channel and with the lead line on the stream bottom, while the second operator "kicked down" the riffle from upstream to downstream, agitating the substrate with his feet. The seine was then lifted quickly by both operators in midstream. We were unable to make fish population estimates on several riffles due to extremely low flow; riffles were either dry or reduced to shallow trickles that we were unable to seine. If small fish were present, they were sampled qualitatively with a dip net.

The number of removal passes (by towing or kicking) ranged from three to five and depended on how quickly we achieved a noticeable reduction in catch. After each removal pass fish were removed from the seine, held in a bucket of aerated water, identified using Holton and Johnson (2003) and Gould (1998), counted, and returned to the stream downstream of the lowest block seine. To estimate the density of fish in isolated riffles and pools, we regressed catch per pass on sum of catches to yield a gross population estimate. This estimate was then divided by the measured surface area of the habitat unit (length x mean width) to yield number of individuals per 100 m<sup>2</sup>.

**Habitat Variables.** The following habitat variables were evaluated at each site after fish sampling was completed unless otherwise noted.

#### Location:

**O** latitude and longitude

• landmark photography and notation on USGS 7.5-min quadrangle map

### Water Quality:

- **o** water temperature and air temperature (initial and final)
- o dissolved oxygen
- o pH
- **o** conductivity
- **o** total suspended solids

#### **Stream Flow:**

- **o** flow velocity
- **o** discharge

## **Channel Morphology:**

- **o** total site length and lengths of component riffle, run, and pool habitats
- **o** stream width (wetted width and channel width)

### o stream depth

#### **Streambed Composition:**

- **o** relative substrate composition
- **o** sediment depth
- o embeddedness

## **Riparian Conditions:**

- **o** adjacent land use (left and right banks)
- **o** buffer width (left and right banks)
- **o** bank erosion (left and right banks)
- **O** bank height and angle at water's edge (left and right banks)
- **O** entrenchment bank height and angle (left and right banks)
- **o** channel canopy shading

## **Cover:**

- **O** vegetative height at water's edge (left and right banks)
- **o** vegetative overhang (left and right banks)
- **O** bank undercut (left and right banks)
- **o** woody and other debris
- **o** emergent and submerged aquatic vegetation

Habitat variables were measured or visually estimated using a transect approach based on Simonson et al. (1994). Using a Hip Chain distance measurer as described above, transects were spaced at five-meter intervals perpendicular to the direction of stream flow, beginning one meter above the lower end of each site and ending at the last five-meter interval below the upper end of each site. For riparian variables transects were extended 10 meters inland from water's edge. Transects were numbered and worked from downstream to upstream, with the left bank and the right bank designated facing downstream. Variables were measured or visually estimated along a 0.3-m wide band centered on the transect line. A 20-m length of 0.25-in (31.8-mm) white nylon line and two metal stakes were used to mark transects while they were worked.

**Location.** Each site was marked on a standard USGS 7.5-min quadrangle map, and landmarks were noted. Latitude and longitude of each site were determined with a GPS Pioneer Satellite Navigator (Magellan Systems Corporation, San Dimas, CA); site coordinates were verified by direct measurement on 7.5-min quadrangle maps as described by NASA (2009). For reference and future identification, four color photographs were taken at each site: (1) upper end facing upstream; (2) upper end facing downstream; (3) lower end facing upstream; (4) lower end facing downstream.



Figure 1. Map of the Thirteen Mile Creek system, Dawson County, Montana, showing 21 sites sampled in 2008; inset shows headwaters sites 20 and 21 between Bloomfield and Red Top at smaller scale (based on U.S. Geological Survey, 7.5 Minute Topographic Series, Intake, Kohlberg Ranch, Red Top, and Bloomfield Quadrangles, provided by TerraServer 2009).

Site No.	County	Quadrangle <sup>a</sup>	Township	Range	Section	Latitude	Longitude	Stream Km	Elevation (m)	Date Sampled
Y <sup>b</sup>	Dawson	Intake	18N	56E	35	47°16'36"	104°32'57"	0.0	603.5	8/04/2008
1	Dawson	Intake	18N	56E	35	47°16'36"	104°32'57"	0.0	603.5	8/04/2008
2	Dawson	Intake	18N	56E	34/35	47°06'43"	104°33'23"	1.1	609.6	8/04/2008
3	Dawson	Intake	18N	56E	34	47°16'32"	104°33'15"	3.5	611.4	8/05/2008
4	Dawson	Intake	18N	56E	34	47°16'28"	104°33'26"	4.9	612.7	8/05/2008
5	Dawson	Intake	18N	56E	34	47°16'41"	104°33'28"	6.0	613.2	8/04/2008
6	Dawson	Intake	18N	56E	34	47°16'52"	104°33'28"	6.2	615.7	8/01/2008
7	Dawson	Intake	18N	56E	27	47°17'30"	104°34'01"	6.8	621.8	8/01/2008
8	Dawson	Intake	18N	56E	21	47°18'02"	104°35'25"	9.6	637.0	7/29/2008
9	Dawson	Intake	18N	56E	21	47°18'12"	104°35'33"	11.6	640.1	7/29/2008
10	Dawson	Intake	18N	56E	21	47°18'16"	104°35'41"	11.8	643.7	7/29/2008
11	Dawson	Intake	18N	56E	17	47°19'09"	104°36'59"	13.7	653.6	7/29/2008
12	Dawson	Intake	18N	56E	17	47°19'31"	104°36'14"	14.7	665.5	7/28/2008
13	Dawson	Intake	18N	56E	18	47°16'29"	104°37'15"	15.8	667.5	7/28/2008
14	Dawson	Intake	18N	56E	7	47°19'36"	104°37'22"	16.2	668.1	7/28/2008
15	Dawson	Intake	18N	56E	7	47°19'43"	104°37'21"	16.8	670.0	7/25/2008
16	Dawson	Intake	18N	56E	7	47°19'55"	104°37'22"	17.9	670.6	7/25/2008
17	Dawson	Kolberg Ranch	18N	56E	7	47°20'10"	104°37'46"	18.7	676.0	7/25/2008
18	Dawson	Kolberg Ranch	18N	56E	7	47°20'22"	104°38'18"	19.3	676.6	7/23/2008
19 <sup>c</sup>	Dawson	Kolberg Ranch	18N	56E	3	47°20'59"	104°41'19"	25.1	725.4	8/09/2008
20	Dawson	Red Top	18N	55E	15	47°25'26"	104°48'42"	48.6	759.9	8/10/2008 <sup>d</sup>
21	Dawson	Bloomfield	19N	54E	13	47°24'23"	104°53'47"	64.1	789.7	8/10/2008 <sup>d</sup>

 Table 1.
 Locations of 21 sampling sites in the Thirteen Mile Creek system, Dawson County, Montana, 2008.

<sup>a</sup>U.S. Geological Survey, 7.5 Minute Series (Topographic); <sup>b</sup>Yellowstone River; <sup>c</sup>South Fork (Lower) Thirteen Mile Creek.



Figure 2. Thirteen Mile Creek system, Dawson County, Montana, July-August 2008: (a) representative section of Thirteen Mile Creek below Dawson County Road 542 (view northwest toward sites 12, 13, 14, and 15); (b) South Fork (Lower) Thirteen Mile Creek (site 19) showing recently installed culvert replacing original wooden bridge; (c) low gradient reach near mouth on Yellowstone River side channel (site 1); (d) ephemeral channel near headwaters (Site 20); (e) northern redbelly dace (*Phoxinus eos*) habitat (site 16) showing dense growth of filamentous algae present during sampling in 2008; (f) typical downstream site on Yellowstone River floodplain terrace (Site 4).

Water Quality. Water and air temperatures were measured with a pocket field thermometer at the lower end of the pool at each site at the beginning and at the end of sampling. Dissolved oxygen concentration, water temperature, and conductivity were determined with a YSI Model 85 oxygen, conductivity, salinity, and temperature meter (Yellow Springs Instruments, Inc., Yellow Springs, OH) and pH with a pHep 3 pH meter (Hanna Instruments, Woonsocket, RI) at the lower end of the pool of each site before fish sampling. Whole water samples were collected in one-pint (0.48-liter) glass jars before fish sampling at the upper ends of sites 2, 6, 10, 11, 12, 15, 16, and 20. These samples were analyzed for total suspended solids (TSS) by Amatec Services, Inc. (Billings, MT) using SM 2540D (Clesceri et al. 1999).

**Stream Flow.** Flow velocity was measured at each site by timing three times the transit of a plastic fishing float over a measured distance of riffle (usually 10 m). An ExTech Digital Stopwatch/Clock (ExTech Instruments Corporation, Waltham, MA) was used for timing, and the results of three runs were averaged to yield mean flow velocity. Discharge was calculated by multiplying the cross-sectional area of the water column (using depth and width measurements as described below) by mean flow velocity (McMahon et al. 1996). Flow velocity in pools was usually too slow to measure using the buoyant object method, especially with wind interference, and low flow in riffles at many sites made estimating flow velocity and discharge impractical.

**Channel Morphology.** Total length of each site and lengths of component riffle and pool habitats were measured using a Hip Chain distance measurer as described above. Wetted width and channel width were measured to the nearest inch (2.54 cm) along transects at each site using a home-made grade stick fabricated from a 10-ft (3.05-m) length of <sup>3</sup>/<sub>4</sub>-in (1.9-cm) aluminum electrical conduit to which an equal length of steel measuring tape graduated in inches was attached with transparent duct tape. At very wide sites we used a 300-ft (91.4-m) fiberglass measuring tape graduated in inches (Model KL-300-18, Keson Industries, Inc., Naperville, IL). When different from wetted width, channel width was delimited by the presence of matted or silt-covered vegetation that had obviously been recently submerged or by the presence of white evaporite deposits. Water depths were measured by grade stick at five equidistant points along transects at each site, beginning at channel center and proceeding shoreward to within 10 cm of the left and right banks. This created six equal-width cells for calculation of mean depth and channel cross-sectional area.

**Streambed Composition.** Substrate material definitions of Simonson et al. (1994) were used. Substrate composition and embeddedness were estimated visually to the nearest five percent along transects at each site. Sediment depth was defined as the depth of yielding material (silt or muck) overlaying a non-yielding substrate (bedrock, boulders, rubble, cobbles, gravel, or sand). In the field, stream bottom material which could be penetrated with the grade stick was considered sediment, and its depth was measured by grade stick along transects at each site at the same points used for measuring water depth. Embeddedness was defined as percent of rubble, cobble, or gravel particles covered by sediment.

**Riparian Conditions.** The land use classification and riparian definitions of Simonson et al. (1994) were used. Riparian land use and canopy shading were estimated visually to the nearest five percent along transects at each site. Buffer width, bank erosion, bank slumping, bank height at water's edge, and entrenchment bank height were measured by grade stick, measuring tape, or 25-ft (7.62-m) extension pole (Model 90180, Crane Enterprises, Inc., Mound City, IL) to the nearest foot (0.30 m) at each site, except for bank height at water's edge, which was measured to the nearest inch (2.54 cm). Bank angle (with respect to horizontal) of both water's edge and

entrenchment banks was measured with an industrial clinometer (Model No. 78555, Shinwa Rules Co., Ltd., Tsubame City, Niigata Pref., Japan 959-1276, <u>http://www.shinwasokutei.co.jp/index.html</u>). Buffer width was defined as the width of undisturbed riparian vegetation extending from water's edge to a point 10 m inland. Bank erosion was defined as the width of bare riparian soil from water's edge to a point 10 m inland, and slumping was defined as the width of detached whole riparian soil masses, vegetated or not, from water's edge to a point 10 m inland. An entrenchment bank was defined as a bank higher than the water's edge bank and separated from water's edge by a distance of 10-30 m. Entrenchment banks obviously contained stream flow during peak flow periods and often coincided with water's edge banks on outside meander bends.

**Cover.** Cover was defined as any instream material, riparian material, or streambank configuration that could provide protection for most fishes of the size range (< 15 cm) found in Upper Sevenmile Creek. The amount of wetted channel bottom covered by woody debris, other debris, emergent vegetation (aquatic or flooded riparian), or submerged vegetation (aquatic or submerged riparian) was visually estimated to the nearest five percent along transects at each site. In order to qualify as cover, debris or vegetation had to occur in water at least 15 cm deep; Simonson et al. (1994) used a one-foot (30.5-cm) water depth criterion, but their approach emphasized cover requirements of larger game fishes. Riparian vegetative overhang, vegetative height at water's edge, and bank undercut were measured by grade stick to the nearest inch (2.54 cm) along transects at each site. In order to qualify as cover, vegetative overhang, vegetative height, and bank undercut had to be at least 15 cm.

**Other.** General observations on flora, fauna, hydrology, geology, and water quality and land use were noted at each site.

## **RESULTS AND DISCUSSION**

**Developments.** Five sampling sites were added to the 16 sites sampled in 2007 and the site numbering sequence modified as follows (Figure 1, Table 1):

Year	Sites																					
2007	1 <sup>a</sup>	2	3	4	-	5	6	7	8	9	10	11 <sup>c</sup>	12 <sup>c</sup>	13	14	15	16	17	18	19 <sup>d</sup>	20 <sup>c</sup>	_
2008	Y <sup>a</sup>	1	2	3	<b>4</b> <sup>b</sup>	5	6	7	8	9	10	11 <sup>c</sup>	12 <sup>c</sup>	13	14	15	16	17	18	19 <sup>d</sup>	20 <sup>c</sup>	21 <sup>b</sup>

<sup>a</sup>Yellowstone River; <sup>b</sup>new site added to sequence; <sup>c</sup>site included in sequence in 2007 but not precisely located or sampled until 2008; <sup>d</sup>South Fork (Lower) Thirteen Mile Creek.

Major habitat changes had occurred at three sites since the 2007 sampling period. A beaver impoundment had been constructed at site 2, inundating the riffle at the site, and the beaver impoundment present at site 5 (old Montana Route 16 crossing) had been removed by the landowner to prevent flooding of adjacent corrals. The old timber bridge over South Fork (Lower) Thirteen Mile Creek (site 19) had replaced by a culvert, and most of the coldwater pool present in 2007 had been filled in (Figure 2b).

					Variab	le <sup>b</sup>					
Site <sup>a</sup>	Т	DO	С	TSS	pH	0	L	W	D		
	(°C)	(mg/l)	(µS)	(mg/l)	•	$(m^{3}/sec)$	( <b>m</b> )	(m)	(cm)		
Y <sup>c</sup>	14.0	4.1	1300	524.7	9.2		>40	13.4/110.3	16.2		
1 P	14.5	3.9	1350	305.6	9.3	—	>40	0.3/9.5	10.3		
2 P <sup>d</sup>	15.0	4.9	1815	365.1	9.1	_	18.5	3.5/3.7	89.3		
3 R						< 0.1	5.0	1.6/2.7	8.7		
3 P	12.0	4.9	1720	478.3	9.2		2.5	2.4/2.8	65.0		
4 R						< 0.1	2.6	2.4/2.8	12.0		
4 P	13.5	5.1	1630	374.6	9.3		12.5	1.7/2.4	16.6		
5 P <sup>e</sup>	18.0	3.8	1745	263.7	9.0		37.8	1.1/3.8	2.2		
6 R						< 0.1	60.0	3.0/4.8	3.7		
6 P	18.5	4.2	1360	265.1	9.1		>40	4.0/4.5	24.1		
7 P	12.0	5.4	1190	255.4	9.2		63.0	2.6/4.0	12.3		
7 R						0	12.0	0/4.6	0		
8 P	17.0	5.2	1310	264.3	9.2		14.0	3.8/12.0	26.0		
8 R						< 0.1	0.5	1.6/6.6	3.3		
9 P <sup>r</sup>							14.0	0/12.0	0		
<b>9 R</b> <sup>1</sup>						0	0.5	0/6.6	0		
10 P	19.0	5.1	1650	345.7	9.3		30.1	2.5/3.8	13.6		
10 R						0	0.5	0.3/2.5	1.0		
11 R						0	16.5	0/3.9	0		
11 P	19.0	5.2	1815	215.4	9.1		38.5	3.6/4.1	17.4		
12 R			10.10			0	15.0	0/4.4	0		
12 P	14.0	4.1	1940	394.3	9.0		38.0	3.1/5.2	19.5		
13 P	19.0	4.2	1850	388.1	9.1		10.0	3.1/4.8	14.1		
13 R						0	5.0	0.7/4.8	3.7		
14 R	20.0	1.0	1 < 1 5	0.67.4	0.1	0	5.0	0/9.5	0		
14 P	28.0	4.0	1615	367.4	9.1	0	25.0	3.2/7.9	14.5		
15 P	17.0	4.5	1500	388.5	9.3	0	28.3	5.8/6.7	53.1		
16 K	10.0	1.6	1505	0764	0.1	<0.1	5.0	2.2/2.8	9.3		
10 P	18.0	4.6	1505	3/6.4	9.1	.0.1	28.3	3.8/4.4	48.0		
17 K	12.0	1.0	005	2015	0.0	<0.1	2.5 15.5	2.6/3.0	12.8		
1/ ľ 10 D	15.0	4.9	905	384.3	8.9	<0.1	15.5	2.4/3.0	<u> </u>		
10 K 10 D	16.0	56	705	215.2	Q /	<0.1	5.U 20.0	2.3/3.0 2.1/2.7	24.0 21_4		
10 P 10 D <sup>g</sup>	10.0	5.0	125 805	213.3 211 4	0.4	0	>40	3.1/3.1 1 $1/2.9$	21.4 7.2		
19 P° 20 D	12.0	5.7	720	211.4	0.7 Q 1	0	>40	1.4/3.0 1 1/4 6	1.5		
20 r 21 P	12.0	3.0 / 8	750	203.3	0.1	0	>40	1.1/4.0	0.3		
41 ľ	12.0	4.0	730	200.9	0.9	U	<i>&gt;</i> 40	0.3/3.8	9.2		

Table 2.Water quality, stream flow, and channel morphology variables for 21 sites in theThirteenMile Creek system, Dawson County, Montana, 23 July – 10 August 2008.

<sup>b</sup>T: mean water temperature; DO: dissolved oxygen; C: conductivity; TSS: total suspended solids; Q: mean discharge; L: habitat unit length; W: mean wetted width/channel width; D: mean depth. A blank indicates that a variable was not determined at the site; a dash (-) indicates that the value was positive but not measurable with the technique used. <sup>c</sup>Sample not stratified into riffles and pools; habitats included riffles, channel pools, and backwater pools.

<sup>d</sup>Riffle present in 2007 but inundated by beaver dam in 2008.

<sup>e</sup>Site consisted of a beaver pond in 2007; subsequently removed by landowner.

<sup>f</sup>Entire site (riffle and pool) dry.

<sup>g</sup>Timber bridge present in 2007 removed and replaced by culvert; most of original pool filled in.

					Variable <sup>b</sup>				
Site <sup>a</sup>	LU	BW	ER	SL	H-1	H-2	A-1	A-2	CS
		( <b>m</b> )	( <b>m</b> )	( <b>m</b> )	(cm)	( <b>m</b> )	( • )	( • )	(%)
Y <sup>c</sup>	MI/WO	>10/>10	—/3	0/0	0/500	_/_	-/90	_/_	0
1 P	MI/MI	>10/>10	—/3	0/0	_/_	1.8/1.8	5/5	20/22	0
2 P	MI/MI	>10/>10	0/0	0/0	130/35	—/95	-/90	-/90	0
3 R	MI/MI	>10/>10	0/0	0/0	200/-	_/_	92/2	_/_	0
3 P	MI/MI	>10/>10	0/0	0/0	90/40	_/_	92/60	_/_	0
4 R	MI/MI	>10/>10	0/0	0/0	50/20	_/_	45/42	_/_	0
4 P	MI/MI	>10/>10	0/0	0/0	80/-	_/_	92/—	_/_	3.3
5 P	MI/MI	>10/>10	0/0	0/0	38/69	_/_	45/75	_/_	12.2 <sup>d</sup>
6 R	PA/PA	0/0	0/-	0/0	33/56	_/_	1/2	_/_	0
6 P	PA/PA	0/>10	0/-	0/0	_/_	_/_	_/_	_/_	0
7 P	PA/MI	0/>10	-/0	0/0	49/134	_/_	30/62	_/_	0
7 R	PA/MI	0/0	—/0	0/0	18/175	_/_	—/81	_/_	0
8 P	EX/PA	0/0	-/0	0/0	-/110	_/_	-/85	_/_	0
8 R	EX/PA	0/0	—/0	0/0	—/30	_/_	-/80	_/_	0
9 P	PA/PA	0/0	0/0	0/0	6/22	_/_	-/22	_/_	0
9 R	PA/PA	0/0	0/0	0/0	35/65	_/_	-/30	_/_	0
10 P	WO/PA	>10/0	0/0	0/0	220/-	0.6/-	81/-	10/-	14.3 <sup>ª</sup>
10 R	WO/PA	>10/0	0/0	0/0	120/-	_/_	70/—	_/_	100 <sup>a</sup>
11 R	PA/PA	>10/0	0/0	0/0	40/48	_/_	_/_	_/_	0
11 P	PA/PA	>10/0	0/0	0/0	45/132	_/_	90/90	_/_	0
12 R	PA/PA	>10/0	0/0	0/0	40/47	_/_	_/_	_/_	0
12 P	PA/PA	>10/0	0/0	0/0	46/129	_/_	85/90	_/_	0
13 P	PA/PA	>10/>10	0/0	0/0	33/53	_/_	—/85	_/_	0
13 R	PA/PA	>10/>10	0/0	0/0	40/105	_/_	-/85	_/_	50
14 R	PA/PA	0/0	-/1.0	0/0	-/100	_/_	10/50	_/_	0
14 P	PA/PA	0/0	-/1.1	0/0	-/110	_/_	10/50	_/_	0
15 P	PA/PA	0/0	0/0	0/0	_/_	_/_	5/1	_/_	100°
16 K	PA/PA	0/0	0/0	0/0	-/40	_/_	-/85	_/_	0
16 P	PA/PA	0/0	0.1/0	0/0	63/45	_/_	44/89	_/_	30
17 K	PA/PA	0/0	—/— 1.(/	0/0	20/25	_/_	12/85	_/_	0
17 P	PA/PA	0/0	1.0/-	0/0	/0/46	_/_	65/38	_/_	0
18 K		0/0	0/0	0/0	-/40	_/_	-/90	_/_	0
10 ľ 10 ľ	PA/PA	0/0	0/0	0/0	20/22	_/_	90/34	_/_	0
19 P 20 D		>10/>10	0/0	0/0	30/22 25/26	_/_	20/20	_/_	0
20 P	PA/PA	>10/>10	0/0	0/0	35/30 01/21	_/_	20/30	_/_	0
21 P	PA/PA	>10/>10	0/0	0/0	21/31	_/_	30/25	_/_	U

Table 3. Mean values of riparian variables for 21 sites in the Thirteen Mile Creek system, Dawson County, Montana, 23 July – 10 August 2008.

<sup>b</sup>Mean values per habitat unit [LU: land use (EX: exposed unvegetated rock or gravel bars; ME: meadow; MI: mixed meadow; OP: open marsh; PA: pasture; SH: shrubs); BW: buffer width; ER: erosion; SL: slumping; H-1: bank height at water's edge; H-2: entrenchment bank height; A-1: bank angle (with respect to horizontal) at water's edge; A-2: entrenchment bank angle; CS: canopy shading]. A zero (0) indicates that the feature was absent. A dash (-) indicates that the value was positive but that it was negligible on the scale used; for entrenchment banks, a dash indicates that the bank was greater than 10 m from water's edge. A slash (/) separates left bank/right bank values.
<sup>c</sup>Sample not stratified into riffles and pools; habitats included riffles, channel pools, and backwater pools.

<sup>d</sup>Bridge shading at sites 5 and 15; cliff shading at site 10.

				Variable <sup>b</sup>			
Site <sup>a</sup>	VH	VO	BU	WD	OD	EV	SV
	( <b>cm</b> )	( <b>cm</b> )	(cm)	(%)	(%)	(%)	(%)
Y <sup>c</sup>	200/>1000	0/0	0/0	0	0	0	0
1 P	_/_	0/0	0/0	0	0	0	0
2 P	100/115	35/45	0/0	0	0	5.0	0
3 R	7/45	35/-	10.0/0	0	0	5.0	1.6
3 P	94/94	50/41	2.7/0	0	0	1.5	1.4
4 R	100/106	50/-	0/0	0	0	5.0	
4 P	100/93	_/_	0/0	0	0	5.0	
5 P	41/32	0/0	0/0	0	0	0	5.0
6 R	10/49	0/0	0/0	0	0	0.4	
6 P	73/34	0/0	0/0	0	0	11.8	
7 P	17/44	0/0	0/0	0	0	7.4	
7 R	—/66	0/0	0/0	0	0		
8 P	100/110	0/0	0/0	0	0	3.3	0
8 R	100/120	0/0	0/0	0	0	5.0	0
9 P	64/84	0/0	0/0	0	0	0	0
9 R	65/30	0/0	0/0	0	0	0	0
10 P	112/-	0/0	4.3/0	5.7	0	0	0
10 R	95/0	0/0	0/0	5	0	0	0
11 R	63/10	0/0	0/0	0	0	0	0
11 P	71/34	0/0	0/0	0	0	0	
12 R	73/8	0/0	0/0	0	0	0	0
12 P	131/42	0/0	0/0	0	0	0	—
13 P	53/20	0/0	0/0	0	0	0	46.7
13 R	60/150	0/0	0/0	0	0	0	0
14 R	—/2	0/0	0/0	0	0	0	0
14 P		0/0	0/0	0	0	0	0
15 P	0/0	0/0	0/0	0	0	0	0
16 R	10/20	0/0	0/0	45.0	0	0	5.0
16 P	15/58	0/0	5/23.3	0	0	1.7	58.3
17 R	30/60	0/0	0/0	0	0	2.5	47.5
17 P	28/10	0/0	0/0	0	0	0	50.0
18 R	0/20	0/0	0/60.0	0	0	2.5	25.0
18 P	50/22	0/0	0/0	0	0	3.8	70.2
19 P	_/_	11/33	0/0	0	0	— •	—
20 P	98/101	_/_	0/0	0	0	90 <sup>a</sup>	0
21 P	47/64	_/_	0/0	0	0	80 <sup>d</sup>	0

Table 4.Cover variables for 21 sites in the Thirteen Mile Creek system, Dawson County, Montana,23 July – 10 August 2008.

<sup>b</sup>Mean values per habitat unit (VH: vegetative height; VO: vegetative overhang; BU: bank undercut; WD: woody debris; OD: other debris; EV: emergent vegetation; SV: submerged vegetation. A slash (/) separates left bank/right bank values. A zero (0) indicates that the feature was absent. A dash (-) indicates that the value was positive but that it was negligible on the scale used.

<sup>c</sup>Sample not stratified into riffles and pools; habitats included riffles, channel pools, and backwater pools.

<sup>d</sup>Mixed terrestrial and emergent aquatic vegetation in ephemeral channel.

<sup>e</sup>Eastern cottonwoods (*Populus deltoides*) on right bank.

		5	Substra	ate Comp	osition <sup>b</sup>			Sedime	ntation <sup>b</sup>
Site <sup>a</sup>	BO	RC	GR	SA	SI	CL	MU	SD	EM
	(%)	(%)	(%)	(%)	(%)	(%)	(%)	(cm)	(%)
Y <sup>c</sup>		15.5	54.5		30.0			—	30
1 P					100			120.0	100
2 P		5.0	75.0		20.0			25.0	80
3 R		80	20					0	0
3 P					100			20.2	100
4 R	10.0	10.0	85.0		5.0			2.5	5
4 P			10.0		90.0			5.0	90
5 P			60.0		20.0	10.0		0	5
6 R		30.0	0.0		10.0			—	10
6 P					100.0			23.0	100
7 P		3.3	16.7		80.0			9.6	100
7 R		40.0	50.0		10.0			0	0
8 P	3.3	8.4	83.3		5.0			0	5
8 R		20.0	75.0		5.0			0	5
9 P		5.0	90.0		5.0			0	5
9 R		5.0	90.0		5.0			0	5
10 P		5.7	85.0		9.3			0	10
10 R		5.0	85.0		10.0			0	10
11 R		6.0	89.0		4.0			0	5
11 P		1.3	86.0		12.7			0	15
12 R		5.0	90.0		5.0			0	5
12 P		1.7	86.0		12.3			0	15
13 P		5.0	86.3		8.7			0	5
13 R		5.0	89.5		5.5			0	10
14 R		5.0	90.0		5.0			0	5
14 P		5.0	90.0		5.0			0	5
15 P		20.0	70.0		10.0			3.1	10
16 R		2.5	90.0		7.5			0	10
16 P		1.7	80.0		18.3			18.3	5
17 R		2.5	90.0		7.5			0	10
17 P		2.0	68.0		30.0			0	30
18 R		5.0	90.0		5.0			0	5
18 P		5.0	80.0		15.0		100 <sup>d</sup>		15
19 P					100 0 <sup>e</sup>		100°	5.5	100
20 P					100.0°			5.5	100
21 P					100.0			6.8	100

Table 5. Streambed composition variables for 21 sites in the Thirteen Mile Creek system, Dawson County, Montana, 23 July – 10 August 2008.

<sup>b</sup>BO: boulders; RC: rubble/cobble; GR: gravel; SA: sand; SI: silt; MU: muck; SD: sediment depth; EM: embeddedness. A blank indicates that a component was not visually apparent.

<sup>c</sup>Sample not stratified into riffles and pools; habitats included riffles, channel pools, and backwater pools.

<sup>d</sup>Black organic muck and loose detritus underlain by gravel.

<sup>e</sup>Mostly dried silt in dewatered channels and shallow silt mixed with gravel in isolated pools.

Stream Flow. Upstream of site 18 the Thirteen Mile Creek system consisted of long sections of dry channel separating a few isolated pools at bridge crossings (Figure 2b). This reach of the system exhibited no active flow during the sampling period, although it is probably responsive to storm events and surface runoff because the associated drainage area is relatively The dry channel sections were occupied by terrestrial vegetation, indicating that duration of large. flow in these channels is brief. From site 18 downstream to site 15 at the Dawson County Road 542 crossing active flow was evident, but it was barely measurable using the buoyant object method, usually less than 0.1 m<sup>3</sup>/s (Figure 2e, Table 2). Downstream of site 15 to the mouth at the Yellowstone River the stream lacked hydrologic continuity. Where zero discharge was recorded, riffles were dry; if riffles were wet it was assumed that flow existed but that it was less than our lower limit of precision using the buoyant object method ( $<0.1 \text{ m}^3/\text{s}$ ). Pools at all sites downstream of site 15, except the pool at site 9, retained some water. Active flow between sites 18 and 15 was probably a result of seepage from several wells located adjacent to the channel and used for livestock water supply.

**Water Quality.** Water quality was relatively uniform longitudinally during the sampling period. Water temperature averaged 15.8°C (range 11.0°C to 28.0°C), cooler than in 2007, and generally responded daily to air temperature, which averaged 21.0°C (range 11.0°C to 36.0°C) (Table 2).

Dissolved oxygen (DO) averaged 4.7 mg/l (range from 3.9 mg/l to 5.7 mg/l) and was marginal for support aerobic aquatic life (Table 2). Low DO concentrations may have been a result of high air temperatures, shallow average pool depth, and lack of hydrologic continuity. Average pH was 9.0 with little variation longitudinally (Table 2).

Concentrations of total suspended solids (TSS) varied little longitudinally, averaging 323.6 mg/l (range 205.3 mg/l to 524.7 mg/l, the latter in the Yellowstone River). TSS concentrations probably reflected local disturbances such as bank erosion, beaver activity, or cattle use, or rather than downstream accumulation of TSS (Table 2). Thirteen Mile Creek probably receives a relatively high natural inorganic sediment load due to weathering of the sparsely vegetated badlands and the unconsolidated glacial outwash that comprise a large part of its watershed.

Conductivity was relatively high, averaging 1390  $\mu$ S (range 725  $\mu$ S to 1940  $\mu$ S) (Table 2). High conductivity reflects high concentrations of total dissolved solids (TDS), which are derived from runoff inputs of a sparsely vegetated watershed in a highly evaporative climate.

**Channel Morphology.** Downstream of site 18, Thirteen Mile Creek exhibited well defined riffle-pool development, although riffles were often dry under prevailing low flow conditions. Pools ranged from 2.5 m to greater than 40 m in length, averaging 2.5 m in wetted width and 24.0 cm in depth. Riffles ranged from 0.5 m to 16.5 m in length, averaging 1.2 m in wetted width and 5.6 cm in depth. Lengths and wetted widths of pools and riffles were generally less than in 2007, reflected lower flow. Because of low flow conditions, channel width was usually greater than wetted width (Table 2).

**Riparian Conditions.** Adjacent land use at most sites on Thirteen Mile Creek was pasture and mixed meadow. Where entrenchment banks were relatively high and steep, undisturbed

meadow, mixed meadow, and shrubs were usually evident, probably because cattle avoided these habitats. Riparian erosion and slumping were minimal and were largely restricted to high angle faces of entrenchment banks. At many sites we evaluated buffer width (width of undisturbed riparian vegetation) as 0 because grazing had modified natural riparian vegetation cover. However, in most locations this indicated a reduction in vegetation height rather than the presence of bare soil or erosion. Banks at water's edge were low, usually less than 0.5 m, and approximately vertical, while most of the channel meandered across a broad floodplain with entrenchment banks, if present, farther than 10 m from water's edge. Eastern cottonwoods were frequently adjacent to the channel, but they were never close enough to provide canopy shading during daylight hours. Significant "canopy" shading was provided most of the day at sites 5 and 15 by bridge cover and before noon by cliff shadow at site 10 (Table 3).

**Cover.** Vegetative height at water's edge averaged 50.0 cm, (range 0.0 cm to 131.0 cm). Vegetative overhang was absent or negligible at many sites; where present, it averaged 37.5 cm (Table 4). Instream cover, including bank undercut, woody debris, other debris, emergent vegetation, and submerged vegetation, was generally scarce. Most pools had some submerged aquatic vegetation, mostly *Ceratophyllum* sp. and *Chara* sp., but it was difficult to visually assess its extent due to high turbidity. Significant amounts of submerged vegetation were associated with deeper pools and riffles at sites 13, 16, 17, and 18 (Table 4). Attached filamentous algae covered the streambed material in most riffles and shallow poolsand the surfaces of pools at sites 16, 17, and 18 (Figure 2e). The most important cover in Thirteen Mile Creek may consist of "virtual" cover provided by tall riparian vegetation.

**Streambed Composition.** The predominant substrate materials in Thirteen Mile Creek were rubble/cobble, gravel, and silt; silt embeddedness was significant at deeper pool sites (Table 5). Riffles generally exhibited higher proportions of rubble and cobble than did pools. Boulders were evident only at sites 4 and 8, where the creek had apparently cut through more resistant geological formations, leaving boulders or slabs of limestone in the channel and riparian areas. Most pools were not deeply sedimented except for several lower gradient pools near the mouth of Thirteen Mile Creek and on the Yellowstone River floodplain terrace (Figures 2c and 2f, Table 5).

**Fish Communities.** Since 1975, 29 species of fishes have been collected in Thirteen Mile Creek and the immediately adjacent side channel of the Yellowstone River, to which we added two new records: sturgeon chub (*Macrhybopsis gelida*) and western silvery/plains minnow (*Hybognathus argyritus/placitus*), both from the Yellowstone River adjacent to the mouth of Thirteen Mile Creek (Tables 6 and 7).

In 2007 we failed to collect 6 species previously reported in Thirteen Mile Creek, including goldeye (*Hiodon alosoides*), mountain sucker (*Catostomus platyrhynchus*), bigmouth buffalo (*Ictiobus cyprinellus*), northern pike (*Esox lucius*), pumpkinseed (*Lepomis gibbosus*), and sauger (*Sander canadense*) (Table 6). Goldeye, mountain sucker, bigmouth buffalo, northern pike, and sauger are riverine species associated with the Yellowstone River and its larger tributaries; they may appear in small numbers in the lower reaches of Thirteen Mile Creek, especially during their spring spawning seasons.

Species <sup>a</sup>	Pre-1981 <sup>b</sup>	<b>MFISH</b> <sup>c</sup>	2007 <sup>d</sup>	2008 <sup>d</sup>	Total
goldeve (Hiodon alosoides)	Р	Р			
common carp ( <i>Cyprinus carpio</i> )	Р	Р		143	143
brassy minnow ( <i>Hybognathus hankinsoni</i> )	Р	Р		5	5
western silvery/plains minnow (Hybognathus					
argyritus/placitus)				24	24
emerald shiner (Notropis atherinoides)	Р			2	2
sand shiner (Notropis stramineus)	Р	Р	33	16	49
northern redbelly dace (Phoxinus eos)	Р	Р	52	9	61
fathead minnow (Pimephales promelas)	Р	Р	19	56	75
flathead chub (Platygobio gracilis)	Р	Р	68	16	84
sturgeon chub (Macrhybopsis gelida)				1	1
lake chub (Couesius plumbeus)	Р	Р		7	7
longnose dace (Rhinichthys cataractae)	Р	Р	212	316	528
creek chub (Semotilus atromaculatus)	Р	Р	2026	1004	3030
river carpsucker (Carpiodes carpio)	Р	Р	3	3	6
longnose sucker (Catostomus catostomus)	Р	Р	15	3	18
mountain sucker (Catostomus platyrhynchus)	Р	Р			
white sucker (Catostomus commersoni)	Р	Р	248	95	343
smallmouth buffalo (Ictiobus bubalus)	Р	Р	1		1
bigmouth buffalo (Ictiobus cyprinellus)	Р	Р			
shorthead redhorse (Moxostoma macrolepidotum)	Р	Р	4		4
black bullhead (Ameiurus melas)	Р	Р	1		1
channel catfish (Ictalurus punctatus)	Р	Р		2	2
stonecat (Noturus flavus)	Р	Р		3	3
northern pike (Esox lucius)	Р	Р			
plains killifish (Fundulus zebrinus)	Р	Р	184	130	314
brook stickleback (Culaea inconstans)	Р	Р	848	162	1010
green sunfish (Lepomis cyanellus)	Р	Р	3	4	7
pumpkinseed (Lepomis gibbosus)	Р	Р			
sauger (Sander canadense)	P	Р			
TOTAL FISH			3717	2001	5718
TOTAL SPECIES	27	26	15	20	29

Table 6.	Species and numbers of	fishes collected in	the Thirteen Mi	ile Creek system,	Dawson County,
Montana.					

<sup>a</sup>Common and scientific names according to Robins et al. (1991).

<sup>b</sup>Elser et al (1980), Morris et al. (1981); check (**P**) means present but numerical data not available.

 $^{\circ}$ MFIS (2008); check (**P**) means present but numerical data not available.

<sup>d</sup>Present study (2007, 2008).

		Number/Site																						
Common Name	Scientific Name	Y <sup>a</sup>	1	2	3	4	5	6	7	8	9 <sup>b</sup>	10	11	12	13	14	15	16	17	18	19 <sup>c</sup>	20	21	Σ
common carp	Cyprinus carpio	21		6	14	102																		143
western silvery/plains	Hybognathus																							
minnow	argyritus/placitus	24																						24
brassy minnow	Hybognathus hankinsoni	5																						5
emerald shiner	Notropis atherinoides	2																						2
sand shiner	Notropis stramineus	16																						16
northern redbelly dace	Phoxinus eos													1			1	3	4		1			9
fathead minnow	Pimephales promelas	31	2	5	1	1	3			7			2					3						56
flathead chub	Platygobio gracilis	16																						16
sturgeon chub	Macrhybopsis gelida	1																						1
longnose dace	Rhinichthys cataractae	1	5	4			1	3		4		15	12	35	55	113	17		37	14				316
lake chub	Couesius plumbeus									2				4					1					7
creek chub	Semotilus atromaculatus	9		5	6	14	1	28	132	79		84	71	134	69	82	117	39	43	91				1004
longnose sucker	Catostomus catostomus									3														3
white sucker	Catostomus commersoni	1						1		36		3	5	5	14	12	1	3	11	3				95
smallmouth buffalo	Ictiobus bubalus																							0
shorthead redhorse	Moxostoma macrolepidotum																							0
river carpsucker	Carpiodes carpio	1		1	1																			3
black bullhead	Ameiurus melas																							0
channel catfish	Ictalurus punctatus	2																						2
stonecat	Noturus flavus	3																						3
plains killifish	Fundulus zebrinus				1	4			125															130
brook stickleback	Culaea inconstans			9	15	13				20		20	15	2	16	1		6	7	20	15	2	1	162
green sunfish	Lepomis cyanellus									4														4
ΤΟΤΑL (Σ)		133	7	30	38	134	5	32	257	155	0	122	105	181	154	208	136	54	103	128	16	2	1	2001

Table 2. Species and numbers of fishes collected in the Thirteen Mile Creek system, Dawson County, Montana, 23 July – 10 August 2008.

<sup>a</sup>Yellowstone River side channel adjacent to mouth of Thirteen Mile Creek.

<sup>b</sup>No water.

<sup>c</sup>South Fork Thirteen Mile Creek.

Deeper vegetated pools provide ample habitat for brook stickleback and northern redbelly dace in Thirteen Mile Creek, especially in the more perennial section between sites 18 and 15, where they may coexist with lake chubs (*Couesius plumbeus*) (Figure 2e). In 2007 we collected 714 brook sticklebacks and no other species in the isolated coldwater pool at site 19 on South Fork (Lower) Thirteen Mile Creek. The installation of a culvert replacing the old timber bridge over South Fork (Lower) Thirteen Mile Creek resulted in the loss of most of this pool, and we collected only 15 brook sticklebacks at site 19 in 2008. The abundance of brook sticklebacks may be limited by nocturnal anoxia when pools become isolated and lack throughflow during the summer.

Records of pumpkinseed in many lower Yellowstone River tributaries may represent escapes from populations stocked in ponds and reservoirs in the watershed. However, we think that many of these records represent misidentification of green sunfish (*Lepomis cyanellus*), which bear orange and red spots superficially resembling pumpkinseeds during their early life history.

The extent of riffle habitat in Thirteen Mile Creek during the sampling period was minimal, and we collected only a few longnose dace (*Rhinichthys cataractae*), creek chubs (*Semotilus atromaculatus*), and plains killifish (*Fundulus zebrinus*) in those riffles that exhibited flow during the 2008 sampling period.. Flathead chub (*Platygobio gracilis*) and sturgeon chub were also encountered in riffles at site Y in the Yellowstone River side channel (Table 7). The dewatering of riffles during the summer is probably a significant factor limiting abundance and diversity of riffle fauna, including longnose dace and stonecat (*Noturus flavus*), in Thirteen Mile Creek.

The most abundant and widely distributed pool species were creek chub, longnose dace, white sucker (*Catostomus commersoni*), and plains killifish (*Fundulus zebrinus*) (Table 7). Longnose dace probably retreat to pools when riffles are dewatered in middle to late summer.

The two headwaters sites (20 and 21) between Red Top and Bloomfield not sampled in 2007 (landowners never home) were sampled in 2008. The sites consisted of long, low gradient, dry channels occupied by terrestrial vegetation and a few widely interspersed, isolated pools containing emergent aquatic vegetation and a few brook sticklebacks. The flow regime of this section of Thirteen Mile Creek is apparently ephemeral and only in direct response to local precipitation

We were unable to effectively estimate densities of fish communities in riffles and pools due to low flows, poor capture rates, or failure to achieve significant catch reduction. The greatest numbers of fish were collected at sites 4, 7, 8, 10, 11, 12, 13, 14, 15, 17, and 18, consisting mostly of creek chubs, longnose dace, young-of-year or 1+ white suckers, and plains killifish. There was no clear relationship between numbers of fish captured at each site and the habitat variables we measured, as was observed in Burns Creek, O'Fallon Creek, and Upper Sevenmile Creek (Barnes 1999, Barnes 2004, Barnes and Siegle 2003, Barnes and Silbernagel 2001, Barnes and Westlind 2000, Barnes et al. 2002, Barnes et al. 2005, Barnes et al. 2006, Barnes et al. 2007).

In Burns Creek, contingent phenomena such as storm flow flushing and distribution of beaver dams in time and space were thought to influence fish community density and diversity as much as relatively "fixed" habitat factors (Barnes and Silbernagel 2001). However, Burns Creek exhibited perennial flow, so fish communities could at least partially stabilize.

In O'Fallon Creek, a similar contingent phenomenon may be the timing and extent of riffle

dewatering in the summer. Fish moving upstream or downstream may be trapped, often at very high densities, in shrinking pools with low quality habitat as riffles dry up and interrupt the hydrologic continuity of the stream. Similarly, drying riffles may exclude fish from preferred habitats. The overall "flashiness" of the stream system probably prevents fish communities from stabilizing and efficiently partitioning available habitats (Barnes 2004, Barnes et al. 2002, Barnes and Siegle 2003). Similar factors probably operate in Thirteen Mile Creek. Sites 2, 3, 4, and 5 featured significant riparian vegetation height/overhang, relatively low surface/depth ratios, and morning/evening canopy or bank shading, yet the densities of fish encountered were lower than would be expected in such relatively "good" habitat. The proliferation and random placement of beaver dams in time and space in this lower reach of Thirteen Mile Creek may significantly limit movement of fish within the system and between the system and the Yellowstone River, as was hypothesized for Burns Creek (Barnes and Silbernagel 2001).

Thirteen Mile Creek is a highly variable system in which extreme high and low flow, hydrologic discontinuity, and spatiotemporal distribution of beaver dams present survival challenges to the resident fish communities. These factors may override other macrohabitat variables, such as substrate composition and instream cover, in controlling density and distribution of these communities.

An understanding of the relationships among fish communities, habitat variables, and contingent phenomena in small intermittent prairie streams such as Thirteen Mile Creek would require several years of study under a full range of stream systems and natural (and anthropocentric) environmental variation.

General Work Plan for 2009. We would like to continue our study of fish communities and habitat variables in Thirteen Mile Creek during the summer of 2009 using the same basic approach and trying to sample the same 21 sites. A disadvantage of trying to understand a system like Thirteen Mile Creek is its longitudinal and seasonal variability. Are the parameters measured at 20 sampling sites during the summer representative of processes in the whole system inter-seasonally and inter-annually? The Thirteen Mile Creek system is similar to the Upper Sevenmile Creek system in size and flashiness but different in lacking extensive sedimentation, similar to the O'Fallon Creek system in flashiness but much smaller in scale, and similar to the Burns Creek system in scale but lacking the hydrologic continuity provided by perennial groundwater driven baseflow.

Acknowledgments. We would like to thank the Montana Department of Fish, Wildlife, and Parks, particularly Brad Schmitz and Vic Riggs in Miles City, for logistical support of this research. Landowners in the Thirteen Mile Creek area kindly provided access privileges and information without which this study would not have been possible.

**Cover Illustration.** Northern redbelly dace (*Phoxinus eos*): original artwork by Ellen Edmonson, New York Biological Survey (1927-1940) conducted by New York State Conservation Department (predecessor to New York State Department of Environmental Conservation) (Kraft et al. 2006).

### LITERATURE CITED

- Alt D, Hyndman DW. 1986. Roadside geology of Montana. Mountain Press, Missoula, MT. 427 p.
- Armantrout NB (ed.). 1998. Glossary of aquatic habitat inventory terminology. Amer. Fish. Soc., Bethesda, MD. 136 p.
- Barfoot CA. 1993. Longitudinal distribution of fishes and habitat in Little Beaver Creek, Montana. MS thesis, MT State Univ., Bozeman. 66 p.
- Barfoot CA, White RG. 1999. Fish assemblages and habitat relationships in a small northern Great Plains stream. Prairie Nat. 31(2): 87-107.
- Barnes MD. 1997. An exploratory survey of fishes and aquatic habitats in Burns Creek, Dawson and Richland counties, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 14 p.
- Barnes MD. 1999. Studies on population density and longitudinal distribution of fishes in Burns Creek, Dawson and Richland counties, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 21 p.
- Barnes MD. 2004. Fish communities and macrohabitat variables in O'Fallon Creek, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 22 p.
- Barnes MD, Chouinard B, Siegle MR. 2002. Fish communities and macrohabitat variables in O'Fallon Creek, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 21 p.
- Barnes MD, Curtis ZJ, JY Shi, Huang ZJ. 2005. Fish communities and macrohabitat variables in Upper Sevenmile Creek, Dawson County, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 23 p.
- Barnes MD, Frank JC, Ye YX, Lin RA. 2007. Fish communities and habitats in Upper Sevenmile Creek, Dawson County, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 25 p.
- Barnes MD, Frank JC, Ye YX, Lin RA. 2008. Fish communities and habitats in Thirteen Mile Creek, Dawson County, Montana. Montana Department of Fish, Wildlife, and Parks, Miles City, MT. 22 p.
- Barnes MD, Mahlum SK, Huang ZJ. 2006. Fish communities and habitats in Upper Sevenmile Creek, Dawson County, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 25 p.
- Barnes MD, Siegle MR. 2003. Fish communities and macrohabitat variables in O'Fallon Creek, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 23 p.
- Barnes, MD, Silbernagel NA. 2001. Density, longitudinal distribution, and habitat relationships of fish communities in Burns Creek, Dawson and Richland counties, Montana. MT Dept. of Fish, Wildlife, and Parks, Miles City. 32 p.
- Barnes MD, Westlind SK. 2000. Studies on population density and longitudinal distribution of fishes in Burns Creek, Dawson and Richland counties, MT. MT Dept. of Fish, Wildlife, and Parks, Miles City. 27 p.
- Bramblett RG, Zale AV, Johnson TR, Heggem D. 2004. Using fish assemblages as indicators of aquatic ecosystem integrity in Montana prairie streams. Paper presented at: 37<sup>th</sup> Annual Meeting, Montana Chapter, American Fisheries Society, Whitefish, MT, 2-6 Feb. 2004.

- Clancey CG. 1978. The fish and aquatic invertebrates in Sarpy Creek, Montana. MS thesis, MT State Univ., Bozeman. 53 p.
- Clesceri LS, Greenburg AE, Eaton, AD (eds.). 1999. Standard methods for the examination of water and wastewater (20<sup>th</sup> ed.). Washington, D.C.: American Public Health Association. 1325 p.
- DeLury DB. 1947. On the estimation of biological populations. Biometrics 3: 145-167.
- Elser A, Clancey C, Morris L, Georges M. 1978. Aquatic habitat inventory of the Beaver Creek drainage and selected tributaries of the Yellowstone River. MT Dept. of Fish and Game and US Dept. of Interior, Bur. Of Land Management, Miles City, MT. 136 p.
- Elser AA, Georges MW, Morris LM. 1980. Distribution of fishes in southeastern Montana. MT Dept. of Fish, Wildlife, and Parks and US Dept. of Interior, Bur. of Land Management, Miles City, MT. 100 p.
- Gould WR. 1998. Key to the fishes of Montana. Dept. of Biology, MT State Univ., Bozeman. 22 p.
- Holton GD, Johnson HE. 2003. A field guide to Montana fishes. MT Dept. of Fish, Wildlife, and Parks, Helena. 95 p.
- Jones-Wuellner MR, Bramblett, RG. 2004. Distribution, species richness, and predictive modeling of Montana prairie fishes. Paper presented at: 37<sup>th</sup> Annual Meeting, Montana Chapter, American Fisheries Society, Whitefish, MT, 2-6 Feb. 2004.
- Kraft CE, Carlson DM, Carlson M. 2006. Inland Fishes of New York (Online), Version 4.0. Dept. of Natural Resources, Cornell Univ., NY State Dept. Environmental Conservation: <u>http://pond.dnr.cornell.edu/nyfish/fish.html</u>
- McDonald, K. 2003. Montana's prairie stream surveys begin to solve a mystery. Montana Department of Fish, Wildlife & Parks, Fisheries Division, <u>http://fwp.state.mt.us/news/</u> <u>article\_2605.aspx</u>.
- McMahon TE, Zale AV, Orth DJ. 1996. Aquatic habitat measurements. In: Murphy BR, Willis DW (eds.). Fisheries techniques (2<sup>nd</sup> ed.). Amer. Fish. Soc., Bethesda, MD. p. 83-120.
- MFIS (Montana Fisheries Information System). 2009. Montana Fisheries Information System. Montana Dept. of Fish, Wildlife and Parks, Helena: <u>http://nris.mt.gov/wis/data/fisheries.htm</u> and <u>http://maps2.nris.state.mt.us/scripts/esrimap.dll?name=MFISH&Cmd=INST</u>.
- Morris L, Hightower T, Elser A. 1981. An aquatic resources assessment of selected streams in the lower Yellowstone River basin. MT Dept. of Fish, Wildlife, and Parks and US Dept. of Interior, Bur. of Land Management, Miles City, MT. 151 p.
- NASA [National Aeronautics and Space Administration]. 2009. Latitude/longitude plotting. NASA, Mobile Aeronautics Education Laboratory, Glenn Research Center, Cleveland, OH: <u>http://www.grc.nasa.gov/WWW/MAEL/ag/llplot.htm</u>.
- NRIS [Natural Resource Information System]. 2009. Watershed information access, lower Yellowstone, O'Fallon (10100005). MT State Library, Helena: <u>http://nris.state.mt.us/</u>.
- Robins CR, Bailey RM, Bond CE, Brooker JR, Lachner EA, Lea RN, Scott WB. 1991. Common and scientific names of fishes of the United States and Canada (5<sup>th</sup> ed.). Amer. Fish.

Soc., Bethesda, MD. 183 p.

- Simonson TD, Lyons J, Kanehl PD. 1994. Guidelines for evaluating fish habitat in Wisconsin streams. US Forest Service General Tech. Rep. NC-164. 36 p.
- Strahler AN. 1952. Hypsometric (area-altitude) analysis of erosional topography. Bull. Geol. Soc. Am. 63: 1117-1142.
- TerraServer. 2009. Online Imagery. Raleigh, NC: http://www.terraserver.com/home.asp.
- USGS [U.S. Geological Survey]. 2008. Water resources of the United States, surface water information, gauging station 06327000, Upper Sevenmile Creek near Glendive, Montana. USGS, Washington, D.C.: <u>http://www.usgs.gov/</u>.
- WRCC [Western Region Climate Center]. 2009. Period of record general climate summary, Glendive, Montana, station 243581, 1893-2004. National Oceanographic and Atmospheric Administration, WRCC, Desert Research Institute, Reno, NV: <u>http://www.wrcc.dri.edu/</u>.