

**MONTANA FISH, WILDLIFE, & PARKS
FISHERIES DIVISION JOB PROGRESS REPORT**

STATE: MONTANA

**PROJECT TITLE: STATEWIDE FISHERIES
INVESTIGATIONS**

PROJECT NO.: F-113-R-9

**STUDY TITLE: SURVEY AND INVENTORY OF
COLDWATER WARMWATER ECOSYSTEMS**

JOB NO.: 3140

**TITLE: NORTHWEST MONTANA COLDWATER
ECOSYSTEM INVESTIGATIONS**

PROJECT PERIOD:

JULY 1, 2008 THROUGH JUNE 30, 2009

**LAKE KOOCANUSA AND
KOOTENAI RIVER BASIN
BULL TROUT MONITORING REPORT**

Prepared by:

**Mike Hensler
Neil Benson**



***Montana Fish,
Wildlife & Parks***

TABLE OF CONTENTS

EXECUTIVE SUMMARY.....	iii
LIST OF TABLES.....	iv
LIST OF FIGURES.....	vi
ACKNOWLEDGEMENTS.....	vii
INTRODUCTION.....	1
DESCRIPTION OF STUDY AREA.....	2
Kootenai River Drainage.....	2
Libby Dam and Lake Koocanusa.....	4
Fish Species.....	5
STREAM ELECTROFISHING/JUVENILE BULL TROUT ABUNDANCE ESTIMATES.....	6
Grave Creek.....	9
West Fork Quartz Creek.....	12
Pipe Creek.....	13
West Fisher Creek.....	14
Bear Creek.....	15
O'Brien Creek.....	16
Keeler Creek.....	17
Callahan Creek.....	18
Libby Creek Above Falls.....	18
STREAMBED CORING.....	19
Findings.....	22
SUBSTRATE SCORING.....	23
Findings.....	25

BULL TROUT REDD COUNTS.....	26
Grave Creek.....	28
Wigwam Drainage.....	30
Quartz Creek.....	31
Pipe Creek.....	32
Bear Creek.....	33
O'Brien Creek.....	34
West Fisher Creek.....	35
Keeler Creek.....	36
Callahan Creek.....	37
KOOTENAI RIVER ADULT BULL TROUT POPULATION ESTIMATE.....	38
Findings.....	39
LAKE KOOCANUSA GILLNET MONITORING.....	42
Bull Trout.....	43
LITERATURE CITED.....	45

EXECUTIVE SUMMARY

We conducted juvenile bull trout population estimates within reference reaches on index creeks. Juvenile estimates for many of the streams including O'Brien Creek, Bear Creek and Callahan Creek have decreased dramatically since survey began. Substrate scores and substrate coring data are presented.

Bull trout redd counts in Grave Creek and the Wigwam River have significantly increased since 1995. Bull trout redds in Grave Creek in 2004 were lower (48%) than the previous year and have since stabilized. Bull trout redd counts in tributaries downstream of Libby Dam including Quartz, Pipe, Bear, and O'Brien creeks, and the West Fisher River have been variable over the past several years, and have not increased in proportion to bull trout redd counts upstream of Libby Dam in recent years. Prior to redd counts in 2007 and 2008, bull trout redd counts in Keeler Creek exhibited a positive trend since monitoring began. Since 2005, there appears to be a negative trend that is substantially different from a stable population.

MFWP conducted four adult bull trout population estimates below Libby Dam during the period 2004 - 2008. The five population estimates ranged from 176 – 972 bull trout. We believe that these substantial swings in population estimates may be a function of the reservoir population. The thought is that cohorts of bull trout, entrained through Libby Dam, reside below the dam and take advantage of entrained kokanee. Through time these groups of fish age and fade out of the population. We determined the proportion of bull trout residing below Libby Dam between 2004 and 2007 that were entrained ranged from 49.1% to 62.7% and that most of the entrained bull trout originated from the Wigwam River.

The length frequency distribution for bull trout captured in the Libby Dam section of the Kootenai River was fairly consistent among years. The exception was that we captured a higher percentage of juveniles during 2007 and 2008 estimates. The bull trout recaptured in 2008 grew an average of 96.3 mm and gained an average of 1,649.9 g.

We continued monitoring fish populations within the reservoir using spring and fall gill netting and present the results for bull trout and kokanee. Spring gill net catch of bull trout during the period 1975-1989 appeared to exist at equilibrium. Bull trout catch per net in Libby Reservoir significantly increased between 1990 and 2004. The trend appears to have stabilized since about 2005. This coincides with the opening of bull trout harvest initiated in 2004 for Koocanusa. We also found that Grave Creek redd counts and bull trout catch per net in Koocanusa have similar trends. Both indices are useful for determining management direction for the harvest of bull trout from Koocanusa.

LIST OF TABLES

Table 1. Morphometric data presented for Lake Koocanusa.....	4
Table 2. Current relative abundance (A=abundant, C=common, R=rare, N = Not Found) and abundance trend from 1975 to 2007 (I=increasing, S = stable, D = decreasing, U = unknown) of fish species present in Lake Koocanusa and the Kootenai River drainage.....	5
Table 3. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Grave Creek, 1997 - 2008.....	9
Table 4. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of West Fork Quartz Creek, 1997 - 2008.....	12
Table 5. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Pipe Creek, 1999 - 2008.....	13
Table 6. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of West Fisher Creek, 2002 -2008.....	14
Table 7. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Bear Creek, 1999 - 2008.....	15
Table 8. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of O'Brien Creek, 1998 - 2008.....	16
Table 9. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Keeler Creek, 1998 - 2008.....	17
Table 10. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Callahan Creek, 2003 – 2008.....	18
Table 11. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Libby Creek, 2003 – 2008.....	18

Table 12. Mesh size of sieves used to gravimetrically analyze hollow core streambed substrate samples collected from Kootenai River basin tributaries.....	21
Table 13. Median percentage of streambed material smaller than 6.35 mm in McNeil core samples collected from bull trout spawning areas in tributary streams to the Kootenai River basin, 1994 – 2006.....	22
Table 14. Characteristics and ranks for computing substrate scores (modified by Leathe and Enk 1985 from Crouse et al. 1981).....	24
Table 15. Summary of Kootenai Drainage substrate scores the stream sections monitored at juvenile population estimate sites in Kootenai River basin stream, 1998 – 2008.....	25
Table 16. Summary of Kootenai Drainage bull trout spawning site inventories from 1993 - 2008 in the stream sections monitored annually.....	29
Table 17. The sampling dates for adult bull trout marked, recaptured, and the estimated total population and fish per mile in the Kootenai River from Libby Dam downstream to the Fisher River confluence, 2004 - 2008	39
Table 18. Spring sinking gill net summary of bull trout catch per net in Lake Koocanusa 1975 - 2008.....	43

LIST OF FIGURES

Figure 1. Kootenai River Basin (Montana, Idaho and British Columbia, Canada).....	3
Figure 2. Photographs of Glen Lake Irrigation District diversion on Grave Creek, Montana. Left photo shows passive wedge wire screen, right photo shows diversion with upper headgate and crossvane structures.....	10
Figure 3. Footprint of Glen Lake Irrigation diversion, Grave Creek, Montana.....	11
Figure 4. Bull trout redd counts and trend analysis for Grave Creek (including Clarence Creek and Big Sky Creek) 1995-2008.....	28
Figure 5. Bull trout redd counts and trend analysis for the Wigwam River (including Bighorn, Desolation and Lodgepole Creeks) 1995 – 2008.....	30
Figure 6. Bull trout redd counts and trend analysis for Quartz Creek (including West Fork Quartz) 1990-2008.....	31
Figure 7. Bull trout redd counts and trend analysis for Pipe Creek 1990-2008.....	32
Figure 8. Bull trout redd counts and trend analysis for Bear Creek, a tributary to Libby Creek, 1995-2008.....	33
Figure 9. Bull trout redd counts and trend analysis for O’Brien Creek 1991-2008.....	34
Figure 10. Bull trout redd counts in West Fisher Creek, 1993-2008.....	35
Figure 11. Bull trout redd counts and trend analysis for Keeler Creek, a tributary to Lake Creek, 1996-2008.....	36
Figure 12. Bull trout redd counts for Callahan Creek, 2003 - 2008.....	37
Figure 13. Length frequency distribution of bull trout captured below Libby Dam, 2004 through 2007.....	41
Figure 14. Average catch per net of bull trout in spring gill nets at the Rexford site on Libby Reservoir 1975-2008.....	44
Figure 15. Bull trout redd count and number per net trends for Grave Creek and Lake Koocanusa, respectively 1993 – 2008.....	44

ACKNOWLEDGEMENTS

We are thankful to the many people that contributed to this monitoring effort. Neil Benson, Jim Dunnigan, Larry Garrow, Tom Ostrowski, Jay Deshazer, Will Young, Amy Ellingson, Monty Benner, Travis Lorentz, Kimber Hrach, Ryan Sylvester, Brian Stephens, Jared Lampton, Cameron Sinclair and Josh Smith proved invaluable with field data collection. Jim Vashro provided impetus and encouragement throughout the process. Thanks to US Forest service personnel from Canoe Gulch district, Rexford district and others. Thanks to Mark Deleray, Scott Rumsey and Tom Weaver for providing the template for this report.

INTRODUCTION

The bull trout that inhabit Lake Koocanusa and Kootenai River represent geographically distinct and important populations within their range. Montana Fish, Wildlife & Parks list bull trout as a species of special concern and in 1996 the United States Fish and Wildlife Service (USFWS), through the Endangered Species Act, listed bull trout as threatened throughout their range.

Forestry practices are the dominant land use in all bull trout core areas and represent the highest risk to bull trout in the middle Kootenai (Libby Dam to Kootenai Falls). This risk to the bull trout population in the middle Kootenai is elevated due to the low number of spawning streams (Quartz, Pipe, O'Brien, Callahan and Libby Creek drainages) available; a direct result of habitat fragmentation caused by Libby Dam. The Kootenai River is a nodal habitat containing critical over-wintering areas, migratory corridors, and habitat required for reproduction and early rearing.

Dam operations are considered a very high risk to the continued existence of the Kootenai drainage population of bull trout (Montana Bull Trout Scientific Group 1996a). Dam operations represent a direct threat to bull trout in the middle Kootenai because of the biological affects associated with unnatural flow fluctuations and real potential gas supersaturation problems arising from spilling water. The dam is a fish barrier, generally restricting a portion of this migratory population to 29 miles of river between Libby Dam and Kootenai Falls.

In the upper Kootenai (above Libby Dam), the threats to bull trout habitat include illegal fish introduction, introduced fish species, rural residential development, and forestry. Additional risks come from mining, agriculture, water diversions, and illegal harvest (Montana Bull Trout Scientific Group 1996b). Critical spawning streams include the Grave Creek drainage in the U.S. and the Wigwam drainage in British Columbia. Transboundary research is ongoing in B.C. tributaries: Elk River, St. Mary River, Skookumchuck Creek, White River, Palliser River, and the Kootenay River upstream (Baxter and Oliver 1997). Nodal habitats for this population are provided in Lake Koocanusa, Tobacco River, and the Kootenay River in Canada.

Bull trout are found below Kootenai Falls in O'Brien Creek, Callahan Creek and in Bull Lake. The latter is a disjunct population that migrates out of Bull Lake, downstream to Lake Creek then upstream in Keeler Creek. These fish inhabit areas in the lower Kootenai River and Kootenay Lake during most of the year.

It is the intention of MFWP to manage bull trout populations as sport fisheries in the future. For this to happen, relevant population information must be compiled. This report will help to provide MFWP and other decision makers with the best available information for bull trout populations in the Kootenai River system. In an effort to maintain consistent survey and analysis throughout the region, we reproduced effort initiated in the Flathead drainage. Much of the background information for this report is excerpted, with thanks, from Deleray et al. (1999).

DESCRIPTION OF STUDY AREA

Kootenai River Drainage

The Kootenai River basin is an international watershed that encompasses parts of British Columbia (B.C.), Montana, and Idaho (Figure 1). The headwaters of the Kootenai River originate in Kootenay National Park, B.C. The river flows south within the Rocky Mountain Trench to the reservoir created by Libby Dam, which is located near Libby, Montana. From the reservoir, the river turns west, passes through a gap between the Purcell and Cabinet Mountains, enters Idaho, and then loops north where it flows into Kootenay Lake, B.C. The waters leave the lake's West Arm and flows south to join the Columbia River at Castlegar, B.C. In terms of runoff volume, the Kootenai is the second largest Columbia River tributary. In terms of watershed area (36,000 km² or 8.96 million acres), it ranks third (Knudson 1994).

Nearly two-thirds of the 485-mile-long channel, and almost three-fourths of the Kootenai watershed is located within the province of British Columbia. Roughly twenty-one percent of the watershed lies within Montana (Figure 1), and six percent is in Idaho (Knudson 1994). The Continental Divide forms much of the eastern boundary, the Selkirk Mountains the western boundary, and the Cabinet Range the southern. The Purcell Mountains fill the center of the river's J-shaped course to Kootenay Lake. Throughout, the basin is mountainous and heavily forested.

Libby Reservoir (Lake Koocanusa) and its tributaries receive runoff from 47 percent of the Kootenai River drainage basin. The reservoir has an annual average inflow of 10,615 cfs. Three Canadian rivers, the Kootenay, Elk, and Bull, supply 87 percent of the inflow (Chisholm et al. 1989). The Tobacco River and numerous small tributaries flow into the reservoir south of the International Border.

Major tributaries to the Kootenai River below Libby Dam include the Fisher River (838 sq. mi.; 485 average cfs), Yaak River (766 sq. mi. and 888 average cfs) and Moyie River (755 sq. mi.; 698 average cfs). Kootenai River tributaries are characteristically high-gradient mountain streams with bed material consisting of various mixtures of sand, gravel, rubble, boulders, and drifting clay and silt, predominantly of glacial/lacustrine origin. Fine materials, due to their instability during periods of high stream discharge, are continually eroded and re-deposited as gravel bars, forming braided channels with alternating riffles and pools.

Streamflow in unregulated tributaries generally peaks in May and June after the onset of snow melt, then declines to low flows from November through March. Flows also peak with rain-on-snow events. Kootenai Falls, a 20-foot-high waterfall and a natural fish-migration barrier, is located eleven miles downstream of Libby, Montana.

Libby Dam and Lake Koocanusa

Lake Koocanusa was created under an International Columbia River Treaty between the United States and Canada for cooperative water development of the Columbia River Basin (Columbia River Treaty 1964). Lake Koocanusa inundated 109 stream miles of the mainstem Kootenai River in the United States and Canada, and 40 miles of tributary streams in the U.S. that provided habitat for spawning, juvenile rearing, and migratory passage for salmonids.

Libby Dam is a 113-m (370-ft) high concrete gravity structure with three types of outlets: sluiceways (3), operational penstock intakes (5 operational, 8 possible), and a gated spillway. The dam crest is 931 m long (3,055 ft), and the widths at the crest and base are 16 m (54 ft) and 94 m (310 ft), respectively. A selective withdrawal system was installed at Libby Dam to allow for temperature-controlled release of water from the reservoir.

Completion of Libby Dam in 1972 created the 109-mile Lake Koocanusa. Specific morphometric data for Lake Koocanusa are presented in Table 1. Filling Lake Koocanusa inundated and eliminated 109 miles of the mainstem Kootenai River and 40 miles of critical, low-gradient tributary habitat. This conversion of a large segment of the Kootenai River from a lotic to lentic environment changed the aquatic community (Paragamian 1994). Replacement of the inundated habitat and the community of life it supported are not possible. However, mitigation efforts are underway to protect, reopen, or reconstruct the remaining tributary habitat to offset the loss. Fortunately, in the highlands of the Kootenai Basin, tributary habitat quality is high. The headwaters are relatively undeveloped and retain a high percentage of their original wild attributes and native species complexes. Protection of these remaining pristine areas and reconnection of fragmented habitats are high priorities for bull trout and other native species.

Table 1. Morphometric data presented for Lake Koocanusa Morphometric data.

Surface elevation	
maximum pool	749.5 m (2,459 ft)
minimum operational pool	697.1 m (2,287 ft)
minimum pool (dead storage)	671.2 m (2,222 ft)
Area	
maximum pool	188 sq. km (46,500 acres)
minimum operational pool	58.6 sq. km (14,487 acres)
Volume	
maximum pool	7.24 km ³ (5,869,400 acre-ft)
minimum operational pool	1.10 km ³ (890,000 acre-ft)
Maximum length	145 km (90 mi)
Maximum depth	107 m (350 ft)
Mean depth	38 m (126 ft)
Shoreline length	360 km (224 mi)
Shoreline development	7.4 km (4.6 mi)
Drainage area	23,271 sq. km (8,985 sq. mi)

Fish Species

Eighteen species of fish are present or have been found in Koocanusa Reservoir and the Kootenai River drainage (Table 2). The reservoir currently supports an important fishery for kokanee *Oncorhynchus nerka* and rainbow trout *Oncorhynchus mykiss* (Kamloops strain), and a bull trout fishery that was re-opened in 2004 (Hensler and Benson, 2007). The annual fishing pressure ranges from 30,000 to over 100,000 angler days. The Kootenai River below Libby Dam is a “blue ribbon” rainbow trout fishery, and the state record fish was harvested there in 1997 (over 33 pounds). Bull trout *Salvelinus confluentus* are captured “incidentally”.

Table 2. Current relative abundance (A=abundant, C=common, R=rare, N = Not Found) and abundance trend from 1975 to 2007 (I=increasing, S = stable, D = decreasing, U = unknown) of fish species present in Lake Koocanusa and the Kootenai River drainage.

Common Name	Scientific name	Relative Abundance Reservoir	Trend	Relative Abundance drainage	Trend	Native
<u>Game fish species</u>						
Westslope cutthroat trout	<i>Oncorhynchus clarki lewisi</i>	R	D	C	S	Y
Rainbow trout	<i>Oncorhynchus mykiss</i>	R	D	C	S	Y
Bull trout	<i>Salvelinus confluentus</i>	C	I	C	I	Y
Brook trout	<i>Salvelinus fontinalis</i>	R	U	A	S	N
Lake trout	<i>Salvelinus namaycush</i>	N	U	R	D	N
Kokanee salmon	<i>Oncorhynchus nerka</i>	A	U	R	U	N
Mountain whitefish	<i>Prosopium williamsoni</i>	R	D	A	S	Y
Burbot	<i>Lota lota</i>	R	D	R	D	Y
Largemouth bass	<i>Micropterus salmoides</i>	R	U	R	U	N
White sturgeon	<i>Acipenser transmontanus</i>	R	D ^ε	R	D	Y ^ε
Northern pike	<i>Esox lucius</i>	R	U	R	U	N
Yellow perch	<i>Perca flavescens</i>	C	I	R	U	N
<u>Non-game fish species</u>						
Pumpkinseed	<i>Lepomis gibbosus</i>	R	U	R	U	N
Redside shiner	<i>Richardsonius balteatus</i>	R	D	C	U	Y
Peamouth	<i>Mylocheilus caurinus</i>	A	I	C	U	Y
Northern pikeminnow	<i>Ptychocheilus oregonensis</i>	A	S	C	U	Y
Largescale sucker	<i>Catostomus macrocheilus</i>	A	S	C	U	Y
Longnose sucker	<i>Catostomus catostomus</i>	C	D	R	U	Y

^ε Five white sturgeon were relocated from below Libby Dam to the reservoir. At least one of these fish moved upriver out of the reservoir while two have been accounted for from angler reports and one verified mortality.

^ε An abundance of anecdotal reports exist of white sturgeon above Kootenai Falls although research to date has failed to validate any reports.

STREAM ELECTROFISHING/ JUVENILE BULL TROUT ABUNDANCE ESTIMATES

Introduction

Estimation of fish population abundance is necessary for understanding basic changes in numbers, species composition and year class strength. Direct enumeration is the most accurate technique, but in most situations indirect methods must be employed. We generally use a combination of techniques in order to minimize errors. Fish populations are dynamic and may fluctuate considerably, even over relatively short periods of time, regardless of human influence. Consequently, managers seeking to assess the effects of various activities on fish populations must understand the nature and causes of such fluctuations as fully as possible. We used the protocols similar to those developed to assess fish abundance in the Flathead Basin using electrofishing techniques (Shepard and Graham 1983). Monitoring focuses on quantifying yearly variation of fish abundance in stream sections sampled consistently year after year. We recommend using electrofishing techniques to assess fish abundance in accessible streams because:

1. The precision of electrofishing can be estimated and reported, providing a measure of reliability;
2. There is less bias associated with changes in field personnel; and
3. Estimates derived using electrofishing techniques are presently more accepted by fisheries professionals.

Two-pass Assumptions (Seber and LeCren 1967):

1. Probability of capture (p) is large enough to have a significant effect upon population total (N).

We can test this assumption by computing (p) after two passes are complete. If p is less than 0.5, assumption 1 probably has been violated (Junge and Libovarsky 1965) and more effort is required. We recommend (p) should be 0.6 or larger.

2. Probability of capture is constant. Fishing effort is the same for both catches and fish remaining after the first pass are as vulnerable to capture as were those that were caught in the first pass.

Assumption 2 has frequently been found to be faulty when electrofishing (Lelek 1965, Gooch 1967, Cross and Stott 1975, Mahon 1980). White et al. (1982) found if p was 0.8 or larger, two-catch estimates were reliable because failure of constant probability of capture (assumption 2) did not matter. We found that as long as p was 0.6 or larger and stream discharge was less than 20 cfs, estimates computed using two-catch estimators were similar to mark-recapture estimates.

Zippin (1958) determined that if the probability of capture (p) decreases with subsequent collections, the estimate was an underestimate of the true population size. These estimates may still be reported, but should be used cautiously. They can be used to compare trends in population abundance, provided the same techniques are used throughout the monitoring program.

3. There is no recruitment, mortality, immigration or emigration between the times of the two collections.

Assumption 3 can be easily met, since both electrofishing collections take place within a single day and the section is isolated using block nets.

4. The first catch is removed from the population or, if returned alive, the individuals are marked so they can be ignored when counting the second catch.

This assumption can be met by removing the first catch from the population.

Bull trout fry are exceedingly difficult to capture by electrofishing. There are several reasons for this:

- 1: Their small surface area makes effective, efficient, repeatable shocking difficult
- 2: Their small size (usually 35 to 50 mm at time of estimates) makes seeing them difficult
- 3: Because of their small size there is a high likelihood they will slip through nets during the estimate
- 4: Because of their small size there is a high likelihood that they will slip through the block nets.

We felt that these reasons led to too much probability to violate our capture assumptions, especially 1,2,and 3. We therefore chose not to include fry in the yearly estimates. We captured a representative sample of bull trout fry and included measurements on the field sheets.

Methods

We incorporated the following fish abundance monitoring guidelines for Kootenai drainage estimates:

1. In streams less than 10 cfs, we used two-pass electrofishing technique. In these small streams adequate numbers of fish were captured using a backpack mounted generator-Variable Voltage Pulsator combination. Probability of capture (p) should be higher than 0.6 to obtain reliable results.
2. In streams 10 to 20 cfs, we used two-pass electrofishing estimation. We used two backpack mounted shocking units. If the p -value falls below 0.6 for a sample site, more effort (third pass) should be made instead of simply reporting the two-catch estimate.
3. In streams larger than 20 cfs, two-pass electrofishing technique was used. We used boat mounted shocking equipment (two cathodes) and, when necessary, backpack mounted equipment simultaneously for these sample sections.

Two-pass Procedure:

We placed a braided nylon block net (1/4 inch mesh) at the lower boundary of the shocking section. When using a block net, we placed the net in the stream with the bottom edge facing upstream and place rocks on the weighted (bottom) edge of the net to hold it in position. We tied the ropes along the top edge of the net to a tree (or any available stable item) on each bank to stretch the net tight and hold it perpendicular to the flow. Rocks placed along the entire bottom edge of the net ensure no fish move under the net. Rebar cut to approximately 1.0 m lengths supported the net upright.

We chose sample sections based on accessibility and proximity to redds that were found in previous years. Though we kept sample sites consistent, section length was not consistent between sites or between years due to considerable shifting of streambeds during some years. Section lengths typically were based on riffle breaks at the top of sections and pools at the bottom.

We sampled each section from the upstream boundary to the lower block net. We found that downstream electrofishing was more efficient than upstream electrofishing, and if two passes were needed for each catch (to provide a reliable estimate), both passes should be downstream. It is important to extend equal efforts during each pass, so that if two passes were used for the first catch; two passes must also be completed for the second catch. Mahon (1980) believed longer time periods between catches improved the accuracy of catch per unit effort estimators. For this reason, we recommend some time between collections. During this time, we worked all fish captured on the first pass.

Two-Pass Estimators:

We used the following formula to estimate population number (Seber and LeCren 1967):

$$N = \frac{C_1^2}{C_1 - C_2}$$

Where N = population size at the time of first pass

C_1 = number of fish > 1+ captured during first pass (by species)

C_2 = number of fish > 1+ captured during second pass (by species)

Variance of the estimate:

$$V(N) = \frac{C_1^2 C_2^2 (C_1 + C_2)}{(C_1 - C_2)^4}$$

Probability of capture (p):

$$p = \frac{C_1 - C_2}{C_1}$$

As stated previously, p should be ≥ 0.6 for a reliable, two-pass estimate to be made. Though there were instances when time constraints made a third pass problematic, if $p \leq 0.6$, the estimate was reported, but must be viewed with caution. If $p \geq 0.6$ we completed the estimate; otherwise, generally more fishing effort was expended. This effort can be expended for computing a multiple estimate (by completing additional electrofishing and computing a multi-catch estimate using formulas presented in Zippin 1958). Population estimates and associated 95% confidence intervals were estimated using *Microfish 2.2* (Van Deventer and Platts 1983).

When reporting the estimates of fish numbers computed by electrofishing, we reported the estimate, the 95 percent confidence interval, the area of the section surveyed, the date, and the density and number of mortalities. When reporting two-pass estimates, we reported the probability of capture (p) with the estimate.

Findings

Grave Creek

The Grave Creek fish abundance section is the only section in the U.S. portion of Lake Koocanusa. It is located just upstream of Clarence Creek and has varied from 190 m to 220 m in length. It is a relatively stable section but has been affected periodically by high flows and beaver activity. We have electrofished this section annually since 1997 (Table 3). Redd counts increased dramatically in this tributary since 1997 (Table 16), but densities of juveniles have not shown similar results.

Table 3. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Grave Creek, 1997 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
Grave Creek	1997	158	+/- 12	0.72	9.7
	1998	186	+/- 9	0.77	11.4
	1999	139	+/- 27	0.57	8.5
	2000	160	+/- 17	0.51	9.8
	2001	165	+/- 18	0.67	11.6
	2002	116	+/- 15	0.66	8.5
	2003	156	+/- 19	0.75	15.6
	2004	153	+/- 10	0.83	13.3
	2005	153	+/- 17	0.76	14.5
	2006	107	+/- 12	0.69	8.8
	2007	145	+/- 10	0.76	12.7
	2008	127	+/- 8	0.77	11.4

In 1998 MFWP initiated a survey to determine entrainment of bull trout through the Glen Lake Irrigation ditch (unpublished). The irrigation ditch is located approximately 7 miles upstream from the confluence with Fortine Creek. The diversion is located downstream from the juvenile estimate section. We installed a screw trap approximately 100 m. down the ditch from the control gate for the diversion. In 1998, 100 percent of the bull trout captured (32) in the screw trap were 1+ and older. In 2001 Montana Fish, Wildlife & Parks and Glen Lake Irrigation District installed a passive screen at the entrance to the ditch and installed an additional headgate upstream (Figures 2 and 3). The screen first functioned on May 12, 2001.

The screen was composed of wedge wire panels with 3/8 inch spacing (Figures 2 and 3). Though the number of entrained bull trout age 1+ and older decreased, during this same time we also saw an increase in Young-of-year bull trout caught. In 2001, we found 204 bull trout in the trap of which ten were 1+ and older; and in 2002, only one of 178 trapped was 1+. The trend continued through 2008 when 9 of 744 bull trout captured were age 1+ none were older. These results lead us to believe that under current habitat conditions, Grave Creek may be near or at carrying capacity for juvenile bull trout and young of year bull trout are forced to out migrate. Although at least for the juvenile estimate section, out migration of young of year bull trout does not appear to have a negative effect on the population of 1+ juveniles. Our assessment of the operation of the diversion structure is an ongoing process that we will evaluate more fully and is beyond the scope of this report but contained in Dunnigan et al. (2009)



Figure 2. Photographs of Glen Lake Irrigation District diversion on Grave Creek, Montana. Left photo shows passive wedge wire screen, right photo shows diversion with upper headgate and crossvane structures.




Figure 3. Footprint of Glen Lake Irrigation diversion, Grave Creek, Montana.

West Fork Quartz Creek

The West Fork Quartz Creek fish abundance section is located at the FS 399 bridge. The section has varied in length from 165 m to 248 meters due to spring flows and downfall from wind events. We chose West Fork Quartz rather than mainstem Quartz Creek because we found the majority of redds from year to year are in that tributary. Densities of juvenile bull trout generally increased between 1997 and 2002 then decreased to a relatively stable level since then (Table 4).

Table 4. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of West Fork Quartz Creek, 1997 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
West Fork Quartz Creek	1997	76	+/- 1	0.94	5.4
	1998	82	+/- 5	0.74	6.6
	1999	Not Sampled			
	2000	87	+/- 14	0.60	9.2
	2001	89	+/- 9	0.67	7.4
	2002	89	+/- 4	0.77	10.6
	2003	70	+/- 6	0.67	7.6
	2004	72	+/- 6	0.81	7.9
	2005	64	+/- 10	0.76	7.3
	2006	57	+/- 7	0.73	6.7
	2007	63	+/- 3	0.86	6.7
	2008	98	+/- 7	0.76	10.2

The trend is similar to but less dramatic than that of redd counts (Figure 4) and is likely due to a number of factors that include considerable lower water years since 1997 and flow operations from Libby Dam that resulted in major spill events in 2002 and 2006.

Pipe Creek

The Pipe Creek fish abundance section is located approximately 3 miles below the confluence with East Fork Pipe Creek. We have found redds above and below the section.. The section has varied in length from 147 to 206 meters due to changes caused by spring flows and downfall from wind events. Densities of juvenile bull trout remained relatively stable to slightly decreasing between 1999 and 2005 (Table 5). This occurred as redd counts decreased substantially in 1999 and 2005 likely from low water conditions, and periodic manmade/natural dams below the spawning sites. We caught no juvenile bull trout in Pipe Creek in 2007.

Table 5. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Pipe Creek, 1999 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
Pipe Creek	1999	31	+/- 1	0.76	2.2
	2000	54	+/- 9	0.68	3.8
	2001	23	+/- 4	0.76	2.1
	2002	18	+/- 1	0.71	1.8
	2003	24	+/- 4	0.77	2.2
	2004	22	+/- 2	0.85	1.6
	2005	12	No Recaptures		1.0
	2006	7	+/- 2	0.78	0.6
	2007	0	-	-	-
	2008	One bull trout caught			0.1

West Fisher Creek

West Fisher Creek was sampled for the first time in 2002. The section is centered on the FS 231 road bridge and was 207 meters long and averaged 7.6 meters in width. Though densities were low, the 2002 estimate of 37 juvenile bull trout was unexpected because of extremely low redd counts and low water during 2001 and 2002 (Table 6). About one-half of the juveniles counted were from the adults that spawned in 2000. The trend continued downward through 2005 and increased again in 2006 and 2007. Much of the decreases is likely due to drought over the last several years that caused low flows and elevated temperatures in the downstream tributaries. The increase in spawning in 2004 (Table 16) appears to have had a positive influence on juvenile densities in 2007.

Table 6. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of West Fisher Creek, 2002 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
West Fisher Creek	2002	37	+/- 2	0.75	2.0
	2003	9	+/- 2	0.81	0.6
	2004	5	+/- 1	0.83	0.3
	2005	2	No Recaptures		0.1
	2006	8	+/- 3	0.73	0.4
	2007	31	+/- 5	0.77	1.6
	2008	3	+/- 1	0.75	0.1

Bear Creek

The Bear Creek fish abundance section is centered on the FS 278 bridge. The section has varied in length from 132 to 213 meters due to changes caused by spring flows and downfall from wind events. This is a relatively stable stretch of stream although there have been some pool changes. Densities of juvenile bull trout increased substantially between 1999 and 2002 (Table 7). We believe the dramatic decrease in 2002, 2004, 2005 and 2008 were caused by low water. Several sections of Libby Creek, to which Bear Creek is a tributary, dried by late July in 2001. The same occurred in 2002 so we expect similar, if not lower densities. Montana Fish, Wildlife & Parks special projects is working with private, corporate and public landholders to reconstruct portions of Libby Creek in hopes that the complete loss of stream flow during low water years can be minimized. This is a slow and laborious process that will likely take 10's of years.

Table 7. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Bear Creek, 1999 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
Bear Creek	1999	101	+/- 9	0.73	8.5
	2000	103	+/- 3	0.87	12.1
	2001	80	+/- 9	0.72	14.0
	2002	67	+/- 3	0.85	6.2
	2003	108	+/- 10	0.79	8.4
	2004	46	+/- 7	0.77	2.6
	2005	79	+/- 17.6	0.69	3.8
	2006	69	+/- 7	0.75	3.5
	2007	17	+/- 1	0.94	1.3
	2008	8	+/- 1	0.89	0.4

O'Brien Creek

O'Brien Creek is one of two tributaries below Kootenai Falls that support bull trout spawning from the Kootenai River (there are bull trout in Callahan Creek, although we have not been successful in trapping upstream migrating adults). The O'Brien Creek fish abundance section currently is centered on the FS 331 bridge above Rabbit Creek. The section has remained stable at 140 meters since the initial survey in 1998. This is a relatively stable stretch of stream with little change in pools from year to year. Densities of juvenile bull trout decreased dramatically between 1998 and 2002 and have remained extremely low since then (Table 8). We could not do estimate in 2002 or 2005 because we got no recaptures.

We believe that the decrease in juvenile densities was caused by a combination of an encroaching beaver population and low water. Redd counts remained relatively stable over this time but the distribution of redds shifted downstream because beaver dams have caused very high sedimentation in some traditional spawning areas. Additionally, estimates of resident brook trout and westslope cutthroat trout and rainbow trout were similar throughout the survey years (unpublished data).

Table 8. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of O'Brien Creek, 1998 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
O'Brien Creek	1998	91	+/- 4	0.84	13.2
	1999	29	+/- 1	0.88	4.2
	2000	21	+/- 7	0.66	3.0
	2001	11	+/- 2	0.61	1.6
	2002	2	No Recaptures		0.3
	2003	5	+/- 1	0.83	0.5
	2004	16	+/- 1	0.89	2.1
	2005	2	No Recaptures		0.3
	2006*	31	+/- 1	0.97	3.3
	2007	5	0	1.0	0.5
	2008	11	+/- 1	0.92	1.1

* Estimate section was moved downstream

Keeler Creek

Bull trout that spawn in Keeler Creek (including the North, South and West Forks) are adfluvial and migrate downstream out of Bull Lake into Lake Creek, then upstream into Keeler Creek. This downstream spawning migration is somewhat unique when compared to other bull trout populations (Montana Bull Trout Restoration Team 1996a). Lake Creek, a tributary of the Kootenai River, has an upstream waterfall barrier isolating this population from the mainstem Kootenai River population. A micro-hydropower dam constructed in 1916 covered the upper portion of the waterfall. A series of high gradient waterfalls are still present below the dam, and are barriers to all upstream fish passage. Keeler Creek likely supplies some recruitment to the Kootenai River through downstream migration.

The Keeler Creek fish abundance section located approximately 1 mile below North Fork Keeler Creek. The section lengths remained fairly constant between 203 and 214 meters since the initial survey in 1998. This is a relatively stable stretch of stream with little change in pools from year to year. The top of the section is controlled by a rock out crop and the bottom is a very stable pool. Densities of juvenile bull trout remained very stable between 1999 and 2002 (Table 9). The estimate in 1998 was an exception. We captured more than twice as many bull trout that year as others. One explanation might be that the flows were very high that year and more juveniles passed into this stable section that includes two fairly large pools. Continued low estimates since 2002 likely reflect low water and beaver activity.

Table 9. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Keeler Creek, 1998 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
Keeler Creek	1998*	159	+/- 50	0.33	7.7
	1999	65	+/- 16	0.69	3.3
	2000	61	+/- 41	0.42	3.1
	2001*	66	+/- 12	0.50	3.0
	2002	74	+/- 13	0.73	3.9
	2003	63	+/- 11	0.74	3.4
	2004	27	+/- 2	0.84	1.4
	2005	18	+/- 1	0.95	1.1
	2006	32	+/- 2	0.87	1.6
	2007	38	+/- 1	0.93	2.4
	2008	27	+/- 2	0.97	1.4

* Three pass estimates.

Callahan Creek

The Callahan Creek fish abundance section is actually located on North Callahan Creek just above the confluence with South Callahan Creek and is located above the FS 414 bridge. The section has varied in length from 154 m to 183 meters due to spring flows and downfall from wind events. We chose North Callahan Creek rather than mainstem Callahan Creek because we found the majority of redds from year to year are above that tributary and logistics of access to mainstem was difficult at best. Densities of juvenile bull trout are quite variable in this section (Table 10). There was a 20+ year flood event during November of 2006 that may have affected population densities in 2007 and 2008. Idaho Fish and Game has monitored redds for this stream and we will continue to track juvenile estimates.

Table 10. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Callahan Creek, 2003 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
Callahan Creek	2003	10	+/- 1	0.83	0.7
	2004	43	+/- 4	0.82	2.4
	2005	35	+/- 4	0.79	2.2
	2006	11	+/- 3	0.85	0.6
	2007	4	+/- 1	0.80	0.2
	2008	Caught 11 bull trout all less than 64 mm (young-of-year)			

Libby Creek Above Falls

The Libby Creek fish abundance section is located above Libby Falls below the Montenore mine site. We have monitored bull trout abundance at this site since 2003 and the section has remained at 150 meters. This particular population is an isolated resident population separated from upstream movement by Libby Falls (approximately 60 ft). The estimate has likely included resident adults. Densities of bull trout varied little from 2003 to 2008 with the exception of 2005 (Table 11). We will continue to monitor this unique population as it is downstream of a proposed adit mine.

Table 11. Population estimates (N), 95 percent confidence intervals (95% C.I.), probability of first pass capture (p) and densities for Age 1 and older bull trout calculated from electrofishing in the permanent section of Libby Creek, 2003 - 2008.

Stream	Year	N	95 % C.I.	p	Density (#100m ²)
Libby Creek Above Libby Falls	2003	27	+/- 2	0.90	3.0
	2004	No estimate			
	2005	55	+/- 8	0.72	5.2
	2006	23	+/- 5	0.77	2.3
	2007	25	+/- 8	0.63	2.1
	2008	33	+/- 3	0.89	2.4

STREAMBED CORING

Introduction

Successful egg incubation and fry emergence are dependent on gravel composition, gravel permeability, water temperature, and surface flow conditions. The female trout begins redd construction by digging an initial pit or depression in the streambed gravel with her tail. After the spawning pair deposits eggs and sperm into this area, the female moves upstream a short distance and continues the excavation, covering the deposited eggs. The process is then repeated several more times, resulting in a series of egg pockets formed by the upstream progression of excavations. The displaced gravel mounds up, covering egg pockets already in place. After egg deposition is complete the female creates a large depression at the upstream edge of the redd. This enhances intra-gravel flow and displaces more gravel back over the entire spawning area. Excavation of the redd causes fine sediments and organic particles to be washed downstream, leaving the redd environment with less fine material than the surrounding substrate. Weather, streamflow, and transport of fine sediment and organic material in the stream can change conditions in redds during the incubation period. Redds can be disturbed by other spawning fish, animals, human activities, or by high flows which displace streambed materials (Chapman 1988).

Redd construction by migratory bull trout in the Flathead drainage disturbs the streambed to a depth of at least 18.0 to 25.0 cm (Weaver and Fraley 1991). Egg pockets of smaller fish (brook trout) tend to be shallower. The maximum depth of gravel displacement is indicative of egg deposition depth (Everest et al. 1987). Freeze coring documented larger substrate particles (up to 15.2 cm) at the base of egg pockets than in overlying substrates (Weaver and Fraley 1991). These particles are likely too large for the female to dislodge during redd construction. Eggs are deposited and settle around these larger particles (Chapman 1988). Continued displacement of streambed materials by the female then covers the eggs.

Redds become less suitable for incubating embryos if fine sediments and organic materials are deposited in interstitial spaces of the gravel during the incubation period. Fine particles impede movement of water through the gravel, thereby reducing delivery of dissolved oxygen to, and flushing of metabolic wastes away from incubating embryos. This results in lower survival (Wickett 1958; McNeil and Ahnell 1964; Reiser and Wesche 1979). For successful emergence to occur fry need to be able to move within the redd, but high levels of fine sediment can restrict their movements (Koski 1966; Bjornn 1969; Phillips et al. 1975). In some instances, embryos that incubate and develop successfully can become entombed (trapped by fine sediments). Sediment levels can alter timing of emergence (Alderdice et al. 1958; Shumway et al. 1964) and affect fry condition at emergence (Silver et al. 1963; Koski 1975).

Measurements of the size range of materials in the streambed are indicative of spawning and incubation habitat quality. In general, research has shown negative relationships between fine sediment and incubation success for salmonids that construct redds (Chapman 1988). A significant inverse relationship exists between the percentage of fine sediment in substrates and survival to emergence of westslope cutthroat trout and bull trout embryos in incubation tests

(Weaver and White 1985; Weaver and Fraley 1991, 1993). Mean adjusted emergence success ranged from about 80 percent when no fine material was present, to less than 5 percent when half of the incubation gravel was smaller than 6.35 mm; about 30 percent survival occurs at 35 percent fines. Entombment was the major mortality factor.

Median percentages of streambed materials smaller than 6.35 mm at fry emergence ranged from 24.8 to 50.3 percent in 29 separate bull trout spawning areas sampled during the Flathead Basin Forest Practice Water Quality and Fisheries Study (Weaver and Fraley 1991). Linear regression of results against output from models assessing ground disturbing activity and water yield increases in these 29 Flathead Basin tributary drainages showed significant positive relationships (Weaver and Fraley 1991). These results demonstrate a linkage between on-the-ground activity and spawning habitat quality. This testing allowed development of models that predict embryo survival to emergence, given the percentage of material smaller than 6.35 mm in the incubation environment. We monitor bull trout spawning and incubation habitat quality by determining the percent fines in a given spawning area through hollow core sampling across years.

Methods

Field crews used a standard 15.2 cm hollow core sampler (McNeil and Ahnell 1964) to collect four samples across each of three transects at each study area. We located actual coring sites at the transects using a stratified random selection process. The total width of stream having suitable depth, velocity, and substrate for spawning was visually divided into four equal cells. We randomly took one core sample in each cell. In some study areas we deviated from this procedure due to limited or discontinuous areas of suitable spawning habitat. We selected study areas based on observations of natural spawning. We only sampled in spawning areas used by adfluvial and fluvial bull trout. During the period of study, these fish spawned in the same general areas, so sampling locations remained similar.

Sampling involved working the corer into the streambed to a depth of 15.2 cm. We removed all material inside the sampler and placed it in heavy duty plastic bags. We labeled the bags and transported them to the Kootenai National Forest Soils Laboratory in Libby, Montana, for gravimetric analysis. We sampled the material suspended in water inside the corer using an Imhoff settling cone (Shepard and Graham 1982). We allowed the cone to settle for 20 minutes before recording the amount of sediment per liter of water. After taking the Imhoff cone sample, we determined total volume of the turbid water inside the corer by measuring the depth and referring to a depth to volume conversion table (Shepard and Graham 1982).

The product of the cone reading (ml of sediment per liter) and the total volume of turbid water inside the corer (liters) yields an approximation of the amount of fine sediment suspended inside the corer after sample removal. We then applied a wet to dry conversion factor developed for Flathead tributaries by Shepard and Graham (1982), yielding an estimated dry weight (g) for the suspended material.

We oven dried the bagged samples and sieve separated them into 13 size classes ranging from >76.1 mm to <0.063 mm in diameter (Table 12). We weighed the material retained on each sieve and calculated the percent dry weight in each size class. The estimated dry weight of the suspended fine material (Imhoff cone results) was added to the weight observed in the pan, to determine the percentage of material <0.063 mm.

Table 12. Mesh size of sieves used to gravimetrically analyze hollow core streambed substrate samples collected from Kootenai River basin tributaries.

76.1 mm	(3.00 inch)
50.8 mm	(2.00 inch)
25.4 mm	(1.00 inch)
18.8 mm	(0.74 inch)
12.7 mm	(0.50 inch)
9.52 mm	(0.38 inch)
6.35 mm	(0.25 inch)
4.76 mm	(0.19 inch)
2.00 mm	(0.08 inch)
0.85 mm	(0.03 inch)
0.42 mm	(0.016 inch)
0.063 mm	(0.002 inch)
Pan	(<0.002 inch)

We refer to each set of samples by using the median percentage <6.35 mm in diameter. This size class is commonly used to describe spawning gravel quality, and it includes the size range typically generated during land management activities. We examined the range of median values for this size class observed throughout the survey area.

Findings

Core sampling in indicator streams has been sporadic since 1994 (Table 13). The current standard for assessing impairment of streams due to increase in sediments continues to be based on fine sediment (<6.35 mm). Weaver and Fraley (1991) found that survival is reduced to one-third when fine sediments reach 35 percent and at 40 percent the survival drops to one-quarter.

Though there is not enough long-term data presented to fully assess trends in the monitored streams, most appear show stable if not decreasing fine sediment levels. Two exceptions are O'Brien Creek and the British Columbia portion of the Wigwam River. Both have been impacted by extra activities in the past several years. O'Brien has increased redd counts but decreased juveniles abundances it is not surprising when viewed in context of the percentage of fine sediments and substrate scores (Tables 13 and 15, respectively).

As was mentioned previously, beavers have become well established in the upper end of O'Brien Creek and are migrating downstream. O'Brien Creek appears to be a fairly high sediment system at the upper end with a large amount of low gradient tortuous stream immediately above the historic spawning areas. Fine sediments may be held back from flushing during high water events and the additional daily activity of the beavers throughout the lower water may release more fine sediments into the stream.

The sediment characteristic in the Wigwam River drainage like most of the bull trout drainages in the Kootenai River basin is a product of natural and anthropogenic disturbances through history. Heavy logging activities in both Montana and British Columbia drainages and 100 year and 200-year flood events have shaped the system in the last 50 years. Oliver and Cope (1999) suggested that "...Frequent lateral channel migrations over time have resulted in erosion of adjacent terraces, coarse sediment delivery to the mainstem river, and have created numerous section of braided channel comprised of sorted gravels and cobbles that provide prime spawning habitat for bull trout". Tepper (2002) found that between 1998 and 2002 the average median of fine sediments (<6.35 mm) increased from 26.5 to 31.7 from the upstream (Montana portion) to downstream (Bighorn Creek) survey sites. It would be advisable to continue monitoring this important tributary as land management activities including considerable new road building occur in the next 10 years.

Table 13. Median percentage of streambed material smaller than 6.35 mm in McNeil core samples collected from bull trout spawning areas in tributary streams to the Kootenai River basin, 1998 – 2006.

Stream	1998	1999	2000	2001	2002	2003	2004	2005	2006
Grave Creek	22.1		25.3	22.9	29.8	28.2	27.4	23.5	23.6
West Fork Quartz Creek	27.5			29.4	27.0	26.4	30.6	24.3	26.3
Pipe Creek	38.1	31.3	31.4	28.9	32.1	35.3	29.7	34.6	28.8
Bear Creek			19.0		27.5	22.0	34.3	34.7	31.3
O'Brien Creek	35.6			35.5	31.5	35.3	34.8	39.8	32.5
North Fork Keeler Creek	27.6		18.6	22.1	26.9	33.1	29.8	27.3	19.9
Wigwam River U.S.	26.5	26.8	24.8				29.6	24.8	26.8

SUBSTRATE SCORING

Introduction

Environmental factors influence distribution and abundance of juvenile bull trout within drainages throughout the range of the species, as well as within specific stream segments (Oliver 1979, Allan 1980, Leathe and Enk 1985, Pratt 1985, Fraley and Shepard 1989, Ziller 1992). Temperature, cover, and water quality regulate general distributions and abundances of juvenile salmonids within drainages. Depth, velocity, substrate, cover, predators, and competitors affect juvenile presence at specific locations in a stream. Although spawning occurs in limited portions of the drainage, juvenile salmonids disperse to occupy most of the areas within the drainage that are suitable and accessible (Everest 1973; Leider et al. 1986).

Juvenile bull trout rear for up to four years in Kootenai Basin tributaries. Snorkel and electrofishing observations during past studies indicate juvenile bull trout are extremely substrate-oriented and can be territorial (Fraley and Shepard 1989). This combination of traits results in partitioning of suitable rearing habitat and a carrying capacity for each stream. We monitor substrate-related habitat potential by calculating substrate scores (Leathe and Enk 1985).

Substrate composition influences distribution of juvenile bull trout and rearing capacities of nursery streams. Sediment accumulations reduce pool depth, cause channel braiding or dewatering, and reduce interstitial spaces among larger streambed particles (Megahan et al. 1980, Shepard et al. 1984, Everest et al. 1987). Juvenile bull trout are almost always found in close association with the substrate (McPhail and Murray 1979, Shepard et al. 1984, Weaver and Fraley 1991). A significant positive relationship existed between substrate score and juvenile bull trout densities in Swan River tributaries (Leathe and Enk 1985) and Flathead River tributaries (Weaver and Fraley 1991), where a high substrate score was indicative of large particle sizes and low score of embeddedness (Crouse et al. 1981). This relationship is thought to reflect substrate types favoring over winter survival (Pratt 1984, Weaver and Fraley 1991).

A substrate score is an overall assessment of streambed particle size and embeddedness. Large particles that are not embedded in finer materials provide more interstitial space that juvenile bull trout favor. This situation generates a higher substrate score. Low substrate scores occur when smaller streambed particles and greater embeddedness limit the interstices within the streambed.

Linear regression of substrate scores against output from a model assessing ground disturbing activity in 28 Flathead Basin tributary drainages showed a significant negative relationship. Researchers also obtained a significant negative relationship between substrate scores and output from a model predicting increases in water yields (Weaver and Fraley 1991). These results demonstrate a linkage between ground disturbance and increased water yield and streambed conditions. Linear regression of juvenile bull trout density against substrate scores in 15 Flathead Basin streams showed a significant positive relationship (Weaver and Fraley 1991). This showed a strong linkage between streambed condition as measured by substrate scoring and actual juvenile bull trout abundance.

Methods

Substrate scoring involves visually assessing the dominant and subdominant streambed substrate particles, along with embeddedness across transects. Surveyors assign a rank to both the dominant and subdominant particle size classes in each cell (Table 14). They also rank the degree to which the dominant particle size is embedded (Table 14). The three ranks are summed, obtaining a single variable for each cell. A mean of all transects in a section results in the substrate score.

Table 14. Characteristics and ranks for computing substrate scores (modified by Leathe and Enk 1985 from Crouse et al. 1981).

Rank	Characteristic
	<u>Particle Size Class</u> ¹
1	Silt and/or detritus
2	Sand (<2.0 mm)
3	Small gravel (2.0-6.4 mm)
4	Large gravel (6.5-64.0 mm)
5	Cobble (64.1-256.0 mm)
6	Boulder and/or bedrock (>256.0 mm)
	<u>Embeddedness</u>
1	Completely embedded or nearly so (75% - 100%)
2	50% - 75% embedded
3	25% – 50% embedded
4	5% – 25% embedded
5	Unembedded
¹ Used for both dominant and subdominant particle ranking	

We obtained the substrate scores using ten equally spaced transects in the juvenile bull trout abundance sections. Again, lower scores indicate poorer quality rearing habitat; higher values indicate good conditions.

Findings

We began collecting substrate scores in 1998 and collected them only sporadically until 2002 (Table 15). We did combine all Kootenai drainage bull trout streams to compare juvenile bull trout population estimates to substrate scores. We found a positive correlation but it was not significant ($P > 0.1$). In the case individual streams, because of limited sampling, a quality assessment was not possible. For the most part, the scores from most of the streams continue to compare favorably with Flathead River basin streams where Flathead Basin Cooperative Forest Practice Study determined that scores of 10.0 or less threatened juvenile bull trout rearing capacity and scores 9.0 or less impaired rearing capacities (Deleray et al. 1999).

Table 15. Summary of Kootenai Drainage substrate scores the stream sections monitored at juvenile population estimate sites in Kootenai River basin stream, 1998 - 2008.

Stream	1998	1999	2002	2003	2004	2005	2006	2007	2008
Grave Creek	13.4		13.2	14.3	13.8	12.8	13.9	13.2	13.1
West Fork Quartz Creek	13.2		13.2	13.3	14.5	14.1	14.1	13.4	13.9
Pipe Creek	13.0	14.0	13.7	12.3	12.1	12.9	12.8	14.1	12.4
Bear Creek		13.0	13.6	14.6	13.8	13.7	13.5	15.5	13.9
West Fisher Creek			13.1	13.8	12.9	14.1	13.3	15.1	13.3
O'Brien Creek	11.5	12.2	10.6	11.9	10.9	11.8	12.3	12.8	12.7
Keeler Creek	12.8	14.4	12.4	13.2	15.5	13.4	14.1	15.5	15.0
Callahan Creek				14.5	15.5	13.9	14.1	14.0	14.8
Libby Creek above falls				14.3		13.6	13.4	15.2	12.8

O'Brien Creek continues to rank lowest of the surveyed streams and also continues to lag behind most core streams in juvenile bull trout abundance, especially considering the relatively high number of redds that are produced each year (Table 16). We changed our population estimate site in 2006 determine if location reflected the population structure in O'Brien Creek or some other factor (environmental or biological) was responsible for this apparent lack of recruitment; fine sediments appear to negatively impact bull trout survival in O'Brien Creek.

BULL TROUT REDD COUNTS

Introduction

A reliable survey of annual spawner escapement is a valuable element of any fisheries monitoring program. These data are frequently used as measures of anticipated production in succeeding generations. They also provide an index of success in regulating the fishery. Observations during past studies indicate that migratory fish populations in the Kootenai System consistently use the same stream sections for spawning. Similar findings resulted from spawning site surveys in the Flathead and Clark Fork River drainages (Montana Fish, Wildlife & Parks, Kalispell, unpublished file data; MBTSG 1996b, 1996c). As a result of specific spawning habitat requirements, the majority of bull trout spawning is clustered in a small portion of the available habitat, making these areas critical to bull trout production.

Field crews annually monitor the number of spawning sites (redds). These counts provided information on trends in escapement into upper basin tributaries and allowed us to choose sampling locations for other monitoring activities. Timing of salmonid spawning has likely evolved in response to seasonal changes in water temperature (Bjornn and Reiser 1991). Initiation of spawning by bull trout appears to be strongly related to water temperature, although photoperiod and streamflow may also be factors (Shepard et al. 1984). Most bull trout spawn between late August and early November (McPhail and Murray 1979; Oliver 1979; Shepard et al. 1984; Pratt 1985; Brown 1992; Ratliff 1992). Spawning in the Flathead drainage (Fraley and Shepard 1989) and in Mackenzie Creek, British Columbia (McPhail and Murray 1979), began when daily maximum water temperatures declined to 9-10° C. Spawning takes place primarily at night (Heimer 1965; Weaver and White 1985), but has been observed during daylight hours (Needham and Vaughan 1952; personal observations).

Bull trout spawning typically occurs in areas influenced by groundwater (Allan 1980; Shepard et al. 1984; Ratliff 1992; Fraley and Shepard 1989). Such areas tend to remain open in the Kootenai drainage during harsh winter conditions, while adjacent stream sections ice over or contain extensive accumulations of anchor ice. Recent investigations in the Swan River drainage found that bull trout spawning site selection occurred primarily in stream reaches that were gaining water from the subsurface, or in reaches immediately downstream of upwelling reaches (Baxter 1997).

Reaches used by spawning adults typically have gradients less than 2 percent (Fraley and Shepard 1989). Water depths at the upstream edges of 80 redds of migratory bull trout in the Flathead drainage ranged from 0.1 to 0.6 m and averaged 0.3 m; water velocities (at 0.6 of the depth below the surface) ranged from 0.09 to 0.61 m/s and averaged 0.29 m/s (Fraley et al. 1981). Similar mean depths (0.3 m) and water velocities (0.31 m/s) at migratory bull trout redds were documented in the Swan River drainage (Kitano et al. 1994).

Migratory bull trout redds ranged from 1.0 to 3.1 m in length (mean 2.1 m) in tributaries of the North and Middle forks of the Flathead River (n=465); width of these redds ranged from 0.8 to 1.5 m and averaged 1.1 m (Fraley et al. 1981). These dimensions are comparable to redds created by fluvial and adfluvial bull trout in the Kootenai drainage.

Areas in which redds are counted on a routine basis are called “index” areas. In some cases these index surveys continue to an upstream barrier. It is important to establish upper and lower limits of index areas. Through repeated annual index surveys we obtain valuable trend information to use in monitoring bull trout populations. Detection of trends will often require at least 10 years of monitoring index areas (Rieman and Meyers 1997).

Methods

We conducted preliminary surveys to determine appropriate timing for final counts. During a basin-wide count we surveyed all habitat that appeared suitable for bull trout spawning (as described above). From this basin-wide survey, index areas were identified for annual surveys. We began final inventories after we observed completed redds, few adult fish, and little evidence of active spawning during the preliminary surveys. Timing of final counts is critical, because as redds age, they lose the characteristic cleaned or bright appearance becoming more difficult to identify. Also, as winter approaches, fall freshets are fairly common in the Kootenai drainage and can wipe out traces of redds if flows get high enough.

We surveyed the Wigwam River and West Fisher, Grave, Quartz, Bear (tributary to Libby Creek), Keeler, Pipe, and O’Brien Creeks. MFWP and U.S. Forest Service (USFS) personnel walked streams in the United States and personnel from the British Columbia Ministry of Water, Land, and Air Protection walked the Wigwam River and associated tributaries. They visually identified redds by the presence of a pit or depression and associated tail area of disturbed gravel. If timing is correct, identification of redds presents little problem. We classified redds differently than in the Flathead. We counted redds only if they were positively identified. We did not include “probable redds” in our counts. We felt that our crews were well trained and confident enough to assess redds as existing or not. We used linear regression to assess population trends.

Findings

Grave Creek

MFWP counted redds in the Grave Creek Basin (including Blue Sky, Clarence, Williams and Lewis Creeks) for the first time in 1983, as well as in 1984, 1985, and 1993 through 2004. Grave Creek was surveyed from its confluence with the Tobacco River upstream to near the mouth of Lewis Creek (approximately 13 miles), where it becomes intermittent. Most redds in Grave Creek were located upstream from the mouth of Clarence Creek to the confluence with Lewis Creek. MFWP found 10 redds between the confluence with the Tobacco River and one mile below Clarence Creek in 1983. However, we did not find redds in this reach during surveys conducted in 1993 and 2000. The distribution of bull trout redds in Blue Sky, Clarence, Williams and Lewis creeks was similar to observations in previous years (Hoffman et al. 2002).

We observed 141 bull trout redds in Grave Creek in 2004, which represented a 48% reduction from the previous year (Table 16) and numbers have since not reached 2003 levels. Nevertheless, bull trout have exhibited a significant positive trend in spawning abundance in Grave Creek since 1993 (Figure 4).




Figure 4. Bull trout redd counts and trend analysis for Grave Creek (including Clarence Creek and Big Sky Creek) 1995-2008.

Table 16. Summary of Kootenai Drainage bull trout spawning site inventories from 1993 - 2007 in the stream sections monitored annually.

Stream	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008
Grave Creek ^a	36	71	15	35	49	66	134	97	173	199	245	141	194	148	208	207
Quartz Creek ^c	89	64	66	47	69	105	102	91	154	62	55	49	71	51	35	46
Pipe Creek	6	7	5	17	26	34	36	30	6	11	10	8	2	6	0	4
Bear Creek			6	10	13	22	36	23	4	17	14	6	3	14	9	14
West Fisher Creek	2	0	3	4	0	8	18	23	1	1	1	13	27	4	18	6
O'Brien Creek	6	7	22	12	36	47	37	34	47	45	46	51	86	65	77	79
Keeler Creek ^d				74	59	92	99	90	13	102	87	126	170	142	84	62
Wigwam River (U.S. & B.C.) ^b		104	247	512	598	679	868	1204	1496	1916	2053	2133	649 ^f	2311	1916	1839
Other (B.C.) ^g										404	373	324	354	474	410	347
Callahan Creek (IDFG) ^e										19	40	25	21	34	3	15
Total				711	850	1053	1330	1592	2060	2776	2924	2876	1577	3249	2760	2619

^a Includes mainstem Grave Creek, Clarence Creek, Blue Sky Creek

^b Includes mainstem Wigwam River, Ram Creek, Lodgepole Creek, Desolation Creek.

^c Includes mainstem Quartz Creek and West Fork Quartz Creek

^d Includes mainstem Keeler Creek, North Fork Keeler Creek, South Fork Keeler Creek.

^e Includes North and South Callahan Creeks

^f Massive slide downstream of traditional count area caused a partial barrier that likely inhibited migration

^g Includes Blackfoot, Skookumchuk and White Rivers.

Wigwam Drainage

Bull trout redd counts for the Wigwam River includes the tributary streams of Bighorn, Desolation, and Lodgepole creeks, and the portion of the Wigwam River within Montana. A large flood event caused a landslide to partially block the Wigwam downstream of the traditional redds count area in 2005. It was obviously a partial barrier for that year and has been passable since. A total of 1916 and 1839 bull trout redds were observed in the Wigwam Drainage in 2007 and 2008, respectively. This was the first time since 1995 in which counts were not higher than the previous year (Figure 5). Anglers have been allowed to harvest bull trout from Koocanusa since 2004. We will continue to monitor redds in the Wigwam River and Grave Creek to identify the potential effect of angling the number of spawning adults to these streams.




Figure 5. Bull trout redd counts and trend analysis for the Wigwam River (including Bighorn, Desolation, and Lodgepole creeks) 1995-2008.

Quartz Creek

Bull trout redd counts in Quartz Creek since 1995 have been variable (Figure 6). Although overall trend is positive, annual variation limits our ability to statistically distinguish this relationship from a stable (zero slope) population. We observed a total of 35 redds in Quartz and West Fork Quartz creeks in 2004 (Table 1). The average number of redds of the period of record was 74.9 redds. The 2004 observation of 49 redds was 34.5% lower than the average over the period of record. A log jam located approximately 0.25 miles upstream of the confluence of West Fork Quartz Creek in 2002 and 2003 may have limited bull trout spawner escapement during these years. This log jam was removed prior to adult bull trout upstream migration in 2004. There was a slight increase in redds found in 2008.




Figure 6. Bull trout redd counts and trend analysis for Quartz Creek (including West Fork Quartz) 1990-2008.

Pipe Creek

Bull trout redd counts in Pipe Creek peaked in 1999 with 36 redds, with redd numbers and have decreased since that peak. Despite the decreasing trend of bull trout redds during the last five years, the overall general trend during the time period 1995-2004 has been variable, with a slope that is not significantly different than a stable population (Figure 7). The mean number of bull trout redds since 1990 has been 14.5 redds. The 8 redds we observed in Pipe Creek in 2004 was 45% lower than the 14 year average. Low water conditions during the fall spawning season during the last several years may partially explain the low spawner escapement into Pipe Creek. During spring 2008 a logjam downstream of the traditional spawning area blew out. This was a possible reason for we found redds in the fall.




Figure 7. Bull trout redd counts and trend analysis for Pipe Creek 1990-2008.

Bear Creek

Bear Creek bull trout redd counts have been variable during the period 1995-2004 (Figure 8). Although the overall general trend has increased since 1995, the relationship is not statistically different than a stable population. Low water conditions in Bear Creek since 2000 also partially explain the low spawner escapement in Bear Creek. The average number of bull trout redds since 1995 in Bear Creek has been 15.1 redds. The 6 redds we observed in Bear Creek in 2004 was 60.3% less than the 9 year average. Figure 8 depicts an increasing trend since 2004, we don't know the effect of entrainment through Libby dam during spill events like 2006 have on bull trout escapement to the tributaries downstream of the dam although it may be an explanation for increases in Bear Creek.




Figure 8. Bull trout redd counts and trend analysis for Bear Creek, a tributary to Libby Creek, 1995-2008.

O'Brien Creek

The trend of bull trout redds in O'Brien Creek is generally increasing since 1996 (Figure 9). We observed a total of 51 bull trout redds in O'Brien Creek in 2004 (Table 14). This is surprising considering the low juvenile survival downstream of the redds. It is quite possible that the adult escapement is more a function of adults moving downstream from Libby Dam over Kootenai Falls (a substantial partial barrier). They likely drop back to the nearest acceptable stream to spawn.




Figure 9. Bull trout redd counts and trend analysis for O'Brien Creek 1991-2008.

West Fisher Creek

We were unable to determine a significant trend in bull trout redd counts in West Fisher Creek over the period of record for this stream (1993-2008). In fact, there appears to be three episodes of redd building. From the period 1993-2000, the general trend was one of increasing abundance.

However, during the period of 2001-2003, we observed only 1 bull trout redd each year (Figure 10).

A second episode of redd building began after 2003 and a third appears to have begun in 2006. These trends are somewhat similar to other spawning tributaries between Libby Dam and Kootenai Falls. Extreme low water since 1998 and two extreme spill events from Libby (2002 and 2006) have periodically influenced redd building. The low water events dramatically increase water temperatures to as high as 76 degrees F in July and August (MFWP unpublished data) in mainstem Fisher River that is the corridor to reach West Fisher Creek. The effect could be to discourage or delay migration to the traditional spawning grounds during those years. Additionally, high spill events in 2002, 2004 and 2006 from Libby dam have likely reduced the number of adults available to spawn in this particular tributary.




Figure 10. Bull trout redd counts for the West Fisher Creek, 1993-2008.

Keeler Creek

Bull trout that spawn in Keeler Creek (including the North, South and West Forks) are an adfluvial stock that migrates downstream out of Bull Lake into Lake Creek, then up Keeler Creek. This downstream spawning migration is somewhat unique when compared to other bull trout populations (Montana Bull Trout Scientific Group 1996).

We observed a total of 13 and 102 bull trout redds in Keeler Creek and associated tributaries in 2001 and 2002, respectively (Table 16). A beaver dam built in lower Keeler Creek during late summer/early fall 2001 impeded upstream bull trout migration. The dam was removed, but a fall freshet increased stream flow substantially and prevented accurate counts. Therefore, the 13 redds observed in 2001 is undoubtedly an underestimate of the true number of redds in Keeler Creek in 2001.

We observed a total of 84 and 62 bull trout redds in Keeler Creek and associated tributaries in 2007 and 2008, respectively. Prior to those counts, bull trout redd counts in Keeler Creek exhibited a positive trend since monitoring began. Since 2005, there appears to be a negative trend that is substantially different from a stable population (Figure 11). The 2007 and 2008 observations represent 10% and 33% decreases relative to the period of record mean. We will continue to monitor redds and potential barriers for this stream.




Figure 11. Bull trout redd counts and trend analysis for Keeler Creek, a tributary to Lake Creek, 1996-2008.

Callahan Creek

The Callahan Creek redd counts were completed on North Callahan Creek from just above the confluence with South Callahan Creek upstream to water fall in Idaho and in South Callahan Creek from the confluence with North Callahan upstream approximately 3 miles. Redd counts for bull trout have quite variable in this drainage (Figure 12). There was a 20+ flood event during November of 2006 that may affect population densities in 2007 and future redd counts. Idaho Fish and Game monitored the redds for this stream from 2003-2007 we did redd counts for this stream in 2008 and will continue in the future.




Figure 12. Bull trout redd counts for Callahan Creek, 2003 - 2008.

KOOTENAI RIVER ADULT BULL TROUT POPULATION ESTIMATE

Methods

We collected adult bull trout using nighttime electrofishing by jet boat to perform a mark-recapture population estimate of bull trout in the Kootenai River from Libby Dam (River mile [RM] 221.7) downstream to the confluence of the Fisher River (RM 218.2). We operated two boat crews during each sampling event. Each boat contained a driver and two netters. Our electrofishing unit on each boat consisted of a Coffelt model Mark 22 electrofishing unit operating with an electrical output ranging from 200-350 volts at 5-8 amps powered by a 5,000 watt gasoline powered generator.

In order to thoroughly electrofish the entire 3.5 miles of Kootenai River, we divided the sample area into 2 sections, and conducted electrofishing on each section on a single evening. Section 1 was from Libby Dam downstream to the Alexander Creek confluence (RM 220.5), and was 1.2 miles long. Section 2 was from the Alexander Creek confluence downstream to the Fisher River Confluence, and was 2.3 miles long.

We recorded the total time (minutes) electrical current was generated in the water as a measure of effort. We measured total length (mm), weighed (g), examined all fish for marks, collected scale samples, and released all bull trout captured near their capture location. All bull trout were marked with individually numbered 134 (ISO) KHz passive integrated transponder (PIT) tags and an adipose fin clip was removed to evaluate PIT tag retention. PIT tags were inserted with an 8-gauge hypodermic needle into the muscle behind the dorsal fin.

We estimated bull trout abundance using a mark-recapture population estimation technique which assumes the population of bull trout is “closed”, suggesting no births, deaths or migrations occurred during sampling periods (Ricker 1975). Additional assumptions were that marked and unmarked fish have equal mortality rates, marked fish were randomly distributed throughout the study area, marks were not lost, and all marked fish captured were recognized and counted (Lagler 1956). We used a computer software program called Mark/Recapture (version 7.0) that uses a log-likelihood estimator to estimate the absolute abundance of adult bull trout within the study reach. We estimated the total population present within the study area after each marking episode, beginning with the second episode.

Findings

We conducted population estimates for bull trout on the Kootenai River between Libby Dam and Fisher River beginning April 2004. Each marking episode consisted of two nights of capturing fish that ranged between 7 to 9 days between mark and recapture episodes. We calculated population estimates and associated 95% confidence intervals. The mean population estimate for all samples periods over the period ranged from 176 in 2006 to 976 in 2005 (Table 17). We then standardized each population estimate and 95% confidence interval into fish per mile (Table 17).

Table 17. The sampling dates for adult bull trout marked, recaptured, and the estimated total population and fish per mile in the Kootenai River from Libby Dam downstream to the Fisher River confluence, 2004 - 2008.

Dates	Number Marked	Number Captured	Number Recaptured	Total Population Estimate (95 % CI)	Fish per Mile (95 % CI)
April, 2004	109	103	13	920 (698 - 1142)	262 (199 – 326)
April, 2005	See Dunnigan et al. 2006			976 (553 - 1399)	279 (158 – 400)
April, 2006	19	52	5	176 (73 – 279)	50 (21 – 80)
April/May, 2007	37	52	4	347 (124 – 571)	99 (35 – 163)
April , 2008	73	40	7	381 (158 – 605)	109 (43 – 175)

We believe that these substantial swings in population estimates may be a function of the reservoir population. The thought is that cohorts of bull trout, entrained through Libby Dam, reside below the dam and take advantage of entrained kokanee. Through time these groups of fish age and fade out of the population. We don't have enough information to determine the mechanism of replacement for this population though it is undoubtedly caused by a combination of the population of bull trout in the reservoir (especially at the forebay), the fluvial population below the dam and dam operations.

Dunnigan et al. (2009) investigated the genetic structure of bull trout in the Kootenai Basin. They found significant differences exist between populations that reside in all of the known spawning tributaries in the Montana and British Columbia portions of the Kootenai drainage. They determined that the proportion of bull trout residing below Libby Dam between 2004 and 2007 that were entrained ranged from 49.1% to 62.7%. They also found that most of the entrained bull trout originated from the Wigwam River.

The length frequency distribution for bull trout captured in the Libby Dam section of the Kootenai River was fairly consisted among years. The exception was that we captured a higher percentage of juveniles during 2007 and 2008 estimates (Figure 13).




Figure 13. Length frequency distribution of bull trout captured below Libby Dam, 2004 through 2008.

The mean total length for bull trout captured during 2008 was 603 mm (range = 237 – 900mm). The mean length of bull trout in 2004, 2005, 2006, and 2007 captured below Libby Dam was 649, 677, 692, and 655 mm, respectively. Dunnigan et al. (2009) found that the mean length of bull trout captured in 2008 was significantly less than all previous years ($p < 0.05$). These were the only comparisons between years that differed.

We recaptured 14 bull trout during our sampling period April 20 and 21, 2005. All the recaptured fish during this sampling episode were originally captured and marked approximately one year prior (± 1 to 12 days). The recaptured bull trout grew an average of 86.4 mm (range 42 – 179 mm), and gained an average of 2094 g (range 1,421 – 3,550 g).

We recaptured 13 bull trout during our sampling period April 11-19, 2006 that we had previously marked in 2004 and 2005 below Libby Dam. The recaptured bull trout grew an average of 113 mm (range 10 – 201 mm) and gained an average of 1802 g (range 224 – 2908 g).

We recaptured 10 bull trout during our sampling period April 30 – May 8, 2007 that were previously marked in 2004, 2005 or 2006 below Libby Dam ranging between 313 to 1,117 days prior. The recaptured bull trout grew an average of 118.3 mm (range 87 – 161 mm), and gained an average of 1,935.4 g (range 1119 - 3926). We also captured two bull trout that we observed in two of the previous three years; one of these fish was captured in 2004 and 2005, and the other fish was captured in 2004, 2006 and 2007.

We recaptured 16 bull trout during our sampling period April 9 –17, 2008 that were previously marked in 2004, 2005, 2006, or 2007 below Libby Dam. Time between captures ranged from 313 to 1,117 days. The recaptured bull trout grew an average of 96.3 mm and gained an average of 1,649.9 g. Two fish exhibited weight loss since the first capture. One of these fish lost 592 g 722 days after being initially tagged (0.82g/day), and the other lost 128 g in 344 days (3.07 g/day); all other recaptured fish gained length and weight between captures (Dunnigan et al. 2009).

LAKE KOOCANUSA GILLNET MONITORING

Methods

Gillnets have been used by MFWP since 1975 to assess annual trends in fish populations and species composition. These yearly sampling series were accomplished using criteria established by Huston et al. (1984).

Netting methods remained similar to those reported in Chisholm et al. (1989). Netting effort was reduced from 128 ganged (coupled) nets in 1975, to 56 in 1988, and 14 ganged floating and 28 single sinking nets in 1991. Netting effort occurred in the spring and fall, rather than the year round effort prior to 1988. Because of their importance to bull trout either as prey or competitors, kokanee salmon (*Oncorhynchus nerka*) and Kamloops rainbow trout (*Oncorhynchus mykiss gairdneri*) were included in this assessment. Kamloops rainbow trout were distinguished from wild rainbow trout by eroded fins (pectoral, dorsal and caudal); these fish are held in the hatchery until release into the reservoir at age 1+.

The year was stratified into two gillnetting seasons based on reservoir operation and surface water temperature criteria:

- 1) Spring (April - June): The reservoir was being refilled, surface water temperatures increased to 9 - 13°C.
- 2) Fall (September - October): Drafting of the reservoir began, surface water temperature decreased to 13 - 17°C.

Seasonal and annual changes in fish abundance within the near-shore zone were assessed using floating and sinking horizontal gillnets. These nets were 38.1 m long and 1.8 m deep and consisted of five equal panels of 19-, 25-, 32-, 38-, and 51-mm mesh.

Fourteen to twenty-eight floating (ganged) and one or two single, sinking nets were set in the fall in the Tenmile, Rexford and Canada portions of the reservoir. Spring netting series consisted of 20 to 111 (standardized to 28 in 1991) sinking nets and an occasional floating net set only in the Rexford area. Spring floating and fall sinking net data are not included in this report due to a lack of standardization in net placement. Nets were set perpendicular from the shoreline in the afternoon and were retrieved before noon the following day. All fish were removed from the nets and identified, followed by collection of length, weight, sex and maturity data. When large gamefish (Kamloops rainbow, cutthroat, bull trout or burbot) were captured alive, only a length was recorded and the fish were released.

Findings

Initially, we netted areas that included the entire reservoir through all seasons. Over time, seasonal netting was reduced to spring and fall series (Chisholm, et al 1989, Dalbey et al 1997). However, our fall gill netting series typically capture few bull trout. The primary reasons are that sampling dates purposely coincided with the period in which adults were in spawning tributaries, and that bull trout are not traditionally captured in floating gillnets. We netted one area (Koocanusa bridge to Montana/B.C. border) from 1988 to 2008. Table 18 summarizes long-term bull trout mean catch per net in Koocanusa from spring sinking nets.

Table 18. Spring sinking gill net summary of bull trout catch per net in Lake Koocanusa 1975 - 2008.

Year	Date	Reservoir Elevation	Mean Catch Per Net
1975	6/9		1.4
1976	5/1	2373	1.9
1978	5/15	2367	2.2
1980	5/5	2389	0.8
1981	5/5	2378	1.3
1982	5/25	2363	1.5
1984	6/12	2412	1.8
1985	6/6	2415	1.3
1986	5/8	2379	1.9
1987	5/5	2390	1.2
1988	5/12	2344	2.0
1989	5/1	2355	1.2
1990	5/10	2358	1.2
1991	5/16	2330	0.5
1992	5/5	2333	2.3
1993	5/17	2352	1.2
1994	5/16	2405	3.0
1995	5/8	2386	2.3
1996	5/12	2365	3.5
1997	5/12	2350	3.1
1998	5/11	2418	2.5
1999	5/17	2352	3.6
2000	5/14	2371	6.7
2001	5/15	2393	5.4
2002	5/13	2384	4.9
2003	5/13	2417	5.4
2004	5/11	2419	6.4
2005	5/10	2425	6.1
2006	5/10	2423	4.4
2007	5/21	2408	4.5
2008	5/13	2397	5.4

Spring gill net catch of bull trout during the period 1975-1989 appeared to exist at an equilibrium with a slope (0.0091) that was not significantly different than zero ($r^2 = 0.011$; $p = 0.751$). Bull trout catch per net in Libby Reservoir significantly increased between 1990 and 2004. The trend appears to have stabilized since about 2005 (Figure 14). This coincides with the opening of bull trout harvest initiated in 2004 for Koocanusa (Hensler and Benson 2008). We also found that Grave Creek redd counts and bull trout catch per net in Koocanusa have similar trends (Figure 15). Both indices are useful for determining management direction for the harvest of bull trout from Koocanusa.




Figure 14. Average catch per net of bull trout in spring gill nets at the Rexford site on Libby Reservoir 1975-2008.




Figure 17. Bull trout redd count and number per net trends for Grave Creek and Lake Koocanusa, respectively 1993 – 2008.

LITERATURE CITED

- Alderdice, D. F. , W. P. Wickett, and J. R. Brett. 1958. Some effects of temporary exposure to low dissolved oxygen levels on Pacific salmon eggs. *Journal of the Fisheries Research Board of Canada* 15: 229-250.
- Arden, W., P. DeHaan and J. Dunnigan. 2007. Genetic Analysis of Bull Trout in the Kootenai River Basin. Montana, Fish, Wildlife & Parks. Libby, MT.
- Allan, J. H. 1980. Life history notes on the Dolly Varden char (*Salvelinus malma*) in the Upper Clearwater River, Alberta. Manuscript report. Alberta Energy and Natural Resources, Fish and Wildlife Division, Red Deer, Alberta.
- Baxter, C. V. 1997. Geomorphology, land-use, and groundwater-surface water interaction: a multi-scale, hierarchical analysis of the distribution and abundance of bull trout (*Salvelinus confluentus*) spawning. Master's thesis. University of Montana, Missoula, Montana.
- Baxter, J.S. and G.G. Oliver 1997. Bull trout metapopulation structure in the upper Kootenay River: fish and habitat summary data. Prepared for Ministry of Environment, Lands and Parks by Interior Reforestation Co. Ltd. Cranbrook, B.C. 117 pp. Plus Appendices.
- Baxter, J. S. , and W. T. Westover. 1999. Wigwam River bull trout - habitat conservation trust fund progress report (1998). Fisheries Project Report KO 54, viii + 21 p. + 1 appendix.
- Bjornn, T. C. 1969. Embryo survival and emergence studies, Job No. 5, Federal Aid in Fish and Wildlife Restoration. Job Completion Report, Project F-49-R-7. Idaho Fish and Game Department, Boise.
- Bjornn, T. C. , and D. W. Reiser. 1991. Habitat requirements of salmonids in streams. *American Fisheries Society Special Publication* 19:83-138.
- Brown, L. G. 1992. Draft management guide for the bull trout, *Salvelinus confluentus* (Suckley), on the Wenatchee National Forest. Washington Department of Wildlife. Wenatchee, Washington.
- Campana, S. E. 1990. How reliable are growth back-calculations based on otoliths? *Canadian Journal of Fisheries and Aquatic Science* 47, 2219-2227.
- Carlander, K. D. 1981. Caution on the use of the regression method of back-calculating lengths from scale measurements. *Fisheries*, Volume 6, Number 1, p 2-4.
- Chapman, D. W. 1988. Critical review of variables used to define effects of fines in redds of large salmonids. *Transactions of the American Fisheries Society*. 117: 1-21.

- Chisholm, I. and P. Hamlin. 1987. 1985 Libby Reservoir Angler Census: May 13 – October 31, 1985. Interim Report. Project No. 84-52.
- Chisholm, I. M.E. Hensler, B. Hansen, D. Skaar. 1989. Quantification of Libby Reservoir Levels Needed to Maintain or Enhance Reservoir Fisheries-Summary Report. Montana Department of Fish, Wildlife and Parks. Project Number 83-467. Libby, Montana.
- Cross, D. G. and B. Stott. 1975. The effect of electric fishing on the subsequent capture of fish. *Journal of Fish Biology*. Volume 7: 349-357.
- Crouse, M. R. , C. A. Callahan, K. W. Malueg, and S. E. Dominguez. 1981. Effects of fine sediments on growth of juvenile coho salmon in laboratory streams. *Transactions of the American Fisheries Society* 110:281-286.
- Dalbey, S., J. DeShazer, L. Garrow, G. Hoffman, and T. Ostrowski. 1997. Quantification of Libby Reservoir Levels Needed to Maintain or Enhance Reservoir Fisheries: Methods and data Summary, 1988 – 1996. Montana Department of Fish, Wildlife and Parks. Project Number 83-467. Libby, Montana.
- Deleray, M., L. Knotek, S. Rumsey, T. Weaver. 1999. Flathead Lake and River System Fisheries Status Report. Montana Fish, Wildlife & Parks Dingell-Johnson report No. F-78-R-1 through 5, Element 1 Project 1 and 2, SBAS Project No. 3131. Kalispell, MT.
- Dunnigan, J.L., J. DeShazer, L. Garrow, B. Marotz, T. Ostrowski, and C. Sinclair. 2006. Mitigation for the construction and operation of Libby Dam: Libby Mitigation Program, 2005 annual progress report. Bonneville Power Administration. Portland, Oregon. Project Number 199500400.
- Dunnigan, J., J. DeShazer, L. Garrow, T. Ostrowski M. Benner, R. Leary, J. Tohtz, and D. Neeley. 2009. Libby Mitigation Program, 2007 Annual Progress Report: Mitigation for The Construction and Operation of Libby Dam. BPA Project Number 199500400
- Everest, F. H. 1973. Ecology and management of summer steelhead in the Rogue River. Oregon State Game Commission, Fishery Research Report 7, Corvallis, Oregon.
- Everest, F. H. , and five others. 1987. Fine sediment and salmonid production: a paradox. Pages 98-142 *in* E. O. Salo and T. W. Cundy, editors. *Streamside management: forestry and fishery interactions*. University of Washington, Institute of Forest Resources Contribution 57, Seattle, Washington.
- Flathead Basin Commission. 1991. Flathead basin forest practices water quality and fisheries cooperative program. Final report. Flathead Basin Commission, Kalispell, Montana.

- Flathead Basin Commission. 1993. 1991-1992 Biennial Report, Kalispell, Montana, USA.
- Fraley, J. J. , D. Read, and P. J. Graham. 1981. Flathead River fisheries study. Montana Department of Fish, Wildlife and Parks. Kalispell, Montana.
- Fraley, J. J. , and B. B. Shepard. 1989. Life history, ecology and population status of bull trout (*Salvelinus confluentus*) in the Flathead Lake and River system, Montana. Northwest Science 63:133-143.
- Gooch, B. 1967. An evaluation of the two catch method of population estimation. Montana Department of Fish, Wildlife & Parks, Helena, Montana.
- Heimer, J. T. 1965. A supplemental Dolly Varden spawning area. Master's thesis, University of Idaho, Moscow, Idaho. Cited in: Goetz 1989. Biology of the bull trout *Salvelinus confluentus*: a literature review. U. S. Forest Service, Willamette National Forest, Eugene, Oregon.
- Hensler, M, and N. Benson. 2007. Angler survey of experimental recreational bull trout fishery for Lake Koocanusa, Montana 2006-2007. Montana Fish, Wildlife & Parks, Helena, MT
- Hensler, M. and N. Benson. 2008. Angler Survey of Experimental Recreational Bull Trout Fishery In Lake Koocanusa, Montana 2007 -2008. Montana Fish, Wildlife & Parks. Kalispell, MT.
- Junge, C. O. and J. Libovarsky. 1965. Effects of size selectivity on population estimates based on successive removals with electrofishing gear. Zoological Listing 14:171-178.
- Kitano, S. , K. Maekawa, S. Nakano, and K. D. Fausch. 1994. Spawning behavior of bull trout in the upper Flathead Drainage, Montana, with special reference to hybridization with brook trout. Transactions of the American Fisheries Society 123:988-992.
- Knudson, K. 1994. Water quality status report: Kootenay (Kootenai) River Basin British Columbia, Montana and Idaho. Kootenai River Network. Libby, MT. and Ecological Resource Consulting, Helena, MT. 57 pp.
- Koski, K. V. 1966. The survival of coho salmon (*Oncorhynchus kisutch*) from egg deposition to emergence in three Oregon coastal streams. Master's thesis. Oregon State University, Corvallis, Oregon.
- Koski, K. V. 1975. The survival and fitness of two stocks of chum salmon (*Oncorhynchus keta*) from egg deposition to emergence in a controlled stream environment at Big Beef Creek. Doctoral dissertation. University of Washington. Seattle, Washington.
- Lagler, K.F. 1956. Freshwater fishery biology. Wm. C. Brown Company, Dubuque, Iowa.

- Leathe, S. A. , and M. D. Enk. 1985. Cumulative effects of microhydro development on the fisheries of the Swan River drainage, Montana. Volume 1. Summary report prepared for the Bonneville Power Administration, Contracts DE-A179-82BP36717 and DE-A179-83BP39802, Project 92-19.
- Leider, S. A. , M. W. Chilcote, and J. J. Loch. 1986. Movement and survival of presmolt steelhead in a tributary and the main stem of a Washington river. *North American Journal of Fisheries Management* 6:526-531.
- Lelek, A. 1965. A field experiment on the receptivity of chub, *Leuciscus cephalus* (L.) To the repeated influence of pulsating direct current. *Zoological Listing* 15:69-78.
- Liknes, G. A. and P. J. Graham. 1988. Westslope cutthroat trout in Montana: life history, status and management. *American Fisheries Society Symposium* 4:53-60.
- Mahon, R. 1980. Accuracy of catch-effort methods for estimating fish density and biomass in streams. *Biology of Fish* 5(4):343-360.
- McNeil, W. J. and W. H. Ahnell. 1964. Success of pink salmon spawning relative to size of spawning bed materials. U. S. Fish and Wildlife Service, Special Scientific Report 169. Washington, DC
- McPhail, J. D, and C. B. Murray. 1979. The early life history and ecology of Dolly Varden (*Salvelinus malma*) in the upper Arrow Lakes. Department of Zoology and Institute of Animal Resources, University of British Columbia, Vancouver, British Columbia.
- Megahan, W. , W. S. Platts, and B. Kulesza. 1980. Riverbed improves over time: South Fork Salmon. In: Symposium on watershed management. American Society of Civil Engineers, New York. 1:381-395.
- MFWP (Montana Fish, Wildlife & Parks). 1998. Montana statewide angling pressure 1997. Montana Fish, Wildlife & Parks, Helena, Montana.
- Montana Bull Trout Scientific Group (MBTSG). 1996a. Lower Kootenai River drainage bull trout status report (below Kootenai Falls). Prepared for The Montana Bull Trout Restoration Group. Helena, MT.
- Montana Bull Trout Scientific Group (MBTSG). 1996b. Upper Kootenai River drainage Bull Trout status report (including Lake Koocanusa, Upstream of Libby Dam). Prepared for The Montana Bull Trout Restoration Group. Helena, MT.
- Needham, P. R. , and T. M. Vaughan. 1952. Spawning of the Dolly Varden, *Salvelinus malma*,

- in Twin Creek, Idaho. *Copeia*, 1952, Number 3, pp. 197-199.
- Oliver, G.G. and R.S. Cope. 1999. Fisheries implications – Wigam River forest development plan review. Correspondence to Lloyd A. Havens, Crestbrook Forest Industries Ltd., Cranbrook, B.C. February 9, 1999.
- Oliver, G. 1979. A final report on the present fisheries of the Wigwam River with emphasis on the migratory life history and spawning behavior of Dolly Varden charr *Salvalinus malma* (Walbaum). Fisheries investigations in tributaries of the Canadian portion of Libby Reservoir. British Columbia Fish and Wildlife Branch, Victoria, BC.
- Paragamian, V. L. 1994. Kootenai River fisheries inventory: stock status of burbot and rainbow trout and fisheries inventory. Idaho Department of Fish and Game, Bonneville Power Administration, Project 88-65. Boise, ID.
- Phillips, R. W. , R. L. Lantz, E. W. Claire, and J. R. Moring. 1975. Some effects of gravel mixtures on emergence of coho salmon and steelhead trout fry. *Transactions of the American Fisheries Society* 104:461-466.
- Platts, W. S. and R. L. Nelson. 1988. Fluctuations in trout populations and their implications for land-use evaluation. *North American Journal of Fisheries Management* 8:333-345.
- Pratt, K. L. 1984. Habitat selection and species interactions of juvenile westslope cutthroat trout (*Salmo clarki lewisi*) and bull trout (*Salvelinus confluentus*) in the upper Flathead River Basin. Master's thesis. University of Idaho, Moscow, Idaho.
- Pratt, K. L. 1985. Pend Oreille trout and char life history study. Idaho Department of Fish and Game. Boise, Idaho.
- Ratliff, D. E. 1992. Bull trout investigations in the Metolius River-Lake Billy Chinook System. Pages 37-44 in P. J. Howell and D. V. Buchanan, editors. *Proceedings of the Gearhart Mountain bull trout workshop*. Oregon Chapter of the American Fisheries Society, Corvallis, Oregon.
- Reiser, D. W. , and T. A. Wesche. 1979. In situ freezing as a cause of mortality in brown trout eggs. *Progressive Fish-Culturist* 41: 58-60.
- Ricker, W. E. 1975. Computation and interpretation of biological statistics of fish populations. *Fisheries Research Board of Canada Bulletin* 191.
- Rieman, B.E. and D.L. Myers. 1997. Use of redd counts to detect trends in bull trout (*Salvelinus confluentus*) populations. *Conservation Biology* 11(4):1015-1018.
- Seber, G. A. F. and E. D. LeCren. 1967. Estimating population parameters from large catches

- relative to the population. *Journal of Animal Ecology* 36:631-643.
- Shepard, B. and P. J. Graham. 1982. Completion report. Monitoring spawning bed material used by bull trout on the Glacier View District, Flathead National Forest. Montana Department of Fish, Wildlife & Parks, Kalispell, Montana. 37 pp.
- Shepard, B. B. and P. J. Graham. 1983. Fish Resource Monitoring Program for the Upper Flathead Basin. Prepared for the Environmental Protection Agency, Contract Number R008224-01-4. Montana Fish, Wildlife & Parks, Kalispell, Montana.
- Shepard, B. B. , K. L. Pratt, and P. J. Graham. 1984. Life histories of westslope cutthroat and bull trout in the upper Flathead River basin, Montana. Prepared for Environmental Protection Agency, Contract No. R008224-01-5. Montana Fish, Wildlife & Parks, Kalispell, Montana.
- Shepard, B. , S. A. Leathe, T. M. Weaver, and M. D. Enk. 1984. Monitoring levels of fine sediment within tributaries to Flathead Lake, and impacts of fine sediment on bull trout recruitment. Unpublished paper presented at the Wild Trout III Symposium. Yellowstone National Park, Wyoming. On file at: Montana Department of Fish, Wildlife and Parks, Kalispell, Montana.
- Shumway, D. L. , C. E. Warren, and P. Doudoroff. 1964. Influence of oxygen concentration and water movement on the growth of steelhead trout and coho salmon embryos. *Transactions of the American Fisheries Society* 93:342-356.
- Silver, S. J. , C. E. Warren, and P. Doudoroff. 1963. Dissolved oxygen requirements of developing steelhead trout and chinook salmon embryos at different water velocities. *Transactions of the American Fisheries Society* 92:327-343.
- Skaar, D., J. DeShazer, L. Garrow, T. Ostrowski and B. Thornberg. 1996. Quantification of Libby Reservoir levels needed to enhance reservoir fisheries. Investigations of fish entrainment through Libby dam, 1990-1994. Final Report. Montana Department of Fish, Wildlife and Parks - Region 1. Prepared for Bonneville Power Administration. Project Number 83-467.
- Tappel, P. D. and T. C. Bjornn. 1983. A new method of relating size of spawning gravel to salmonid embryo survival. *North American Journal of Fisheries Management* 3:123-135.
- Tepper, H. 2002. Wigwam River McNeil sampling program: 1998 – 2002 Summary report. Prepared for Bonneville Power Administration. BPA project Number 2000-004-00. Bonneville Power Administration, Fish and Wildlife Program, Portland, Oregon, USA.
- Van Deventer, J.S., and W.S. Platts. 1983. Sampling and estimating fish populations from

- Streams: North American Wildlife and Natural Resources Conference. 48:349-354.
- Weaver, T. M. , and J. J. Fraley. 1991. Fisheries habitat and fish populations. Flathead basin forest practices, water quality and fisheries cooperative program. Flathead Basin Commission, Kalispell, Montana.
- Weaver, T. M. and J. J. Fraley. 1993. A method to measure emergence success of westslope cutthroat trout fry from varying substrate compositions in a natural stream channel. North American Journal Fisheries Management 13:817-822.
- Weaver, T. M. , and R. G. White. 1985. Coal Creek fisheries monitoring study No. III. Quarterly progress report. U. S. Forest Service, Montana State Cooperative Fisheries Research Unit, Bozeman, Montana.
- Westover, W. 2002. Kootenai Kokanee Enumeration (2002). Ministry of Water, Land and Air Protection. Cranbrook, British Columbia.
- White, G. C. , D. R. Anderson, K. P. Burnham, D. L. Otis. 1982. Capture-recapture and removal methods for sampling closed populations. United States Department of Energy. Contract W-7465-ENG-36. Project Report LA-8787-NERP. Los Alamos National Laboratory, Los Alamos, New Mexico.
- Wickett, W. P. 1958. Review of certain environmental factors affecting the production of pink and chum salmon. Journal of the Fisheries Research Board of Canada. 15:1103-1126.
- Ziller, J. S. 1992. Distribution and relative abundance of bull trout in the Sprague River subbasin, Oregon. Pages 18-29 In: P. J. Howell and D. V. Buchanan, editors. Proceedings of the Gearhart Mountain bull trout workshop. Oregon Chapter of the American Fisheries Society, Corvallis, Oregon.
- Zippin, C. 1958. The removal method of population estimation. Journal of Wildlife Management 22(1):82-90.