

FURBEARER PROGRAM

STATEWIDE HARVEST
& MANAGEMENT REPORT
2012-2013

MONTANA



FURBEARER PROGRAM

2012-13 STATEWIDE

HARVEST AND MANAGEMENT REPORT

PERIOD COVERED: July 2012 – June 2013

PREPARED BY: Brian Giddings
State Furbearer Coordinator
Montana Fish, Wildlife & Parks
P.O. Box 200701
Helena, MT 59620-0701

DATE: June 2014

TABLE OF CONTENTS

List of Tables	4
List of Figures	6
Program Goals	8
Statewide Objectives	8
Management Strategies	8
Harvest and Management Activities	9
Statewide Harvest and Management Results	10
Montana Furbearer Program Related Bibliography in Chronological Order	58

LIST OF TABLES

Table

1	Montana trapping license sales, 2012-2013.....	14
2	Montana furbearer, predator and nongame species harvest summary, 2012-2013...	14
3	Montana furbearer, predator and nongame species 10-year harvest summary, 2003-04 to 2012-13.....	15
4	Average pelt price reported by species, 1990-91 to 2012-13.....	16
5	Beaver harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	17
6	Otter harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	19
7	Muskrat harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	23
8	Mink harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	25
9	Marten harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	27
10	Fisher harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	31
11	Wolverine harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	36
12	Bobcat harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.....	40
13	Weasel harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.....	44
14	Skunk harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.....	46
15	Coyote harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.....	48
16	Fox harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.....	50
17	Raccoon harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.....	52
18	Badger harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.....	54

LIST OF FIGURES

Figure

1	Map of Montana delineating furbearer trapping districts and counties.....	13
2	Montana trapper license sales trend, 1975-76 to 2012-13.....	13
3	Statewide beaver harvest by trapping district, 1994-95 to 2012-13.....	18
4	Statewide trend in beaver harvest from CPUE, 1995-96 to 2012-13.....	18
5	Statewide otter harvest by trapping district, 1994-95 to 2012-13.....	20
6	Statewide trend in otter harvest from CPUE, 1995-96 to 2012-13.....	20
7	Otter population parameter of juveniles per adult female ratio, 2003-04 to 2012-13.....	21
8	Otter population parameter of age structure, 2003-04 to 2012-13.....	21
9	Otter population parameter of sex ratio, 2003-04 to 2012-13.....	22
10	Otter population parameter of median ages, 2003-04 to 2012-13.....	22
11	Statewide muskrat harvest by trapping district, 1994-95 to 2012-13.....	24
12	Statewide trend in muskrat harvest from CPUE, 1995-96 to 2012-13.....	24
13	Statewide mink harvest by trapping district, 1994-95 to 2012-13.....	26
14	Statewide trend in mink harvest from CPUE, 1995-96 to 2012-13.....	26
15	Statewide marten harvest by trapping district, 1994-95 to 2012-13.....	28
16	Statewide trend in marten harvest from CPUE, 1995-96 to 2012-13.....	28
17	Marten population parameter of juveniles per adult female ratio, 1996-97 to 2005-06.....	29
18	Marten population parameter of age structure, 1996-97 to 2005-06.....	29
19	Marten population parameter of sex ratio, 2003-04 to 2012-13.....	30
20	Marten population parameter of median ages, 1996-97 to 2005-06.....	30
21	Statewide fisher harvest by trapping district, 1994-95 to 2012-13.....	32

22	Statewide trend in fisher harvest from CPUE, 1995-96 to 2012-13.....	32
23	Fisher population parameter of juvenile per adult female ratio, 2003-04 to 2012-13.....	33
24	Fisher population parameter of age structure, 2003-04 to 2012-13.....	33
25	Fisher population parameter of sex ratio, 2003-04 to 2012-13.....	34
26	Fisher population parameter of median ages, 2003-04 to 2012-13.....	34
27	Statewide wolverine harvest by trapping district, 1994-95 to 2012-13.....	36
28	Statewide trend in wolverine harvest from CPUE, 1995-96 to 2012-13.....	37
29	Wolverine population parameter of juvenile per adult female ratio, 2002-03 to 2011-12.....	37
30	Wolverine population parameter of age structure, 2002-03 to 2011-12.....	38
31	Wolverine population parameter of sex ratio 2002-03 to 2011-12.....	38
32	Wolverine population parameter of median ages, 2002-03 to 2011-12.....	39
33	Statewide bobcat harvest by trapping district, 1994-95 to 2012-13.....	41
34	Statewide trend in bobcat harvest from CPUE, 1995-96 to 2012-13.....	41
35	Bobcat population parameter of juvenile per adult female ratio, 2003-04 to 2012-13.....	42
36	Bobcat population parameter of age structure, 2003-04 to 2012-13.....	42
37	Bobcat population parameter of sex ratio, 2004-03 to 2012-13.....	43
38	Bobcat population parameter of median ages, 2003-04 to 2012-13.....	43
39	Statewide weasel harvest by trapping district, 1995-96 to 2012-13.....	45
40	Statewide trend in weasel harvest from CPUE, 1995-96 to 2012-13.....	45
41	Statewide skunk harvest by trapping district, 1994-95 to 2012-13.....	47
42	Statewide trend in skunk harvest from CPUE, 1995-96 to 2012-13.....	47
43	Statewide coyote harvest by trapping district, 1994-95 to 2012-13.....	49
44	Statewide trend in coyote harvest from CPUE, 1995-96 to 2012-13.....	49

45	Statewide fox harvest by trapping district, 1994-95 to 2012-13.....	51
46	Statewide trend in fox harvest from CPUE, 1995-96 to 2012-13.....	51
47	Statewide raccoon harvest by trapping district, 1994-95 to 2012-13.....	53
48	Statewide trend in raccoon harvest from CPUE, 1995-96 to 2012-13.....	53
49	Statewide badger harvest by trapping district, 1994-95 to 2012-13.....	55
50	Statewide trend in badger harvest from CPUE, 1995-96 to 2012-13.....	55
51	Statewide harvest trend comparison of species group from CPUE, 1995-96 to 2012-13.....	56
52	Statewide harvest trend comparison of species group from CPUE, 1995-96 to 2012-13	56
53	Statewide harvest trend comparison of species group from CPUE, 1995-96 to 2012-13	57
54	Statewide harvest trend comparison of species group from CPUE, 1995-96 to 2012-13.....	57

Program Goals

- 1) Maintain well-distributed and healthy furbearer populations and associated habitats.
- 2) Provide ecological, recreational, cultural, educational, economic, and scientific benefits of the state's furbearers through sound resource management.
- 3) Address the social impacts of furbearers on human health, private property, and agricultural values.

Statewide Objectives

- 1) Monitor population trends and the distribution of each furbearer species.
- 2) Maintain Montana's viable populations of each species by promoting the conservation and enhancement of furbearer habitats.
- 3) Address the interest by resident publics for consumptive and non-consumptive uses of the state's furbearer resource.
- 4) Optimize recreational harvest opportunities through a sustained use management approach under regulatory protections.
- 5) Minimize animal damage and/or nuisance wildlife problems utilizing Department policies and management practices.
- 6) Promote trapping practices that minimize the take of non-target species and maximizes the humane harvest of furbearers.
- 7) Develop a public understanding and acceptance for the basis of the consumptive use of furbearers.

Management Strategies

- 1) Identify and associate species distribution and population trends with delineated habitats.
- 2) Investigate species population trends through species/habitat surveys, species occurrence reports, harvest data, and research information.
- 3) Utilize regulatory mechanisms to provide trapper/hunter participation, harvest data, and biological information.
- 4) Include furbearer species in land management decisions.

Harvest and Management Activities

- 1) Population information and harvest data are collected by county and/or trapping district and reported by trapping district and statewide in this report. This method is intended to more closely describe the association between species diversity, distribution, and abundance with identified ecosystems and to use reconcilable legal units in the state. Furbearer species with harvest seasons are beaver, otter, muskrat, mink, marten, fisher, wolverine, bobcat, and swift fox. Furbearers with a closed season are lynx, and are not included in this report. Weasel, skunk and coyote are state classified predators while red fox, raccoon, and badger are nongame species of which limited harvest data is collected so they are included in this report.
- 2) The annual harvests of otter, marten, fisher, wolverine and bobcat are monitored through a statewide reporting, pelt tagging and harvest registration system. Registration is initiated under 24-hour mandatory reporting through an automated telephone call-in system referred to as the Mandatory Reporting Response Entry (MRRE) system. All pelt tag sealing and completion of species harvest registration forms, which are generated in MRRE, are conducted by FWP personnel. Marten, fisher, wolverine and swift fox pelts are tagged under the authority of the state, while otter and bobcat are tagged under oversight of the U.S. Fish & Wildlife Service to meet federal CITES pelt export requirements.
- 3) Harvest data on the three remaining furbearers (beaver, muskrat, mink) and six fur-producing animals (weasel, skunk, coyote, fox, raccoon, badger) was collected through a trapper harvest survey questionnaire. In addition, the same harvest data is collected on the five tagged/registered furbearers through the same survey questionnaire to specifically measure trapper effort and catch rates. Trapper effort will be used in developing long-term species population trend indices. The trapping and fur harvest survey was mailed to all resident and nonresident license holders. No reminder was sent to non-respondents. Expanded estimates of furbearer trapping, hunting, and harvest activities were made from the returned sample. The survey requests information on the estimated number of species harvested by county and trapping district, harvest method, and harvest effort. Summary harvest statistics and calculated catch rates were generated by a software package through FWP's Research & Technical Services Unit.
- 4) Mandatory carcass collections are required for fisher and wolverine, and skulls must be surrendered from harvested otter, bobcat and swift fox. Marten skulls have not been required to be turned in since the 2008-09 season, but were collected in prior years. All carcasses and skulls are forwarded to FWP's Wildlife Laboratory in Bozeman for biological analysis to determine specimen age, sex, body condition, food habits, reproductive history, and to collect tissue samples for potential genetic analysis.
- 5) A Montana fur dealer survey conducted by the state furbearer coordinator has been replaced by checking the North American Fur Auction (NAFA) website after the winter and spring sales in order to obtain average pelt values for each fur-producing species. An increasing number of Montana trappers are shipping directly to NAFA. This information can be used to calculate economic fur value of each species as a predictor of harvest pressure (i.e. higher prices = greater harvest pressure).

- 6) Annual winter furbearer snow track surveys had been conducted by regional wildlife biologists following a standardized survey protocol and track identification methods in Trapping Districts 1-4 (NW and SW mountainous forest habitats). However, track surveys were discontinued after the 2011-12 winter until a further evaluation is conducted.
- 7) Biologists in trapping districts 4-7 are in the process of developing lagomorph prey indices through the use of headlight surveys. The numbers of lagomorphs are counted on established routes three times each survey period. This index to prey availability is utilized to predict bobcat population fluctuations by anticipating changes in annual rabbit production (March surveys) or recruitment levels (September surveys).
- 8) Department furbearer occurrence/distribution report forms are distributed and collected annually. Reports are completed only by Department personnel from verified reports or personal observations. Accumulated reports provide species occurrence and location data to assist in delineating statewide and trapping district distribution of selected furbearer species (otter, fisher, wolverine, lynx and swift fox).
- 9) Furbearer research is an ongoing statewide activity that is utilized to address management related issues on a species-specific basis when funding is available.

Statewide Harvest and Management Results

Harvest and management results were analyzed by county and trapping district and reported as a statewide summary. The seven legally defined trapping districts (TDs) and 56 Montana counties are shown in Fig. 1.

License Sales

The 6,299 trapping license sold during the 2012-13 season was a 19% increase from the previous year of 5,053 licenses and 37% above the 10-year average (Fig. 2). License purchases at the seven regional offices and the Helena headquarters are somewhat mixed each year. However, in 2012-13 all regional offices and Helena had increased sales from the previous year (Table 1). The ability to purchase licenses online, but probably more so was the first gray wolf trapping season in 2012-13 with a requirement that wolf trapper's purchase a general trapper license. Again, a general upward trend in statewide license sales is apparently continuing through 2012-13 from the lowest sales at any time in 1990-91 when 1,736 licenses were sold.

Annual Harvest Summary

Montana's furbearer harvest for the 2012-13 season is presented in Table 2. A 10-year harvest summary for years that species harvest data is available is presented in Table 3. These figures represent the known legal harvest of registered furbearer species and an estimated harvest of the remaining six species based on the trapper harvest survey. Detailed harvest statistic estimates by species, trapping district and county are available in the Trapping and Fur Harvest Reports (K. Podruzny, pers. comm.). During the most recent year available, trapper survey questionnaires were returned from approximately 35% of the 6,299 people who purchased a trapper's license during the 2012-13 furbearer season. The total number of animals reported being taken during the 2012-13 season increased by 8% over the 2011-12 season (Table 3). This increase may be the

result of mild weather conditions, generally stable populations of most furbearing animal species in various portions of the state, and higher than average pelt prices for most species.

Pelt Prices

Pelt prices continued a general increase for most species during the 2012-13 season, with several species demonstrating large increases in value, particularly beaver, otter, muskrat, mink, fisher, bobcat, and red fox (Table 4). The most significant increases in harvest numbers were reported for beaver, muskrat, marten, bobcat, coyote, fox and raccoon with decreases in the harvest of weasel and skunk, despite slightly higher prices from the previous year.

Species Harvest Summary

Statewide species harvest trends by trapping districts and statewide are presented in the Species Harvest Summary section (pages 17 to 60). The statewide harvest of most species was generally stable to increasing with larger increases in the mink, marten and coyote harvests during 2012-13. These changes are variable, however, among the seven trapping districts. Harvest numbers may correspond to species abundance within each habitat type, although other variables, such as trapper effort and catch rate, may be more useful indicators to correlate harvest data with population trends. Under this assumption, there are specific implications for habitat and species management opportunities.

Population Monitoring

Results calculated from the trapper harvest survey that reports trapper effort for all species, including the known registered species harvest, provided Catch Per Unit Effort (CPUE = # animals harvested/1,000 trap days) which is used to help monitor population trends (Species Harvest Summary (pages 17 to 60). Using estimated catch rates (trap days/catch) from the annual harvest survey continues to be evaluated as a population monitoring parameter. Metrics such as these will be examined further to determine how well they may reflect species population trend. Graphs of the CPUE for species groups, to compare trends among similar species, are presented on pages 56 and 57 in the Species Harvest Summary section.

Results of carcass collections from fisher and wolverine, and skull turn-in from otter and bobcat are shown under each of these species sections in the Species Harvest Summary (pages 17 to 60). The most important aspect of these collections is to extract a tooth for age determination. The graphs illustrate analysis of the biological parameters reported, which are juvenile/adult female ratios, age structure, sex ratios, and median ages of the harvest sample which should represent population parameters. Not all years or most recent years may be available for age data, as processing adult teeth can take 1-2 years for results. Marten skull collections were discontinued beginning with the 2008-09 season, partially because stable population parameters have remained unchanged. Fisher and wolverine sample sizes are extremely small, so they do not necessarily represent a population trend. Any additional species information from wildlife laboratory analysis will be reported as it becomes available in future reports.

The number of FWP occurrence/distribution reports received showed a consistent trend during the past several years with the majority of reports collected for wolverine and swift fox. Recent reports will to be entered in a locational referenced database, similar to the furbearer harvest database that

provides species distribution data. The number of counties in the state for reported swift fox observations continues to increase.

Furbearer Research

Research related to furbearers that were supported and/or permitted by FWP or the state furbearer program during the report period includes a lynx project in northwestern Montana and an otter genetics connectivity investigation was initiated during the report period. The USFS lynx project is ongoing. No results are available from the otter genetics analysis. Wolverine work conducted from 2002-2010 has been completed and continues to be published in peer review journals R. Inman, pers. comm.). A bibliography of most furbearer program related research to date is listed on pages 61 to 67 in this report.




Figure 1. Map of Montana delineating furbearer regulation trapping districts and counties.




Figure 2. Montana trapper license sales trend, 1975-76 to 2012-13.

Table 1. Montana trapping license sales, 2012-2013.

License Type	Kalispell	Missoula	Bozeman	Great Falls	Billings	Glasgow	Miles City	Helena	Statewide
General	850	677	1,039	625	746	183	490	1,502	6,112
Youth	9	13	14	4	5	1	1	1	47
Landowner	6	8	10	23	31	10	32	8	128
Nonresident	0	0	6	0	0	1	4	1	12
Total	865 (+27%)	698 (+33%)	1,069 (+15%)	652 (+12%)	782 (+15%)	195 (+17%)	527 (+13%)	1,512 (+22%)	6,299 (+20%)

Table 2. Montana furbearer, predator and nongame species harvest summary, 2012-2013.

Trapping District	1	2	3	4	5	6	7	Total*
Beaver	653	1,104	1,931	1,711	835	532	309	7,085
Otter	26	16	32	8	3	0	0	85 (9)
Muskrat	4,352	8,247	8,548	3,089	1,437	1,727	330	27,731
Mink	183	255	486	153	102	298	13	1,491
Marten	293	656	459	8	27	--	--	1,443
Fisher	3	3	--	--	--	--	--	6 (0)
Wolverine	0	0	0	0	0	--	--	0
Bobcat	280	196	273	281	299	53	557	1,939
Swift Fox	-	-	-	-	-	21	-	21
Weasel	172	70	24	13	3	8	11	301
Skunk	115	102	140	244	626	201	282	1,711
Coyote	655	894	1,335	3,919	2,334	5,093	5,899	20,131
Red Fox	223	596	290	309	440	207	771	2,837
Raccoon	83	274	964	588	2,557	763	1,327	6,557
Badger	29	0	91	497	72	293	309	1,292
Total	7,067	12,413	14,573	10,820	8,735	9,106	9,808	72,630

*Figure may include animals harvested in unknown trapping district and () indicates incidental harvest.

Table 3. Montana furbearer, predator and nongame species 10-year harvest summary, 2003-2004 to 2012-2013.

Year	2003-04	2004-05	2005-06	2006-07	2007-08	2008-09	2009-10	2010-11	2011-12	2012-13
Beaver	9,361		8,918	7,421	7,219	7,124	5,795	5,445	6,833	7,086
Otter	80	88	93	78	67	60	64	58	68	88
Muskrat	11,915		21,270	17,014	10,042	10,699	12,754	18,494	27,236	27,731
Mink	808		1,306	1,348	1,018	655	584	760	872	1,491
Marten	1,062	1,248	952	856	1,141	844	711	932	1,083	1,721
Fisher	8	7	9	7	6	7	6	7	7	7
Wolverine	10	11	11	9	9	4	3	4	2	0
Bobcat	1,783	2,114	2,201	2,228	2,389	2,428	1,738	1,644	1,975	1,638
Weasel	321		243	503	310	175	120	488	342	301
Skunk	2,996		2,325	1,933	2,599	1,845	2,717	3,975	1,735	1,711
Coyote	12,286		9,412	10,886	9,723	6,969	9,048	8,489	16,398	20,131
Fox	2,056		2,473	3,164	1,862	1,696	1,471	1,418	2,469	2,837
Raccoon	5,936		4,540	4,368	4,506	4,052	4,099	2,201	6,409	6,557
Badger	1,788		1,166	1,330	871	643	450	609	1,474	1,292
Total	50,411		54,939	51,145	41,762	37,201	39,110	45,531	66,919	72,591

Table 4. Average pelt price reported by species, 1990-91 to 2012-13.

Year	Beaver	Otter	Muskral	Mink	Marten	Fisher	Wolverine	Bobcat	Coyote	Red Fox	Raccoon	Weasel	Skunk	Badger
1990-91	9.52	25.15	0.73	13.84	25.47	35.00	140.00	90.98	13.01	8.45	4.32	0.27	4.05	5.29
1991-92	11.81	17.50	1.30	20.50	25.58	40.00	130.00	87.00	23.95	22.50	8.28	2.25	4.25	7.65
1992-93	8.02	39.76	1.18	10.21	17.24	35.00	135.00	85.37	22.18	11.17	10.68	3.50	4.52	8.38
1993-94	12.35	33.30	1.54	10.02	21.74	32.74	147.80	90.43	15.78	10.68	10.10	2.00	3.01	6.82
1994-95	14.95	30.00	1.67	9.31	15.00			81.75	20.61	15.33	9.30	2.66	3.40	11.87
1995-96	16.13	35.95	2.82	9.16	19.17		200.00	75.42	19.46	18.58	10.97	1.75	6.15	10.00
1996-97	23.59	30.98	3.83	14.48	25.01			124.05	24.68	17.74	15.26	1.83	3.86	11.19
1997-98	21.18	20.00	1.94	9.54	17.25			95.25	17.15	12.72	14.67	1.00	2.85	11.73
1998-99								85.50						
1999-00					19.33			98.67	22.06					
2000-01	15.98	59.17	1.71	8.37	19.95	28.62	212.94	106.05	18.93	16.24	10.02	1.50	3.73	15.98
2001-02	12.40	47.93	2.07	10.05	18.70	25.12	225.00	135.25	23.70	22.65	19.30	2.00	5.00	18.50
2002-03	14.00	75.00	2.10	10.50	19.50	25.00	225.00	203.00	30.70	24.00	11.00	3.00	7.00	21.50
2003-04	14.50	90.00	2.15	11.00	20.50	28.10	275.00	280.25	28.50	20.00	11.50	3.00	5.50	23.00
2004-05	15.25	94.00	2.25	11.50	19.50	28.25	275.00	325.00	30.70	21.50	11.00	3.00	7.00	23.50
2005-06	20.50	100.00	3.50	15.00	45.50	35.00	300.00	345.00	38.50	25.00	11.50	3.00	6.50	27.50
2006-07	23.49	80.00	3.20	12.88	61.57	74.31	217.85	257.33	43.36	20.84	22.05	4.96	4.04	27.57
2007-08	24.80	40.91	3.23	15.22	77.29	87.51	280.35	449.45	37.90	22.49	33.22	5.69	5.27	42.60
2008-09	25.21	30.85	2.55	11.53	37.58	42.83	254.67	281.35	30.70	21.59	17.86	4.02	2.32	24.80
2009-10	16.74	51.10	4.23	17.39	47.76	50.08	211.42	346.54	35.29	22.34	18.02	4.07	2.34	72.56
2010-11	16.57	57.63	6.66	17.48	61.98	47.58	253.15	411.84	73.16	24.37	18.50	3.13	2.11	24.12
2011-12	38.22	102.29	10.19	23.14	55.94	74.99	319.67	426.31	77.30	57.49	19.45	3.16	7.30	38.61
2012-13	30.91	112.58	\$11.51	20.05	84.70	145.30	235.74	589.08	93.98	65.78	27.56	3.13	4.26	25.45

SPECIES HARVEST AND MANAGEMENT SUMMARY

BEAVER

The statewide beaver harvest has been relatively stable over the past several years, but has increased in 2011-12 and again in 2012-13 over the previous several years, but still remains at a lower level from the most recent peak harvests in the late 1990s (Fig.3). The estimated 2012-13 harvest of 7,085 beaver is 16% below the 10-year average harvest level, despite above average reported pelt price (Table 5).

Examining the trend in CPUE it appears harvest effort decreased during the 2012-13 season after an increase the previous year, indicating that less beaver are being taken per unit of effort (Fig. 4). Population monitoring activities for beaver are based completely on reported trapper survey data, with the CPUE considered to indicate relative population trend, which could be considered as relatively stable, with a recent decreasing to stable trend since 2007-08. A comparison of CPUE for beaver with the other semi-aquatic species is shown in Fig. 51.

Generally higher pelt prices will lead to more trapper effort, as reflected in the CPUE during 2011-12 for beaver and an increase in harvest numbers, particularly in certain portions of the state, however that trend declined in 2012-13. Habitat conditions may also be influencing beaver numbers by expanding water areas and riparian tributaries as a result of good spring moisture conditions, which could lead to less damage control complaints, at least in some regional areas of the state, particularly in western and central Montana.

Table 5. Beaver harvest, pelt price, and quota level if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	823	1173	2795	2637	2164	1847	248	11699	14.95	
1995-96	679	846	1854	2118	2127	711	285	8620	16.13	
1996-97	626	1118	2961	5681	3453	1590	1122	16550	23.59	
1997-98	698	1194	4460	3005	2227	972	959	13515	21.18	
1998-99	510	1045	3243	3942	1900	718	276	11634		
1999-00	908	1298	2821	2966	1961	2265	587	12805		
2000-01	399	1095	2623	1756	2528	407	247	9056	15.98	
2001-02	499	1394	3242	2953	1266	1273	460	11156	12.41	
2002-03	685	1071	2296	2040	1201	777	399	8475	14.01	
2003-04	424	1485	2336	2074	2175	477	389	9361	14.51	
2004-05									15.25	
2005-06	767	628	2852	1970	856	1626	219	8918	20.51	
2006-07	479	944	2067	1450	1509	661	310	7421	23.49	
2007-08	209	812	1409	788	698	994	313	7219	24.81	
2008-09	415	513	2015	1199	618	460	107	7124	25.21	
2009-10	466	836	1021	1034	437	233	295	5795	16.74	
2010-11	315	825	963	1356	709	16	267	5445	16.57	
2011-12	357	1225	1805	1931	567	696	252	6833	38.22	
2012-13	653	1104	1931	1711	835	532	309	7085	30.91	




Figure 3. Statewide beaver harvest by trapping district, 1994-95 to 2012-13.




Figure 4. Statewide trend in beaver harvest from CPUE, 1995-96 to 2012-13.

OTTER

Otter are one of five furbearers that are required to be reported, registered and pelt tagged so that the actual number of harvested animals is known. The 2012-13 harvest of 85 otters is about 13% above the 10-year average of 74 otters (Table 6). The otter harvest has always been managed through a trapper limit and since the 2002-03 season also under trapping district (TD) quotas. Up through the 2001-02 season a one otter per trapper limit was in place, which also changed in 2002-03 to a two otter limit per trapper under the regulated quota in each of the seven TD's. The two otter limit and quota changes were made in response to healthy populations, to reduce surrendered incidental take in beaver sets, and more interest by trappers as pelt prices were increasing at that time. Quotas are now used as a harvest management tool to maintain well distributed and healthy otter populations, while trapper limits provide more opportunity and flexibility to harvest otter by the trapping community. The total quota for the state has increased from 84 in 2002-03 to 95 in 2007-08, at which level it has remained (Table 6). The statewide otter harvest increased with pelt prices until a peak price and corresponding harvest occurred during the 2005-06 season. Harvest has since declined through the 2011-12 season, but increased by 20% in 2012-13 along with higher pelt price over the last two years. However, the long-term harvest level and proportion of the harvest by TD has remained relatively stable (Fig. 5).

The statewide trend in otter harvest CPUE has been relatively stable, however an increase occurred in 2012-13, indicating less effort per otter harvest which may suggest an increasing otter population in the state (Fig. 6). A comparison of otter CPUE with the other semi- aquatic species is presented in Fig. 51. Population monitoring for otter consists of the collection and analysis of biological data from the harvest sample through mandatory carcass turn-in from trappers through the 2011-12 season. Starting with the 2012-13 season, only otter skulls are being collected. Although not all data is available for some years, trends in population parameters of juveniles per adult female, age structure, sex ratios, and median ages shown in Fig. 7 to 10 indicate a strong juvenile segment and show a relatively stable population on a statewide basis.

Table 6. Otter harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	23	7	23	4	5	0	0	62	30.01	
1995-96	17	8	22	6	7	0	1	61	35.95	
1996-97	17	8	27	7	6	0	0	65	30.98	
1997-98	15	8	41	13	7	0	0	84	20.01	
1998-99	17	4	34	9	3	0	0	67		
1999-00	18	9	26	8	3	0	0	64		
2000-01	13	15	18	1	1	0	0	48	59.17	
2001-02	28	23	39	5	1	0	0	96	47.93	
2002-03	21	13	35	8	4	0	1	83	75.01	84
2003-04	19	18	33	8	2	0	0	80	90.01	84
2004-05	25	19	32	8	3	0	1	88	94.01	92
2005-06	20	22	36	8	5	0	2	93	100.01	93
2006-07	21	17	29	6	5	0	0	78	80.01	93
2007-08	24	14	17	5	2	0	1	67	40.91	95
2008-09	21	14	22	0	3	0	0	60	30.85	95
2009-10	21	20	17	8	2	0	0	68	51.10	95
2010-11	20	14	18	4	3	0	0	59	57.63	95
2011-12	22	19	21	3	3	0	0	68	102.29	95
2012-13	26	16	32	8	3	0	0	85	112.58	95




Figure 5. Statewide otter harvest by trapping district, 1994-95 to 2012-13.




Figure 6. Statewide trend in otter harvest from CPUE, 1995-96 to 2012-13.




Figure 7. Otter population parameter of juveniles per adult female ratio, 2003-04 to 2012-13.




Figure 8. Otter population parameter of age structure, 2003-04 to 2012-13.




Figure 9. Otter population parameter of sex ratios, 2003-04 to 2012-13




Figure 10. Otter population parameter of median ages, 2003-04 to 2012-13.

MUSKRAT

The statewide muskrat harvest has continued to increase over the past several years. The 2012-13 season produced the highest estimated harvest of 27,731 animals over the last 19 years and remained at a similar level to the 2011-12 harvest of 27,236 muskrat (Table 7). This increasing harvest may have been influenced by higher than average pelt prices of \$11.51 in 2012-13 compared to the previous 10-year average pelt price of \$4.00. In addition, the estimated 2012-13 muskrat harvest was 46% above the 10-year average harvest of 15,194 animals (Fig. 11).

Population monitoring activities for muskrat are based completely on trapper harvest survey data, with CPUE from the harvest survey considered to be an indicator of relative population trend, which could be considered as stable to increasing, with a recent decline during 2011-12 and 2012-13 (Fig. 12). Examining the trend in CPUE it appears catch rates were generally increasing until the past two years, indicating that in 2012-13 less muskrat are being taken per unit of effort, possibly indicative of the above average harvest numbers and trapping pressure (Fig.12). A comparison of CPUE for muskrat with the other semi-aquatic species is shown in Fig. 51.

Table 7. Muskrat harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	1393	4905	4394	2152	925	404	83	14256	1.67	
1995-96	716	4177	3271	1791	1276	181	39	11727	2.82	
1996-97	2980	3992	2732	3712	1799	772	134	16121	3.83	
1997-98	2552	3887	5043	3519	1499	2122	205	18826	1.94	
1998-99	2270	2240	3495	2609	709	811	111	12243		
1999-00	1643	3156	2651	3049	794	763	1191	13247		
2000-01	897	6170	2905	536	2844	129	361	13842	1.71	
2001-02	556	5681	3409	599	596	132	43	11070	2.07	
2002-03	1427	3915	4571	952	308	156	119	11448	2.11	
2003-04	869	3923	5625	864	318	45	270	11915	2.15	
2004-05									2.25	
2005-06	1561	4902	9862	2203	888	1217	637	21270	3.51	
2006-07	1850	4821	5210	2418	1868	728	117	17014	3.21	
2007-08	510	806	1188	761	522	442	146	10042	3.23	
2008-09	485	1131	2037	801	567	0	0	10699	2.55	
2009-10	852	2564	3054	1953	546	404	48	12754	4.23	
2010-11	949	1977	4452	4684	628	822	51	18494	6.66	
2011-12	1740	6304	11057	3180	705	3452	799	27236	10.19	
2012-13	4352	8247	8548	3089	1437	1727	330	27731	11.51	




Figure 11. Statewide muskrat harvest by trapping district, 1994-95 to 2012-13.




Figure 12. Statewide trend in muskrat harvest from CPUE, 1995-96 to 2012-13.

MINK

The statewide mink harvest has continued to increase over the past several years since the lowest harvest during 2008-09 in recent years. The 2012-13 estimated harvest of 1,491 mink is the highest since 2000-01 (Table 8). Mink harvest is considered to be somewhat correlated to interest in muskrat trapping, which appears to be the case with the 2012-13 peak in muskrat harvest numbers. The estimated 2012-13 mink harvest was 37% above the 10-year harvest average (Fig. 13), along with above average pelt prices. The average value of mink pelts was the second highest level in at least 19 years (Table 8). Mink harvest levels are likely tied to landownership patterns and trapper access to streams and wetlands, and where muskrat can be found on public land wetlands.

Population monitoring activities for muskrat are based completely on trapper harvest survey data, with the CPUE considered to be an indicator of relative population trend, which could be considered as stable, despite the above average estimated harvest during the 2012-13 season. When examining the trend in CPUE for mink, it appears harvest effort has generally remained stable, with some changes in harvest effort that may be related over time to interest in muskrat trapping, indicating that mink are being harvested at about a similar scale per unit of effort, at least through the 2012-13 season (Fig. 14). A comparison of CPUE for mink with the other semi-aquatic species is shown in Fig. 51.

Table 8. Mink harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	187	215	274	234	97	121	17	1145	9.31	
1995-96	140	290	111	126	128	87	34	919	9.16	
1996-97	252	134	339	488	126	280	20	1638	14.48	
1997-98	220	174	381	248	289	133	49	1493	9.54	
1998-99	285	162	309	171	120	27	3	1078		
1999-00	218	183	428	325	38	476	41	1709		
2000-01	95	198	1038	103	57	15	30	1536	8.37	
2001-02	111	300	307	89	61	43	32	959	10.05	
2002-03	92	229	564	94	13	38	40	1071	10.51	
2003-04	43	290	331	71	45	3	25	808	11.01	
2004-05									2.25	
2005-06	62	151	563	92	92	340	6	1306	15.01	
2006-07	94	269	678	129	158	18	3	1348	12.88	
2007-08	122	101	80	51	86	182	98	1018	15.22	
2008-09	62	85	127	20	28	0	0	655	11.53	
2009-10	40	62	118	171	35	13	5	584	17.39	
2010-11	57	154	175	129	27	3	21	760	17.48	
2011-12	53	190	415	102	58	23	29	872	23.14	
2012-13	183	255	486	153	102	298	13	1491	20.05	




Figure 13. Statewide mink harvest by trapping district, 1994-95 to 2012-13.




Figure 14. Statewide trend in mink harvest from CPUE, 1995-96 to 2012-13.

MARTEN

Marten are one of the five furbearers that are required to be reported, registered and pelt tagged so that the actual number of harvested animals is known. The statewide marten harvest continued to increase, with an increasing harvest trend during the past several years including 2012-13 (Fig. 15). The 2012-13 harvest of 1,443 marten was 32% above the 10-year average and at the highest harvest level over the past 19 years. The higher harvest in 2012-13 may correspond to a similar increase in pelt prices (Table 9). Examining the trend in CPUE it appears harvest effort has remained relatively stable on a statewide basis, indicating more marten are being taken with consistent per unit of effort (Fig.16). Also, the distribution of the marten harvest is apparently shifting somewhat back to TD 1 in northwestern Montana as well as TD 2 in the west central part of the state while southwestern Montana remains similar to previous years (Fig. 15). Primary marten habitat is located almost exclusively on public forest lands.

Population monitoring for marten has consisted of analyzing harvest data and using the collection and analysis of biological data from the harvest sample through mandatory skull turn-in from trappers. However, marten skull collections were discontinued beginning with the 2008-09 season because of the difficulty in reconciling individual skulls to male/female categories for age data. The statewide trend in marten using CPUE appears to be a stable trend (Fig.16) and a comparison of marten CPUE with the other terrestrial species is presented in Fig 52. The last year of aging marten was 2005-006 so this prior 10-year period is presented in Figures 17-19. From the available data collected prior to 2006-07 season, the previous long-term trend in population parameters continued to demonstrate a stable trend that showed an average of three juveniles per adult female (Fig. 17), a positive age structure bias to juveniles (Fig. 18), a stable sex ratio (Fig. 19), with a slight decrease in median age of adults and median age of total harvest at one and a half, indicating a strong proportion of juveniles in the population (Fig. 20). Again, these parameters indicate a relatively stable population over time that is maintaining a healthy population status on a statewide basis.

Table 9. Marten harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	868	315	131	4	5			1323	15.01	
1995-96	433	167	202	0	0			802	19.17	
1996-97	513	172	143	0	2			830	25.01	
1997-98	403	291	192	9	5			900	17.25	
1998-99	473	172	61	3	7			716		
1999-00	313	183	149	1	7			653	19.33	
2000-01	560	326	174	1	3			1064	19.95	
2001-02	359	220	266	0	0			845	18.71	
2002-03	419	241	390	3	0			1053	19.51	
2003-04	459	339	259	2	3			1062	20.51	
2004-05	290	374	560	3	21			1248	19.51	
2005-06	280	265	370	1	36			952	45.51	
2006-07	143	268	418	2	25			856	61.57	
2007-08	245	446	441	0	9			1141	77.29	
2008-09	170	366	282	0	26			844	37.58	
2009-10	99	402	192	0	18			711	47.76	
2010-11	184	363	333	0	52			932	61.98	
2011-12	353	420	308	2	1			1083	55.94	
2012-13	293	656	459	8	27			1443	84.70	




Figure 15. Statewide marten harvest by trapping district, 1994-95 to 2012-13.




Figure 16. Statewide trend in marten harvest from CPUE, 1995-96 to 2012-13.




Figure 17. Marten population parameter of juveniles per adult female ratio, 1996-97 to 2005-06.




Figure 18. Marten population parameter of age structure, 1996-97 to 2005-06.




Figure 18. Marten population parameter of sex ratios, 2003-04 to 2012-13.




Figure 19. Marten population parameter of median ages, 1996-97 to 2005-06.

FISHER

The fisher harvest has been managed through a trapper limit of one and quotas in trapping districts (TD) 1 and 2. Over time, conservative adjustments have been made to quota levels that were based on harvest rates, population parameters and snow track survey data. These previous changes have provided a sustainable trapper harvest that is conservatively matched with maintenance of the current fisher population size and distribution relative to available habitats. A predictive habitat model indicates that moderate to high suitability fisher habitats comprise approximately 6,504 mi² in west central and northwestern Montana, with TD 2 having over 50% more high suitability habitat than TD 1. A female sub-quota is also in place of 2 females to add an additional measure of protection for the reproductive segment of the population, to further insure harvest has no influence on statewide population status. Given fisher distribution relative to habitat availability, fisher habitat capacity appears to be correlated with similar levels of occupancy that is not impacted by a history of highly managed harvest. Fisher are one of the five furbearers that are required to be reported, registered and pelt tagged so the actual number of harvested animals is known. The fisher harvest continues to remain very stable under the current quota system (Fig. 21), despite a much higher pelt price (Table 10). The 2012-13 harvest of 6 fishers was near the 10-year average annual harvest level.

Population monitoring of fisher consists of analyzing harvest data and using the collection and analysis of biological data from the harvested animals through mandatory carcass turn-in from trappers. The trend in fisher harvest effort using CPUE has been a stable trend with an increase during 2012-13 (Fig.22). A comparison of fisher CPUE with the other terrestrial species is presented in Fig 52. Harvested fishers provide an extremely small sample size, so population parameters do not allow a lot of interpretation. However, the small amount of data that is available appears to show that the population trend from these parameters is about two juveniles per adult female (Fig. 23), a mixed age structure with a good representation of juveniles in most years (Fig. 24), a low female sex ratio in most years (Fig. 25), with a higher than expected median age of adults and expected median age of the total harvest (Fig. 26).

Table 10. Fisher harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	3	5						8		10
1995-96	0	2						2		10
1996-97	2	4						6		7
1997-98	1	6						7		7
1998-99	2	6						8		7
1999-00	0	5						5		7
2000-01	0	7						7	28.62	7
2001-02	2	5						7	25.12	7
2002-03	2	5						7	25.01	7
2003-04	2	6						8	28.11	7
2004-05	0	7						7	28.25	7
2005-06	3	6						9	35.01	7
2006-07	2	5						7	74.31	7
2007-08	1	5						6	87.51	7
2008-09	1	6						7	42.83	7
2009-10	1	5						6	50.08	7
2010-11	1	7						8	47.58	7
2011-12	2	5						7	74.99	7
2012-13	3	3						6	145.30	7




Figure 21. Statewide fisher harvest by trapping district, 1994-95 to 2012-13.




Figure 22. Statewide trend in fisher harvest from CPUE, 1995-96 to 2012-13.




Figure 23. Fisher population parameters of juveniles per adult female ratio, 2003-04 to 2012-13.




Figure 24. Fisher population parameter of age structure, 2003-04 to 2012-13.




Figure 25. Fisher population parameter of sex ratios, 2003-04 to 2012-13.




Figure 26. Fisher population parameter of median ages, 2003-04 to 2012-13.

WOLVERINE

Since wolverines were first classified as a state furbearer in the late 1970s, harvest was regulated by a one wolverine per trapper limit. Wolverine harvest was considered to be recovered in Montana from a low point in the 1930s and now occupied the western third of the state. A study in the mid-1970s found that wolverines were at relatively high densities in the South Fork of the Flathead River drainage. Since then, statewide harvest during a 30-year period was considered stable and somewhat self-regulating with an average of 10.5 wolverines harvested annually (range 2 - 22 per year) during the period. However, recent research on the species has provided new information regarding wolverine ecology, better defined wolverine habitat, examined genetic relationships, survival, and landscape connectivity. FWP's furbearer program provided funds and logistical support to these studies. Research results were used to develop a habitat model for Montana with corresponding population numbers and estimated sustainable harvest rates which were considered sustainable at a more regulated level. Therefore, FWP changed trapping regulations to reflect emerging information and additional data from wolverine research that suggested conservative quota levels were appropriate. Over the past several years quotas were adjusted to associate harvest levels with the three largest intact ecosystems in the state (Northern Continental Divide, Bitterroot-Idaho and Greater Yellowstone) and to recognize the lower population sizes in insular mountain ranges in the central portion of Montana. Further analysis tied to genetic make-up of the state's wolverine population, the issue of maintaining population connectivity, and recognizing the core population areas of the three major ecosystems led to additional regulation changes. These most recent adjustments included delineating four wolverine management units (WMUs) with the three major ecosystems having reduced quotas for a statewide total of 5 animals and a central Montana WMU with a quota of zero to promote population connectivity between the three major ecosystems in the state where harvest is allowed. Managing the WMU/quota system has maintained biologically sound harvest opportunity for trappers, that does not jeopardize conservation of the species. However, with the pending decision by the USFWS to list wolverine under the ESA, the quota was reduced to zero during the 2012-13 season, so no harvest occurred.

Wolverines are one of five furbearers that are required to be reported, registered and pelt tagged so that the actual number of harvested animals is known (Table 11). The statewide wolverine harvest continues to remain stable through 2011-12 (Fig. 27). Conservative quotas implemented beginning with the 2008-09 season were reflected in the harvest of two wolverines during the 2011-12 season, which was 75% below the 10-year average harvest. This restrictive quota system has achieved the management goal of redistributing and lowering the wolverine harvest in the state. Harvest of wolverine is considered independent of pelt prices (Table 10) but does provide a unique harvest opportunity for resident trappers. Examining the trend in CPUE through 2011-12, it appears harvest effort has been relatively stable to more recently declining on a statewide basis, indicating that less wolverine are being taken per unit of effort, which would be expected (Fig.28).

Population monitoring for wolverine has consisted of analyzing harvest data and using the collection and analysis of biological data from the harvest sample through mandatory carcass turn-in from trappers. Up through the 2011-12 season, the statewide trend in wolverine using CPUE is a stable to declining trend (Fig.28) and a comparison of wolverine CPUE with the other terrestrial species is presented in Fig 52. The harvest sample of wolverine has provided only a very small sample size, so population parameters are more difficult to interpret. However, the long-term trend in population parameters have showed about .5 juveniles per adult female (Fig. 29), a mixed age structure with a relatively continuous representation of juveniles (Fig. 30), about a 50% female sex ratio (Fig. 31), with a median age of adults higher than the median age of the total harvest (Fig. 32).

Table 11. Wolverine harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	2	1	5	1	0			9		
1995-96	5	2	4	1	0			12	200.01	
1996-97	6	0	3	2	1			12		
1997-98	1	5	6	3	0			15		
1998-99	0	2	2	5	0			9		
1999-00	0	0	3	1	0			4		
2000-01	1	6	4	2	0			14	212.94	
2001-02	1	0	9	0	0			10	225.01	
2002-03	2	2	8	2	1			15	225.01	
2003-04	1	2	3	2	2			10	275.01	
2004-05	3	1	6	1	0			11	275.01	12
2005-06	0	4	4	2	1			11	300.01	12
2006-07	2	0	5	2	0			9	217.85	12
2007-08	2	1	5	1	0			9	280.35	10
2008-09	2	0	0	2	0			4	254.67	5
2009-10	1	1	1	0	0			3	211.42	5
2010-11	0	3	1	0	0			4	253.15	5
2011-12	0	2	0	0	0			2	319.67	5
2012-13	0	0	0	0	0			0	235.74	0




Figure 27. Statewide wolverine harvest by trapping district, 1994-95 to 2012-13.




Figure 28. Statewide trend in wolverine harvest from CPUE, 1995-96 to 2012-13.




Figure 29. Wolverine population parameter of juveniles per adult female ratio, 2002-03 to 2011-12.




Figure 30. Wolverine population parameter of age structure, 2002-03 to 2011-12.




Figure 31. Wolverine population parameter of sex ratios, 2002-03 to 2011-12.




Figure 32. Wolverine population parameter of median ages, 2002-03 to 2011-12.

BOBCAT

Bobcats are one of five furbearers that are required to be reported, registered and pelt tagged so that the actual number of harvested animals is known (Table 12). The bobcat harvest has always been managed through trapping district (TD) quotas with changes in trapper limits or removal of trapper limits in some districts. In the late 1990s trapper limits were increased in response to low trapper interest in bobcats because of relatively low pelt prices (Table 12) and later removed altogether in the eastern districts (TD 4 – 7). Adjustments in bobcat quotas have been used as a management tool to maintain healthy bobcat populations, while providing opportunity and flexibility to harvest bobcat by the trapping community. As bobcat populations in the state have increased over time, along with trapper interest, TD quotas have generally increased proportionately. The statewide total quota has increased from 1,415 in 1994-95 to 2,480 in 2008-09 and 1,970 during the 2012-13 season (Table 12). The bobcat harvest has increased from 1,052 in 1994-95 to 2,428 in 2008-09 and 1,939 in 2012-13 (Fig. 33). Pelt prices have jumped dramatically beginning with the 2003-04 season and continue to remain at a high level through 2012-13 (Table 12). Examining the trend in CPUE it appears harvest effort has been stable to slightly declining, indicating that fewer bobcat are being taken per unit of effort (Fig.34).

The statewide trend in bobcat using CPUE is declining slightly (Fig.34) and a comparison of bobcat CPUE with the other terrestrial species is presented in Fig 52. Population monitoring for bobcat consists of analyzing harvest data and the collection and analysis of biological data from the harvest sample through mandatory skull turn-in from trappers to extract a tooth to determine age. Population parameters show a recent upward trend in juveniles per adult female (Fig. 35), an older age structure (Fig. 36), less than a 50% female sex ratio (Fig. 37), and an increasing trend in the median age of the total harvest (Fig. 38). Again, these parameters indicate a stable to a somewhat declining population trend on a statewide basis.

Table 12. Bobcat harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	148	117	121	145	157	75	289	1052	81.75	1415
1995-96	169	113	105	105	109	12	149	762	75.42	1440
1996-97	166	108	133	174	165	45	250	1041	124.05	1440
1997-98	167	158	139	163	191	40	348	1206	95.25	1490
1998-99	173	159	134	133	197	68	229	1093	85.51	1490
1999-00	199	170	145	184	212	91	410	1411	98.67	1510
2000-01	222	168	128	173	230	86	391	1398	106.05	1630
2001-02	244	178	173	177	267	121	542	1702	135.25	1730
2002-03	201	146	199	193	315	135	597	1786	203.01	1805
2003-04	210	182	229	211	356	88	507	1783	280.25	1880
2004-05	225	172	218	312	424	135	628	2114	325.01	2030
2005-06	230	158	291	287	392	122	721	2201	345.01	2255
2006-07	243	177	294	320	426	91	677	2228	257.33	2255
2007-08	264	182	314	316	489	100	724	2389	449.45	2355
2008-09	258	184	292	298	503	71	822	2428	281.35	2480
2009-10	248	108	203	214	487	42	465	1767	346.54	2275
2010-11	278	113	216	245	406	26	360	1644	411.84	2175
2011-12	259	104	275	311	308	91	627	1975	426.31	1925
2012-13	280	196	273	281	299	53	557	1939	589.08	1970




Figure 33. Statewide bobcat harvest by trapping district, 1994-95 to 2012-13




Figure 34. Statewide trend in bobcat harvest from CPUE, 1995-96 to 2012-13.




Figure 35. Bobcat population parameter of juvenile per adult female ratios, 2003-04 to 2012-13.




Figure 36. Bobcat population parameter of age structure, 2003-04 to 2012-13.




Figure 37. Bobcat population parameter of sex ratios, 2003-04 to 2012-13.




Figure 38. Bobcat population parameter of median ages, 2003-04 to 2012-13.

WEASEL

The statewide weasel harvest continues to remain relatively stable, although generally at a lower level during the past decade within a range of 200 to 500 animals with some years below this level (Table 13). The majority of weasels taken over most years is in northwestern Montana's trapping district (TD) 1 (Fig. 39). The estimated 2012-13 statewide harvest of 301 animals was right at the 10-year average harvest, along with average pelt prices (Table 13). Despite the moderate harvest, average pelt prices offered for 2012-13 were higher than a decade ago.

Examining the trend in CPUE it appears harvest effort has generally increased over the past several years, indicating that more weasels are being taken per unit of effort (Fig.40). Population monitoring activities for weasel are based completely on trapper harvest survey data, with CPUE considered to be an indicator of relative population trend, which could be considered stable, despite the decline in 2012-13. A comparison of CPUE for weasel with the other classified predator species is shown in Fig. 53.

Table 13. Weasel harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	286	222	161	109	5	19	0	802	2.66	
1995-96	264	53	24	0	2	0	0	343	1.75	
1996-97	217	16	154	618	8	4	79	1094	1.83	
1997-98	123	54	153	56	0	0	0	386	1.01	
1998-99	144	48	9	42	3	0	0	246		
1999-00	211	86	24	155	0	0	3	480		
2000-01	87	11	19	42	0	0	8	167	1.51	
2001-02	75	7	14	4	0	0	0	100	2.01	
2002-03	248	124	32	0	0	0	0	405	3.01	
2003-04	88	164	51	13	3	0	3	321	3.01	
2004-05									3.01	
2005-06	118	77	9	27	12	0	0	243	3.01	
2006-07	213	161	79	35	12	0	3	503	4.96	
2007-08	185	45	21	12	3	0	0	310	5.69	
2008-09	45	76	6	0	0	0	0	175	4.02	
2009-10	54	24	8	13	0	3	0	121	4.07	
2010-11	164	84	181	13	3	3	3	488	3.13	
2011-12	199	105	15	6	0	15	3	342	3.16	
2012-13	172	70	24	13	3	8	11	301	3.13	




Figure 39. Statewide weasel harvest by trapping district, 1994-95 to 2012-13.




Figure 40. Statewide trend in weasel harvest from CPUE, 1995-96 to 2012-13.

SKUNK

The statewide skunk harvest continues to remain stable, and within a general range of 1,500 to 3,000 animals with some years below or above this level (Table 14). The majority of skunks taken over most years come from the central and southern portions of Montana in trapping districts (TD) 4 and 5 (Fig. 41). The estimated 2012-13 statewide harvest of 1,711 animals was 26% below the 10-year average harvest, with a slightly lower than average pelt price (Table 14). A lower harvest than the previous several years may be a result of the pelt price offered for 2012-13 of \$4.26 that was lower than average over the last decade.

Examining the trend in CPUE it appears harvest effort again declined during the 2012-13 season, indicating that fewer skunks are being taken per unit of effort (Fig.42). Population monitoring activities for skunk are based completely on trapper harvest survey data, with CPUE considered to be an indicator of relative population trend, which could be considered to be declining trend over previous several years. A comparison of CPUE for skunk with the other classified predator species is shown in Fig. 53.

Table 14. Skunk harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	69	194	336	222	532	579	1287	3219	3.41	
1995-96	75	198	167	89	401	162	619	1784	6.15	
1996-97	142	169	638	260	705	539	929	3382	3.86	
1997-98	102	138	573	394	445	281	749	2682	2.85	
1998-99	84	246	345	342	306	15	228	1567		
1999-00	7	90	238	780	1015	0	632	2762		
2000-01	72	213	445	175	361	163	141	1570	3.73	
2001-02	46	182	578	442	71	150	146	1616	5.01	
2002-03	40	224	421	248	154	100	235	1422	7.01	
2003-04	167	177	616	397	493	937	210	2996	5.51	
2004-05									7.01	
2005-06	195	145	652	492	252	296	293	2325	6.51	
2006-07	99	187	251	503	477	44	371	1933	4.04	
2007-08	27	209	161	442	152	510	471	2599	5.27	
2008-09	48	113	180	361	643	0	299	1845	4.02	
2009-10	107	53	212	1407	447	27	112	2717	2.34	
2010-11	51	294	267	2567	464	48	113	3975	2.11	
2011-12	32	120	292	140	436	117	597	1735	7.30	
2012-13	115	102	140	244	626	201	2	1711	4.26	




Figure 41. Statewide skunk harvest by trapping district, 1994-95 to 2012-13.




Figure 42. Statewide trend in skunk harvest from CPUE, 1995-96 to 2012-13.

COYOTE

The statewide coyote harvest increased dramatically during the 2011-12 and again in the 2012-13 seasons from both trapping and hunting (Table 15). The majority of coyotes taken over most years are in the northcentral and eastern Montana trapping district's (TD) 4, 6 and 7 (Fig. 43). The estimated 2012-13 statewide harvest of 20,131 animals was nearly 50% above the 10-year average harvest, along with the highest average pelt price of \$93.98 (Table 15). This average coyote pelt prices offered for 2012-13 was the highest reported in the past 19 years.

Examining the trend in CPUE it appears harvest effort has generally remained the same, indicating an increasing coyote population and/or that there is an increase in the number of successful trappers and hunters (Fig. 40). Population monitoring activities for coyote are based completely on trapper harvest survey data, with CPUE considered to be an indicator of relative population trend, which could be considered stable. A comparison of CPUE for coyote with the other classified predator species is shown in Fig. 53.

Table 15. Coyote harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

1994-95	284	851	1774	2112	1227	788	3034	10079	20.61	
1995-96	312	728	991	1216	1197	389	624	5495	19.46	
1996-97	189	1193	1594	2953	1445	925	1055	9354	24.68	
1997-98	524	1424	2163	2496	1493	821	1588	10510	17.15	
1998-99	267	874	1387	1486	688	453	904	6059		
1999-00	514	798	1429	3142	1526	1060	2651	11134	22.06	
2000-01	167	593	1483	1836	1563	559	2988	9303	18.93	
2001-02	114	745	2086	2211	774	1783	2004	9726	23.71	
2002-03	175	971	1452	1357	567	3386	2817	10725	30.71	
2003-04	306	1046	2311	3198	1485	1632	2309	12286	28.51	
2004-05									30.71	
2005-06	278	823	1291	1650	569	2431	2346	9412	38.51	
2006-07	433	789	1485	2269	1058	2713	2137	10886	43.36	
2007-08	197	546	1200	1716	451	2286	1946	9723	37.91	
2008-09	387	437	494	1453	494	827	1780	6969	30.71	
2009-10	193	396	544	651	571	153	1112	9048	35.29	
2010-11	485	661	464	1764	1135	677	2203	8489	73.16	
2011-12	292	605	1243	4660	1834	3487	4276	16398	77.3	
2012-13	655	894	1335	3919	2334	5093	5899	20131	93.98	




Figure 43. Statewide coyote harvest by trapping district, 1994-95 to 2012-2013.




Figure 44. Statewide trend in coyote harvest from CPUE, 1995-96 to 2012-13.

RED FOX

The statewide red fox harvest increased during the 2012-13 season following a general increasing harvest since 2009-10 after a declining trend in harvest over the prior 15 years (Table 16). The majority of the red fox taken over most years is across all trapping districts (TD) except TD 1 (Fig. 45). The estimated 2012-13 statewide harvest of 2,837 animals was only 22% above the 10-year average harvest level, despite the highest pelt price in 19 years of \$65.78 (Table 16).

Examining the trend in CPUE it appears harvest effort has generally been stable with a decrease during 2011-12 that remained constant in 2012-13, indicating that fewer fox are being taken per unit of effort (Fig.46). Population monitoring activities for red fox are based completely on trapper harvest survey data, with CPUE considered to be an indicator of relative population trend, which could be considered as declining. A comparison of CPUE for fox with the other unclassified nongame species is shown in Fig. 54.

Table 16. Fox harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	19	284	1133	738	2039	691	1963	6872	15.33	
1995-96	73	280	498	411	1267	181	790	3573	18.58	
1996-97	87	402	898	1795	909	677	996	5764	17.74	
1997-98	54	355	1327	795	898	307	1074	4810	12.72	
1998-99	27	210	321	495	438	129	534	2156		
1999-00	10	414	701	842	483	494	684	3629		
2000-01	19	243	521	608	293	270	240	2201	16.24	
2001-02	7	478	770	735	364	435	285	3074	22.65	
2002-03	8	483	523	380	216	364	577	2552	24.01	
2003-04	23	465	434	523	296	68	248	2056	20.01	
2004-05									21.51	
2005-06	38	358	178	509	145	569	670	2473	25.01	
2006-07	55	380	465	409	441	757	655	3164	20.84	
2007-08	45	164	248	266	227	155	277	1862	22.49	
2008-09	20	234	130	367	265	56	299	1695	21.59	
2009-10	16	195	166	80	335	16	129	1471	22.34	
2010-11	113	377	167	162	232	59	156	1418	24.37	
2011-12	29	541	333	328	450	196	591	2469	57.49	
2012-13	223	596	290	309	440	207	771	2837	65.78	




Figure 45. Statewide red fox harvest by trapping district, 1994-95 to 2012-13.




Figure 46. Statewide trend in red fox harvest from CPUE. 1995-96 to 2012-13.

RACCOON

The statewide raccoon harvest had been declining but has increased in recent years, and jumped substantially during the 2011-12 and 2012-13 seasons relative to previous years (Table 17). The majority of raccoon harvested by trapping or hunting over most years has been in southern Montana's trapping districts (TD) 3, 5 and to a lesser degree TD 7 (Fig. 47). The estimated 2012-13 statewide harvest of 6,557 animals is the third highest in 19 years and was 28% above the 10-year average harvest. This high harvest was accompanied by the highest average pelt price of \$27.56 in 19 years (Table 17).

Examining the trend in CPUE it appears harvest effort has declined in the past couple of years, indicating that fewer raccoon are being taken per unit of effort (Fig.48). Population monitoring activities for raccoon are based completely on trapper harvest survey data, with CPUE considered to be an indicator of relative population trend, which could be considered as declining. However, with a higher raccoon harvest level in 2011-12 and 2012-13 with higher pelt prices may be an indication of increased interest in trapping and/or hunting for raccoons. A comparison of CPUE for raccoon with the other unclassified nongame species is shown in Fig. 54.

Table 17. Raccoon harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	64	220	627	520	1724	272	965	4392	9.31	
1995-96	41	111	205	728	2335	471	795	4687	10.97	
1996-97	220	189	1012	1807	3547	976	1465	9216	15.26	
1997-98	61	338	1146	1422	2363	706	921	6956	14.67	
1998-99	144	198	871	736	1855	129	267	4200		
1999-00	69	200	977	908	1661	394	735	4944		
2000-01	11	205	1057	342	2091	281	399	4387	10.02	
2001-02	29	307	1484	485	1337	289	1273	5203	19.31	
2002-03	62	283	939	410	1160	380	1427	4662	11.01	
2003-04	78	258	1008	371	1869	904	1447	5936	11.51	
2004-05									11.01	
2005-06	121	154	1146	524	1125	500	814	4540	11.51	
2006-07	108	240	889	532	1517	266	816	4368	22.05	
2007-08	60	161	421	555	1277	358	651	4506	33.22	
2008-09	39	99	711	717	1343	70	307	4052	17.86	
2009-10	37	155	268	171	1037	145	137	4099	18.02	
2010-11	75	285	359	372	1294	89	218	3201	18.5	
2011-12	73	322	1141	503	2989	380	1000	6409	19.45	
2012-13	83	274	964	588	2557	763	1327	6557	27.56	




Figure 47. Statewide raccoon harvest by trapping district, 1994-95 to 2012-13.




Figure 48. Statewide trend in raccoon harvest from CPUE, 1995-96 to 2012-13.

BADGER

The statewide badger harvest has been relatively inconsistent over the years with low harvest numbers during several previous years and then a significant increase during 2012-13 and the previous season (Table 18). The majority of the badgers taken over most years are in north central and southeastern Montana's trapping districts (TD) 4 and 7 (Fig. 49). The estimated 2012-13 statewide harvest of 1,292 animals was 22% above the 10-year average harvest, along with a below average pelt price of \$25.45 (Table 18). This higher harvest following several years of lower than average harvests occurred despite generally good pelt prices.

Examining the trend in CPUE it appears harvest effort has been stable to slightly declining in 2012-13, indicating that a relatively stable number of badger are being taken per unit of effort (Fig. 50). Population monitoring activities for badger are based completely on trapper harvest survey data, with CPUE considered to be an indicator of relative population trend, which could be considered as stable. A comparison of CPUE for badger with the other unclassified nongame species is shown in Fig. 54.

Table 18. Badger harvest, pelt price, and harvest quota if applicable, 1994-95 to 2012-13.

Year	TD 1	TD 2	TD 3	TD 4	TD 5	TD 6	TD 7	State	Pelt Price	Quota
1994-95	12	17	114	289	26	135	338	931	11.87	
1995-96	2	2	85	280	29	5	85	491	10.01	
1996-97	4	4	102	1260	24	157	268	1819	11.19	
1997-98	0	5	174	563	38	146	146	1071	11.73	
1998-99	0	3	51	87	9	42	69	261		
1999-00	7	3	166	400	21	41	352	991		
2000-01	8	15	114	209	30	84	38	498	15.98	
2001-02	4	4	160	360	57	82	75	742	18.51	
2002-03	13	24	229	378	27	116	224	1012	21.51	
2003-04	8	20	361	765	336	66	232	1788	23.01	
2004-05									23.51	
2005-06	3	39	187	394	122	113	308	1166	27.51	
2006-07	0	32	269	178	190	324	336	1330	27.57	
2007-08	3	27	72	173	54	95	286	871	42.61	
2008-09	0	6	42	51	25	0	169	643	24.81	
2009-10	5	24	5	27	16	27	45	450	72.56	
2010-11	5	48	40	65	48	51	151	609	24.12	
2011-12	3	12	237	714	12	284	213	1474	38.61	
2012-13	29	0	91	497	72	293	309	1292	25.45	




Figure 49. Statewide badger harvest by trapping district, 1994-95 to 2012-13.




Figure 50. Statewide trend in badger harvest from CPUE, 1995-96 to 2012-13.




Figure 51. Statewide harvest trend comparison of species group from CPUE, 1995-96 to 2012-13.




Figure 52. Statewide harvest trend comparison of species group from CPUE, 1995-96 to 2012-13




Figure 53. Statewide harvest trend comparison of species group from CPUE, 1995-96 to 1012-13.




Figure 54. Statewide harvest trend comparison of species group from CPUE, 1995-96 to 2012-13.

Montana FWP Furbearer Program Related Bibliography in Chronological Order

- Wright, P.L. **1947**. The sexual cycle of the male long-tailed weasel (*Mustela frenata*). J. Mammalogy 28(4):343-352.
- Townsend, J.E. **1952**. A study in beaver ecology in western Montana with special reference to movements. M.S. thesis. Montana State University, Bozeman, MT. 44 p. (J. Mammalogy 34(4):459-479).
- Townsend, J.E. **1954**. Beaver bank accounts. Montana Wildlife. 4(1):16-17.
- Greer, K.R. **1953**. Yearly food habits of the river otter in the Thompson Lakes region, northwestern Montana, as indicated by scat analysis. M.S. thesis. Montana State University, Bozeman, MT 30 p. (Amer. Mid. Naturalist 54(2):299-313).
- Lechleitner, R.R. **1953**. An analysis of same age criteria in a domestic and a wild population of mink (*Mustela vison*). M.S. thesis. University of Montana, Missoula, MT. 60 p. (J. Mammalogy 35(4):496-503).
- Quick, H.F. **1953**. Wolverine, fisher, and marten studies in a wilderness region. Transactions of North American Wildlife Conference 18:512-533.
- Casagrande, L.G. **1955**. A study of beaver-waterfowl relations in the mountainous area of Beaverhead County, Montana. M.S. Thesis. Montana State University, Bozeman, MT. 33 p.
- Hawley, V.D. **1955**. The ecology of the marten in Glacier National Park. M.S. thesis. University of Montana, Missoula, MT. 131 p. (J. Mammalogy 38(2):174-184; J. Wildl. Mgmt. 26(1):55-74).
- Newby, F.E. and P.L. Wright. **1955**. Distribution and status of the wolverine in Montana. Journal of Mammology 36:248-253.
- Newby, F.E. **1956**. Fur production in Montana. Montana Wildlife. 6(1):7-13.
- Quick, H.F. **1956**. Effects of exploitation on a marten population. J. Wildl. Manage. 20:267-274.
- Hawley, V.D., and F.E. Newby. **1957**. Marten home ranges and population fluctuations. J. Mamm. 38:174-184.
- Mitchell, J.L. **1957**. A study of mink movements and populations on the lower Madison River, Montana. M.S. thesis. Montana State University, Bozeman, MT. 19 pp.
- Newby, F.E. **1957**. Marten population status. Pages 63-70 in Completion report project W-49-R-6. Montana Fish and Game Department, Helena, MT.
- Weckwerth, R.P. **1957**. The relationship between the marten population and the abundance of small mammals in Glacier National Park. M.S. thesis. University of Montana, Missoula, MT 76 p. (J. Wildl. Mgmt 26(1):55-74; J. Wildl. Mgmt 27(1):93-98).
- Newby, F.E. **1958**. A new approach to predator management in Montana. Montana Wildlife. August, pgs. 22-27.
- Adams, L. **1959**. An analysis of a population of snowshoe hares in northwestern Montana. Ecology Monograph 29(2):141-170.

Jonkel, C.J. **1959**. An ecological and physiological study of pine marten. M.S. thesis. University of Montana, Missoula, MT. 81 p. (J. Wildl. Mgmt. 27(1):93-98).

Newby, F.E. **1960**. Unpublished records. Montana Fish and Game, Helena, Montana.

Hawley, V.D. **1960**. Fisher are returned to Montana forests. Montana Wildlife, July 1960: 16-18.

Mitchell, J.L. **1961**. Mink movements and populations on a Montana river. Journal of Wildlife Management 25(1):48-54.

Weckwerth, R.P., and V.D. Hawley. **1962**. Marten food habits and population fluctuations in Montana. J. Wildl. Manage. 26:55-74.

Jonkel, C.J. and R.P. Weckwerth. **1963**. Sexual maturity and implantation of blastocysts in the wild pine marten. Journal of Wildlife Management 27:93-98.

Newby, F.E. and J.J. McDougal. **1964**. Range extension of the wolverine in Montana. Journal of Mammology. 45:485-487.

Weckwerth, R.P., and P.L. Wright. **1968**. Results of transplanting fishers in Montana. J. Wildl. Manage. 32(4):977-979.

Mitchell, J., K. Greer, and R. Weckwerth. **1971**. Furbearers. Pages 197-205 in Mussehl, T.W. and F.W. Howell, editors. Game management in Montana. Montana Fish and Game Department, Helena, MT. 238 pp.

Mitchell, J. and K. Greer. **1971**. Predators. Pages 206-217 in Mussehl, T.W. and F.W. Howell, editors. Game management in Montana. Montana Fish and Game Department, Helena, MT. 238 pp.

Moore, R.E. and N.S. Martin. **1980**. A recent record of swift fox (*Vulpes velox*) in Montana. J. Mamm. 61:161.

Schladweiler, P. **1980**. The effects of coyotes on big game populations in Montana. Montana Dept. Fish and Game, Job Final Report, Fed. Aid Project. W-120-R-11. 78 pp.

Burnett, G.W. **1981**. Movements and habitat use of the American marten in Glacier National Park, Montana. M.S. Thesis. University of Montana, Missoula, MT. 130 pp.

Greer, K.R., and D. Palmisciano. **1981**. Wildlife investigations laboratory job progress report. Appendix 2: bobcat carcasses from the registered trapping harvest for 1978-80. Fed. Aid Wildl. Restoration Proj. W-120-R-12. Montana Dept. Fish, Wildlife and Parks, Bozeman, MT 14 pp.

Hash, H.S. **1981**. Ecology of the bobcat in a coniferous forest environment in western Montana. Fed Aid Wildl. Restoration Proj. W-120-R-12, Project 3, Job 1. Montana Dept. Fish, Wildlife and Parks, Helena, MT. 13 p.

Hornocker, M.G. and H.S. Hash. **1981**. Ecology of the wolverine in northwestern Montana. Canadian Journal of Zoology 59:1286-1301.

Knowles, P.R. **1981**. Habitat selection, home range size, and movements of bobcats in northcentral Montana. M.S. Thesis. Univ. of Montana, Missoula.

Hash, H.S. **1982**. Statewide population studies, distribution, and habitat use of bobcat, Canada lynx, otter, wolverine, marten, and fisher. Fed. Aid Wildl. Job Prog. Rep. W-120-R-13. Montana Dept. Fish, Wildlife and Parks, Helena.

- Zackheim, H. **1982**. Ecology and population status of the river otter in southwestern Montana. M.S. Thesis. University of Montana, Missoula. 100 p.
- Swenson, J.E., S.J. Knapp, P.R. Martin, and T.C. Hinz. **1983**. Reliability of aerial cache surveys to monitor beaver population trends on prairie rivers in Montana. *Journal Wildlife Management* 47(3):697-703.
- Hamlin, K.L., S.J. Riley, D. Pyrah, A.R. Dood, and R.J. Mackie. **1984**. Relationships among mule deer fawn mortality, coyotes, and alternative prey species during summer. *J. Wildl. Manage.* 48:489-499.
- Pyrah, D. **1984**. Social distribution and population estimates of coyotes in north-central Montana. *J. Wildl. Mgmt.* 48(3):679-690.
- Smith, D.S. **1984**. Habitat use, home range and movements of bobcats in western Montana. M.S. Thesis. University of Montana, Missoula.
- Brainard, S.M. **1985**. Ecology of the bobcat in a coniferous forest environment in western Montana. M.S. Thesis. University of Montana, Missoula, MT 58 p.
- Knowles, P.R. **1985**. Home range size and habitat selection of bobcats (*Lynx rufus*) in north-central Montana. *Canadian Field-Nat.* 99:6-12.
- Giddings, B.J. **1986**. Home range size, movements and habitat use by bobcats in a prairie rangeland environment. M.S. Thesis. Montana State University, Bozeman. 62 p.
- Bissell, G.N. and R. Bown. **1987**. Effects of water level fluctuations on aquatic furbearer distribution, abundance, and habitat in the northern Flathead Valley. Final Report. Montana Dept. Fish, Wildlife and Parks, Kalispell
- Hash, H. **1987**. Wolverine. Pages 574-585 in Novak, M., J.A. Baker, M.E. Obbard, and B. Malloch, editors. *Wild furbearer management and conservation in North America*. Ministry of Natural Resources, Toronto, Ontario, Canada.
- Risdahl, G.L. **1988**. Ecology and reproduction of bobcats in southeastern Montana during a period of low lagomorph density. M.S. Thesis. Montana State University, Bozeman.
- Easter-Pilcher, A. **1990**. Cache size as an index to beaver colony size in northwestern Montana. *Wildlife Society Bulletin* 18:110-113.
- Giddings, B.J., G.L. Risdahl, and L.R. Irby. **1990**. Bobcat habitat use in southeastern Montana during periods of high and low lagomorph abundance. *Prairie Naturalist* 22(4):249-258.
- Dronkert-Egnew, A.E. **1991**. River otter population status and habitat use in northwestern Montana. M.S. Thesis. University of Montana, Missoula. 112 p.
- Fager, C. **1991**. Harvest dynamics and winter habitat use of the pine marten in southwest Montana. M.S. Thesis. Montana State University, Bozeman, MT. 73 p.
- FaunaWest Wildlife Consultants. **1991**. An ecological and taxonomic review of the swift fox (*Vulpes velox*) with special reference to Montana. Montana Dept. Fish, Wildlife and Parks, Helena, MT. 49 pp.
- FaunaWest Wildlife Consultants. **1991**. A bibliography of literature and papers pertaining to the swift and kit foxes. Montana Dept. Fish, Wildlife and Parks, Helena, MT. 49 pp.

Roy, K.D. **1991**. Ecology of reintroduced fishers in the Cabinet Mountains of northwestern Montana. M.S. Thesis. University of Montana, Missoula, MT

Waller, A.J. **1992**. Seasonal use of river otters in northwestern Montana. M.S. Thesis. University of Montana, Missoula. 75 p.

Bergman, H.L., M.J. Szumski, and D.J. Dziak. **1993**. Final Report: Exposure to and injury from environmental metal contamination on semi-aquatic mammals in the upper Clark Fork River, Montana. Red Buttes Environmental Biology Laboratory, Univ. of Wyoming, Laramie.

Heinemeyer, K.S. **1993**. Temporal dynamics in the movements, habitat use, activity and spacing of reintroduced fishers in northwestern Montana. M.S. thesis. Univ. of Montana, Missoula. 158 p.

Kujala, Q.J. **1993**. Winter habitat selection and habitat status of pine marten in southwest Montana. M.S. thesis. Montana State Univ., Bozeman. 58 p.

Vore, J. **1993**. Guidelines for the reintroduction of beaver into southwest Montana streams. Montana Dept. Fish, Wildlife and Parks, Helena, MT. 34 p.

Coffin, K.W. **1994**. Population characteristics and winter habitat selection by pine marten in southwest Montana. M.S. thesis. Montana State Univ., Bozeman. 94 p.

Giddings, B. and C.J. Knowles. **1995**. The current status of swift fox in Montana. Pages 101-120 in Allen, S.H., J. Whitaker-Hoagland, and E. Dowd-Stukel, editors. Report of the swift fox conservation team. North Dakota Game and Fish Dept., Bismark, ND. 170 pp.

Giddings, B. **1996**. Distribution and investigations of swift fox in Montana. Pages 25-29 in Luce, B. and F. Lindsey, editors. Annual report of the swift fox conservation team. Wyoming Game and Fish Dept., Lander, WY. 110 pp.

Aune, K.E., and P. Schlandweiler. **1997**. Age, sex structure, and fecundity of the American marten. Pages 61-77 in G. Proulx, H.N. Harestad, and P.M. Woodard, eds. *Martes: Taxonomy, Ecology, and Management*. The Provincial Museum of Alberta, Edmonton, Alberta, Canada.

Coffin, K.W., Q.J. Kujala, R.J. Douglas, and L.R. Irby. **1997**. Interactions among marten prey availability, vulnerability, and habitat structure. Pages 199-210 in G. Proulx, H.N. Harestad, and P.M. Woodard, eds. *Martes: Taxonomy, Ecology, and Management*. The Provincial Museum of Alberta, Edmonton, Alberta, Canada.

Kahn, R., L. Fox, P. Horner, B. Giddings, and C. Roy. **1997**. Conservation Assessment and conservation strategy for swift fox in the United States. Colorado Division of Wildlife, Denver, CO. 54 p.

Zimmerman, A.L. and B. Giddings. **1997**. Preliminary findings of swift fox studies in Montana. Pages 27-44 in Giddings, B., editor. Swift fox conservation team annual report. Montana Fish, Wildlife & Parks, Helena, MT. 105 pp.

Wisz, M. **1998**. Survey for American marten (*Martes Americana*) in the Little Belt mountains, Lewis and Clark National Forest. Report for Montana Fish, Wildlife and Parks, Helena, MT. 5 pp.

Zimmerman, A.L. **1998**. Reestablishment of swift fox in north central Montana. M.S. thesis. Montana State University, Bozeman. 44 pp.

- Giddings, B. **1999**. Swift fox management activities in Montana. Pages 18-22 in Roy, C., editor. 1998 swift fox conservation team annual report. 107 pp.
- Wis, M.S. **1999**. Islands in the Big Sky: equilibrium biogeography of isolated mountain ranges in Montana. M.S. thesis. Dept. of Environmental, Population and Organismic Biology. University of Colorado. Boulder, CO. 41 pp.
- Giddings, B. **2000**. Montana swift fox management activities. Pages 191-201 in Schmitt, G.C., editor. Swift fox conservation team 1999 annual report. New Mexico Dept. of Game and Fish, Santa Fe, NM. 201 pp.
- Cegelski, C. **2001**. An evaluation of genetic diversity, gene flow, and population genetic structure among wolverine (*Gulo gulo*) populations in the Rocky Mountains. M.S. Thesis. University of Idaho, Moscow, Idaho. 67 p.
- Giddings, B. **2001**. Montana swift fox management activities. Pages 40-44 in Schmitt, G.C. and B. Oakleaf, editors. Swift fox conservation team 2000 annual report. New Mexico Dept. of Game and Fish, Santa Fe, NM. 103 pp.
- Moehrensclager, A., and C. Moehrensclager. **2001**. Census of swift fox (*Vulpes velox*) in Canada and northern Montana: 2000-2001. Alberta Sustainable Resource Development, Fish and Wildlife Division, Alberta Species at Risk Report No. 24. Edmonton, AB. 21 p.
- Coffin, K., C. Fager, Q. Kujala, L. Irby, and R. Douglas. **2002**. Winter ecology of American marten in southwestern Montana. Montana Fish, Wildlife and Parks, Wildlife Division, Technical Bulletin, Fed. Aid Project W-100-R. Helena, MT. 54 p.
- Giddings, B. **2002**. Montana swift fox management activities, 2001-2002. Pages 48-52 in Peek, M.S., editor. Swift fox conservation team 2001 annual report. Kansas Dept. of Wildlife and Parks, Emporia, KS. 120 pp.
- Cegelski, C.C., L.P. Waits, and N.J. Anderson. **2003**. Assessing population structure and gene flow in Montana wolverines (*Gulo gulo*) using assignment-based approaches. *Molecular Ecology* 12, 2907-2918.
- Dowd-Stukel, E., C. Slivinski, and B. Giddings. **2003**. A design for species restoration – development and implementation of a conservation assessment and conservation strategy for swift fox in the United States. In Sovada, M.A. and L.Carbyn, editors. The swift fox: ecology and conservation of swift foxes in a changing world. Canadian Plains Research Center, University of Regina, Regina, Saskatchewan. 250 pp.
- Giddings, B. **2003**. Management activities for swift fox in Montana. Pages 10-13 in Grenier, M., editor. Swift fox conservation team 2002 annual report. Wyoming Game and Fish Dept., Lander, WY. 73 pp.
- Knowles, C.J., P.R. Knowles, B. Giddings, and A.R. Dood. **2003**. The historic and recent status of the swift fox in Montana. In Sovada, M.A. and L.Carbyn, editors. The swift fox: ecology and conservation of swift foxes in a changing world. Canadian Plains Research Center, University of Regina, Regina, Saskatchewan. 250 pp.
- Kolbe, J.A., J.R. Squires, and T.W. Parker. **2003**. An effective box trap for capturing lynx. *Wildlife Society Bulletin* 31(4):1-xxx.
- Vinkey, R.S. **2003**. An evaluation of fisher (*Martes pennanti*) introductions in Montana. M.S. Thesis. University of Montana, Missoula, Montana. 97 p.
- Zimmerman, A.L., L. Irby, and B. Giddings. **2003**. The status and ecology of swift foxes in north-central Montana. In Sovada, M.A. and L.Carbyn, editors. The swift fox: ecology and conservation of swift foxes in a changing world. Canadian Plains Research Center, University of Regina, Regina, Saskatchewan. 250 pp.

- Squires, J.R., K.S. McKelvey, and L.F. Ruggiero. **2004**. A snow-tracking protocol used to delineate local lynx, *Lynx Canadensis*, distributions. *Canadian Field-Naturalist* 118(4):583-589.
- Ausband, D.E. **2005**. Assessing the success of swift fox reintroductions on the Blackfeet Indian Reservation, Montana. M.S. Thesis. University of Montana, Missoula, MT. 52 pp.
- Giddings, B. and R. Rauscher. **2005**. Monitoring population status of swift fox in Montana. Pages 29-33 in Grenier, M. and H. Whitlaw, editors. Swift fox conservation team 2003 annual report. Wyoming Game and Fish Dept., Lander, WY. 108 pp.
- Kolbe, J.A. **2005**. The effects of snowmobile trails on coyote movements within lynx home ranges. M.S. Thesis. University of Montana, Missoula, MT. 35 pp.
- Ulizio, T.J. **2005**. A noninvasive survey method for detecting wolverine. M.S. Thesis. University of Montana, Missoula, MT. 56 pp.
- Ausband, D.E. and E.A. Ausband. **2006**. Notes: observations of interactions between swift fox and American badger. *Prairie Naturalist* 38(1):63-64.
- Giddings, B. **2006**. Swift fox monitoring activities in Montana. Pages 7-9 in Stuart, J.N. and S. Wilson, editors. Swift fox conservation team annual report for 2004. New Mexico Dept. of Game and Fish, Santa Fe, NM. 86 pp.
- McKelvey, K.S., J. Von Kienast, K.B. Aubry, G.M. Koehler, B.T. Maletzke, J.R. Squires, E.L. Lindquist, S. Loch, and M.K. Schwartz. **2006**. DNA analysis of hair and scat collected along snow tracks to document the presence of Canada lynx. *Wildlife Society Bulletin*:461-455
- Moehrensclager, A., and C. Moehrensclager. **2006**. Population census of reintroduced swift fox (*Vulpes velox*) in Canada and northern Montana: 2005/2006. Centre for Conservation Research Report No. 1. Calgary Zoo, Calgary, Alberta, Canada. 32 p.
- Moehrensclager, A., S. Alexander, and T. Bricchieri-Colombi. **2006**. Habitat suitability and population viability analysis for reintroduced swift foxes in Canada and northern Montana. Centre for Conservation Report No. 2, Calgary, Alberta, Canada. 30 pp.
- Vinkey, R.S., M.K. Schwartz, K.S. McKelvey, K.R. Foresman, K.L. Pilgrim, B.J. Giddings, and E.C. LoFroth. **2006**. When reintroductions are augmentations: The genetic legacy of fishers (*Martes pennanti*) in Montana. *Journal of Mammalogy*, 87(2):265-271.
- Ulizio, T.J., J.R. Squires, D. Pletscher, M.K. Schwartz, J. Claar, and L.F. Ruggiero. **2006**. The efficacy of obtaining genetic-based identifications from putative wolverine snow tracks. *Wildlife Society Bulletin* 34:1326-1332.
- Aubry, K.B., K.S. McKelvey, and J.P. Copeland. **2007**. Distribution and broadscale habitat relations of the wolverine in the contiguous United States. *Journal of Wildlife Management* 71(7):2147-2158.
- Ausband, D.E. and K.R. Foresman. **2007**. Swift fox reintroductions on the Blackfeet Indian Reservation, Montana, USA. *Biological Conservation* 136:423-430.
- Ausband, D.E. and K.R. Foresman. **2007**. Dispersal, survival, and reproduction of wild-born, yearling swift foxes in a reintroduced population. *Canadian Journal of Zoology* 85:185-189.

- Brock, B.L., R.M. Inman, K.H. Inman, A.J. McCue, M.L. Packila, and B. Giddings. **2007**. Broad-scale wolverine habitat in the conterminous Rocky Mountain states. Pages 21-53 in Cumulative Report. Greater Yellowstone Wolverine Program, Wildlife Conservation Society. Bozeman, MT. 128 pp.
- Giddings, B. **2007**. Monitoring resident swift fox populations during 2005 and 2006 in Montana. Pages 9-15 in Dowd Stukel, E. and D.M. Fecske, editors. Swift fox conservation team report for 2005-2006. South Dakota Dept. Game, Fish and Parks, Pierre, SD. 97 pp.
- Inman, R.M., K.H. Inman, A.J. McCue, and M.L. Packila. **2007**. Wolverine Harvest in Montana: Survival rates and spatial considerations for harvest management. Pages 85-97 in Cumulative Report. Greater Yellowstone Wolverine Program, Wildlife Conservation Society. Bozeman, MT. 128 pp.
- Kolbe, J.A., J.R. Squires, D.H. Pletscher, and L.F. Ruggiero. **2007**. The effects of snowmobile trails on coyote movements within lynx home ranges. *Journal of Wildlife Management* 71(5):000-000.
- Kolbe, J.A. and J.R. Squires. **2007**. Circadian activity patterns of Canada lynx in western Montana. *Journal of Wildlife Management* 71(5):000-000.
- Schwartz, M.K. **2007**. Ancient DNA confirms native Rocky Mountain fisher (*Martes pennant*) avoided early 20th century extinction. *Journal of Mammalogy*, 88(4):921-925.
- Squires, J.R., J.P. Copeland, T.J. Ulizio, M.K. Schwartz, and L.F. Ruggiero. **2007**. Sources and patterns of wolverine mortality in western Montana. *Journal of Wildlife Management* 71(7):2213-2220.
- Squires, J.R. and L.F. Ruggiero. **2007**. Winter prey selection of Canada lynx in northwestern Montana. *Journal of Wildlife Management* 71:310-315.
- Anderson, N.J. and K.E. Aune. **2008**. Fecundity of female wolverine in Montana. *Intermountain Journal of Sciences*. Vol. 14, No. 1-3:17-30.
- Copeland, J.P. and R.E. Yates. **2008**. Wolverine population assessment in Glacier National Park: Comprehensive summary update. USDA Forest Service, Rocky Mountain Research Station, Missoula, MT. 15 pp.
- Giddings, B. **2008**. Swift fox populations in Montana, 2007. Pages 5-7 in Krueger, B. and M. Ewald, editors. Swift fox conservation team report for 2007. Wyoming Game and Fish Dept., Laramie, WY. 70 pp.
- Kudray, G.M., and T. Schemm. **2008**. Wetlands of the Bitterroot Valley: Change and ecological functions. Montana Natural Heritage Program. Prepared for Montana Dept. Environmental Quality, Helena, MT. 32 p.
- Squires, J.R., N.J. DeCesare, J.A. Kolbe and L.F. Ruggiero. **2008**. Hierarchical den selection of Canada lynx in western Montana. *Journal of Wildlife Management* 72:1497-1506.
- Giddings, B. **2009**. Montana annual swift fox report, pages 17-20 in Isakson, P. editor. Swift fox conservation team: Report for 2008. North Dakota Game and Fish Dept., Bismarck, ND. 67 p.
- Inman, R.M., M.L. Packila, K.H. Inman, B. Aber, R. Spence, and D. McCauley. **2009**. Greater Yellowstone Wolverine Program, Progress Report-December 2009. Wildlife Conservation Society, North America Program, General Report, Bozeman, Montana.
- Rauscher, R.L. **2009**. Fort Peck Indian Reservation swift fox translocation report. Montana Fish, Wildlife and Parks, Region Six, Glasgow, MT. 5 pp.

- Schwartz, M.K., J.P. Copeland, N.J. Anderson, J.R. Squires, R.M. Inman, K.S. McKelvey, K.L. Pilgrim, L.P. Watts, and S.A. Cushman. **2009**. Wolverine gene flow across a narrow climatic niche. *Ecology* 90(11):3222-3232.
- Sovada, M.A., R.O. Woodward, and L.D. Igl. **2009**. Historical range, current distribution, and conservation status of the swift fox, *Vulpes velox*, in North America. *The Canadian Field-Naturalist* Vol. 123: 346-367.
- Tremain, K.M. **2009**. Assessing climate induced range shifts in bobcats (*Lynx rufus*) in the northern Rockies. M.S. Thesis. Columbia University, New York, NY. 73 pp.
- Squires, J.R., N.J. DeCesare, J.A. Kolbe, and L.F. Ruggiero. **2010**. Seasonal resource selection of Canada lynx in managed forests of the Northern Rocky Mountains. *Journal of Wildlife Management* 74:1648-1660.
- Dowd Stukel, E., editor. **2011**. Conservation assessment and conservation strategy for swift fox in the United States – 2011 update. South Dakota Game, Fish and Parks, Pierre, SD. 98 pp.
- Giddings, B. **2011**. Montana 2009 and 2010 swift fox report. Pages 13-17 in Bly, K. , editor. Swift fox conservation team: report for 2009-2010. World Wildlife Fund, Bozeman, MT and Montana Fish, Wildlife and Parks, Helena, MT.
- Inman, R.M., M.L. Packila, K.H. Inman, A.J. McCue, G.C. White, J. Persson, B.C. Aber, M.L. Orme, K.L. Alt, S.L. Cain, J.A. Fredrick, B.J. Oakleaf, and S.S. Sartorius. **2011**. Spatial Ecology of wolverines at the southern periphery of distribution. *Journal of Wildlife Management, Research Article* 9999:1-15
- Inman, R.M, B.L. Brock, K.H. Inman, S.S. Sartorius, B.C. Aber, B. Giddings, S.L. Cain, M.L. Orme, J.A. Fredrick, B.J. Oakleaf, K.L. Alt, E. Odell, and G. Chapron. **2011**. Developing a spatial framework and conservation priorities for a wolverine metapopulation.
- Magoun, A.J., C.D. Long, M.K. Schwartz, K.L. Pilgrim, R.E. Lowell, and P. Valkenburg. **2011**. Integrating motion-detection cameras and hair snags for wolverine identification. *Journal of Wildlife Management* 75(3):731-739.
- Newton, D.E. **2011**. Northern river otter population assessment and connectivity in western Montana. M.S. Thesis. University of Montana, Missoula, MT. 57 pp.
- Alexander, J.L. **2012**. Swift fox distribution and population connectivity in eastern Montana. M.S. Thesis. St. Cloud State University, St. Cloud, MN. 74 pp.
- Inman, R.M, M.L. Packila, K.H. Inman, A.J. McCue, G.C. White, J. Persson, B.C. Aber, M.L. Orme, K.L. Alt, S.L. Cain, J.A. Fredrick, B.J. Oakleaf, and S.S. Sartorius. **2012**. Spatial ecology of wolverines at the southern periphery of distribution. *Journal of Wildlife Management* 76(4):778-792.
- Newton, D. and K.R. Foresman. **2012**. Status of Northern river otters in the Upper Clark Fork River. Report to Montana Natural Resource Damage Program and Montana , Fish, Wildlife and Parks. 40 pp.
- Inman, R.M., A.J. Magoun, J. Persson, and J. Mattisson. **2012**. The wolverine's niche: Linking reproductive chronology, caching, competition, and climate. *Journal of Mammalogy* 93(3):634-644.
- Squires, J.R., L.E. Olson, D.L. Turner, N.J. DeCesare, and Jay Kolbe. **2012**. Estimating detection probability for Canada lynx *lynx canadensis* using snow-track surveys in the northern Rocky Mountains, Montana, USA. *Wildlife Biology* 18:215-224.

Giddings, B. **2013**. Swift fox monitoring and management in Montana. Pages 9-11 in Stratman, M.R., editor. Swift fox conservation team: report for 2011-2012. Colorado Division of Parks and Wildlife. Fort Collins, CO.

Inman, R.M. **2013**. Wolverine ecology and conservation in the western United States. PhD Thesis No. 2013:4. Swedish University of Agricultural Sciences, Uppsala, Sweden.

Inman, R.M., B.L. Brock, K.H. Inman, S.S. Sartorius, B.C. Aber, B. Giddings, S.L. Cain, M.L. Orme, J.A. Fredrick, B.J. Oakleaf, K.L. Alt, E. Odell, and G. Chapron. **2013**. Developing priorities for metapopulation conservation at the landscape scale: Wolverines in the Western United States. *Biological Conservation* 166 (2013) 276-286.

Newbury, R.K. **2013**. Behavioral ecology of the bobcat in a region with deep winter snows. University of British Columbia, Okanagan, B.C. 181 pp.

Schwartz, M.K., N.J. DeCesare, B.S. Jimenez, J.P. Copeland, and W.E. Melquist. **2013**. Stand- and landscape-scale selection of large trees by fishers in the Rocky Mountains of Montana and Idaho. *Forest Ecology and Management* 305(2013):103-111.

Squires, J.R., N.J. DeCesare, L.E. Olson, J.A. Kolbe, M. Hebblewhite, and S.A. Parks. **2013**. Combining resource selection and movement behavior to predict corridors for Canada lynx at their southern range periphery. *Biological Conservation* 157:187-195.