Montana Fish, Wildlife & Parks Region 2 Wildlife Quarterly September 2017



Montana Fish, Wildlife & Parks Region 2 Wildlife Quarterly

September 2017



Region 2, 3201 Spurgin Road, Missoula MT 59804, 406-542-5500

Find the Quarterly online at fwp.mt.gov/ regions/r2/WildlifeQuarterly

Region 2 Wildlife Staff

Liz Bradley, Wildlife Biologist, Missoula-West, <u>Ibradley@mt.gov</u>, 406-542-5515

Dave Dickson, Wildlife Management Areas Maintenance, <u>ddickson@mt.gov</u>, 406-542-5500

Kristi DuBois, Wildlife Biologist, Nongame, <u>kdubois@mt.gov</u>, 406-542-5551

Julie Golla, Wildlife Biologist, Upper Clark Fork, <u>jgolla@mt.gov</u>, 406-381-1268

Scott Eggeman, Wildlife Biologist, Blackfoot, <u>seggeman@mt.gov</u>, 406-542-5542

James Jonkel, Bear and Cougar Management Specialist, <u>jajonkel@mt.gov</u>, 406-542-5508

Kendra McKlosky, Hunting Access Coordinator, <u>kmcklosky@mt.gov</u>, 406-542-5560

Rebecca Mowry, Wildlife Biologist, Bitterroot, <u>rmowry@mt.gov</u>, 406-363-7141

Adam Sieges, Wildlife Management Areas Maintenance, 406-693-9083

Tyler Parks, Wolf-Carnivore Management Specialist, <u>tparks@mt.gov</u>, 406-542-5500

Tyler Rennfield, Conservation Specialist, <u>trennfield@mt.gov</u>, 406-542-5510

Brady Shortman, Wildlife Management Areas Maintenance Supervisor, <u>bshortman@mt.gov</u> 406-693-9083

Mike Thompson, Regional Wildlife Manager, <u>mthompson@mt.gov</u>, 406-542-5516

Bob White, Wildlife Management Areas Maintenance, 406-542-5500

Bob Wiesner, Cougar and Bear Management Specialist, 406-542-5508

Statewide Research Staff Housed at Region 2 Headquarters:
Nick DeCesare, Wildlife Biologist, Moose Research Project, ndecesare@mt.gov, 406-542-5500
Ben Jimenez, Research Technician, bjimenez@mt.gov, 406-542-5500

Communication & Education Division:

Vivaca Crowser, Regional Information & Education Program Manager, vcrowser@mt.gov, 406-542-5518

The Region 2 Wildlife Quarterly is a product of Montana Fish, Wildlife & Parks; 3201 Spurgin Road; Missoula 59804. Its intent is to provide an outlet for a depth of technical information that normally cannot be accommodated by commercial media, yet we hope to retain a readable product for a wide audience. While we strive for accuracy and integrity, this is not a peer-refereed outlet for original scientific research, and results are preliminary. October 2015 was the inaugural issue.



stantly learning and practicing the terms of coexistence.

everywhere, there is tolerance, mixed with conflict.



Bob Wiesner handles an immobilized, fall cub grizzly (left).

Eric Graham checks vital signs of an immobilized adult before fitting a radio collar on its neck (below).



Every year, FWP Wildlife Specialists in Region 2 participate in capturing and collaring adult female grizzly bears, as part of Montana's protocol for monitoring survival and other vital rates, which are used to estimate the population trend.



Tracking the Grizzly Population

In 2016, FWP researchers Cecily M. Costello, Richard D. Mace and Lori Roberts published a report entitled, *Grizzly Bear Demographics in the Northern Continental Divide Ecosystem, Montana: Research Results (2004-2014) and Suggested Techniques for Management of Mortality.*

A principal finding of that report was that the rates of births and deaths among radioed male and female grizzly bears can be used in a model to estimate the trend of the grizzly bear population. Radios must be de-

ployed on a representative sample of adult female and male bears across the entirety of the Northern Continental Divide Ecosystem, of which the Blackfoot Watershed in Region 2 (including the Clearwater) comprises its southern extreme.

Between May and July 2017, Jonkel's team trapped five male grizzlies in Region 2 as their latest contribution toward the larger FWP monitoring effort. One of the five bears was fitted with a GPS collar; the other males were captured and released incidentally while hoping to capture and collar a female.

This spring, four additional male grizzlies, which were captured and collared last year, dropped their collars. All four collars were retrieved and the GPS locations downloaded. These bears had been instrumented with GPS collars in conjunction with a griz-


About 60 grizzly bears live in the Blackfoot-Clearwater Watershed.


> - James Jonkel, FWP Region 2 Bear Specialist, 2017

zly/black bear project that began in May 2015. Brittani Johnson, a graduate student at Montana State University, is studying electric fence permeability to bears and the impacts of electric fence on grizzly bear movements and behavior in the Blackfoot Watershed.

One additional female grizzly bear, wearing an ear transmitter, is also being tracked on a weekly basis. This female was captured in July 2016, and was reinstrumented with an ear transmitter rather than a neck collar because of a wound on its neck.









Why?

Why does the bear rub in the woods?

"I've always thought of rubbing as a marking, territorial display," says Jonkel. "They're leaving their calling card. And, males seem to use them more in breeding season when they're crossing other territories looking for mates."

Unlike most of us, who'll never observe a bear rubbing on a tree, post, or most anything else that offers resistance, he's personally observed bears rubbing, and bears coming behind the first one, and the second, to rub, seemingly in response.

"They definitely behave as if they're gathering information at rub sites. They sniff and I've seen them step hard in the footsteps of a bear that rubbed earlier, grinding their feet as if to rub-out the other bear."

After hanging with Jamie, we've all become more aware of the claw and bite marks that bears leave on trees and posts, way more often than we used to notice them. Typically, bite marks can be seen running horizontally across the tree or post. Claw marks usually run vertically.

"We collect hairs that the bears leave behind when they rub, which provides us the opportunity to improve our bank of DNA data for the identification of individual bears. And we use posts, and sometimes set posts, to attract bears to our remote camera sets, which helps us document travel routes and range extensions."

DNA collected from rub trees and other hair collection sets in the Northern Continental Divide Ecosystem (NCDE) was used to establish a baseline estimate of grizzly bears, in ground breaking work led by Kate Kendall of the U. S. Geological Survey in Glacier National Park (retired). Jonkel's team and others working in

the NCDE are collecting the data that allows FWP and partners to assess whether the bear population is increasing, decreasing, shrinking or expanding in relation to that baseline estimate.







Blackfoot Community



Eric Graham is a Wildlife Technician supported by the Blackfoot Challenge. Eric has spent his life in western Montana acquiring and practicing the skills needed to help wildlife and people live better together. A hard worker and a practical man, Eric is a perfect fit with the Challenge, which is a landowner-based group whose mission is to *coordinate* efforts that conserve and enhance the natural resources and rural way of life in the Blackfoot Watershed for present and future generations. A friend of wildlife, of landowners, and of FWP.

It would be hard to imagine a community of people who derive deeper satisfaction from wildlife than the ranchers whose lands grow it alongside their livestock and crops. All they require in return is the opportunity to make a good life for them-




selves and promise for the future. FWP's role is as a partner, in the role of supplying wildlife management expertise and some of the wherewithal to help landowners develop solutions for conflicts as they arise and can be foreseen.






It's hay barley, tall enough to hide these bucks when they're not jumping to scout their next lunge ahead. Increasingly, grizzlies have learned to feed and live in this particular crop, and in crops nearby, digging day beds in the cool ground to escape the midsummer heat. A variety of electric fencing options are being

tested in the Blackfoot to guide grizzlies around conflict areas, which can help bears avoid making mistakes. "It's not something you do and then walk away," Jonkel says. "Every day is a new day when we'll need to watch and adapt."







Type "Ethyl grizzly bear" in your Internet browser to access published stories of this famous, wide-ranging bear. Ethyl was a radio-collared female grizzly that journeyed down to the Missoula area from Bigfork and denned just outside of Missoula in the fall of 2012. In March of 2013, she emerged from her den on Wisherd Ridge and journeyed back to the Flathead Valley. In September of 2013 she returned to the Missoula Valley where she spent a month in the vicinity of the Snowbowl Ski Area. From there she began an incredible journey that took her east to the Coeur d'Alene Mountains in North Idaho and then south into Idaho's Clearwater National Forest. Ethyl was the first grizzly documented to den south of Interstate 90, near St Regis, since the 1930's.

The 2,800-mile track of this grizzly bear is useful for helping FWP identify and consider travel routes for bears and other wildlife. While too much might be made of a single bear's

travels, it's remarkable how many species and individuals follow similar routes as they read the lay of the land and navigate passage. And while few, if any, other bears would make this entire journey, it's like-

ly that some bears would use the portions of this mapped route within their local ranges.





The Living With Bears Project

Western Montana, as many Montanans know it, is soccer, gymnastics, biking and gardening. And, we sometimes take for granted that this modern lifestyle is set in a surprisingly wild place, where it's still possible for a person—or a bear—to look overtop the development and inhabit a land that remains in large part undeveloped, if not unchanged.


FWP Region 2—Jonkel's team of Wildlife Specialists, along with FWP game wardens and biologists—invest countless hours and resources, almost every day, toward making room for wildlife around the "mine fields" of human development that wild animals encounter in their travels. Jonkel, in particular, has been a magician at cultivating partnerships and relationships with homeowner's associations, neighborhood networks, nonprofit organizations, other governmental agencies and boards, schools and universities, refuse collection businesses—just about anyone with an interest in getting involved.

Learning to live with wildlife by eliminating or reducing human-food rewards is an ethic and a lifestyle that's taking hold in western Montana. In the fall, black bears are keyed-in to chokecherries ripening along stream bottoms, drawing bears in close proximity to home-sites, often well within subdivisions


and longer established neighborhoods. In order to keep bears from ranging beyond the natural berry crops in the thick riparian corridors, partners help harvest apples on the wildland-urban interface. They offer solutions to homeowners for minimizing bears' exposure to garbage, using a variety of strategies. They help residents devise innovative ways to provide seed at their bird feeders, and to protect their chickens and bee-yards from depredations by bears. These and other efforts require continual education and information sharing to help residents new to the area understand the issues and needs.

When preventative measures fail, whether due to the failure of people to work together, or due to the behaviors of the occasional "bad bear," FWP is called upon to remove habituated bears, sometimes lethally. While the removal of habituated bears is critical to the success of the conflict prevention program overall, FWP is careful to remove bears as a last resort, only after property owners have secured the attractants on their properties and given the offending bear a fighting change to move on and remain in the wild.

For more information on living with bears and other wildlife, go to *missoulabears.org*.




Estimating Black



BLACK BEAR HARVEST PESEARCH & MAINAGEMENT IN MONTAINA

2011 FINAL REPORT

RICHARD D. MACE AND TONYA CHILTON-RADANDT





Bear Populations

In 2001, FWP began a research study to better assess from the barbs, and removed the contraptions from black bear harvest rates and population densities (Mace and Chilton-Radandt 2011).

In FWP Region 2, the Garnet Mountains (Hunting District 292) were selected as one of 11 areas across Montana for sampling the density of black bears.

Locations of sampling stations in the Garnets were randomly generated and crews of FWP employees were charged with finding and setting hair traps as close to each random sampling location as practical.

Each sampling station consisted of a stinking liquid bait, encircled by a strand of barbed wire. It was intended that bears would scoot under the wire to reach the bait, leaving hair on the barbs.

About one-two weeks after placing the baits, FWP employees revisited the hair traps, collected hairs

the field. Hair samples were sent to a laboratory for identification of the species and individual.

In the Garnets, an estimate of 169 black bears was generated, with 90% confidence that the actual number of bears would lie between 114 and 280. The Garnets sampling area encompassed 800 square miles, yielding an average density of 1 black bear per 5 square miles.

The mean population estimate for Montana was 13,307 black bears. The statewide average density was about 1 bear per 3 square miles. Highest densities were found in northwest Montana.

In related studies, Mace and Chilton-Radandt (2011) found an average litter size of 2.08 cubs per litter, and an average reproductive interval of 2.2 years.

