

2012 Annual Report

Pallid Sturgeon Population Assessment and Associated Fish Community Monitoring for the Missouri River: Segment 1



**Prepared for the U. S. Army Corps of Engineers - Missouri River Recovery Program
By:**

Tyler Haddix, Landon Holte, John Hunziker and Ryan Lott

**Montana Fish, Wildlife & Parks
Government Bldg.
East Kansas Street
Fort Peck, MT 59223**

June 2013

EXECUTIVE SUMMARY

Segment 1 of the Missouri River has been sampled during each year starting in 2006 to 2012, during both the sturgeon and fish community seasons. Segment 1 is situated directly downstream of Fort Peck Dam and is the most highly altered area of the Missouri River downstream of Fort Peck Dam within Montana. Segment 1 is used as a reference area, whereas the same river bend is sampled annually.

Over the seven years of sampling Segment 1 it has become apparent that the altered nature of the river precludes several native species from thriving. No pallid sturgeon *Schaphirynchus albus* have been collected during standardized sampling. In addition, no sicklefin chub *Macrhybopsis meeki* and only one sturgeon chub *M. gelida* have been collected during the fourteen sampling events over seven years. While some target species have been absent or somewhat rare, adult shovelnose sturgeon *S. platyrhynchus*, blue suckers *Cycleptus elongates* and sand shiners *Notropis stramineus* have consistently been collected.

During 2012 a total of 83 shovelnose sturgeon were collected in Segment 1. Shovelnose sturgeon averaged 647.3 mm in length and 1182.5 g in weight. The smallest shovelnose sturgeon captured measured 539 mm, with the largest specimen measuring 779 mm. No young-of-the-year or age-1 sized shovelnose sturgeon have been sampled within Segment 1 through the seven years of sampling.

The relative abundance of shovelnose sturgeon has remained relatively constant through the past seven years with the exception of 2009. During 2009 trammel net catch-per-unit-effort (CPUE) was at a seven year high with 4.5 fish/100 m during the fish community season. However, during the sturgeon season of the same year, CPUE was estimated at only 0.05 fish/100m, the lowest recorded trammel net CPUE for the same seven year period. During 2012 trammel net CPUE was higher during the sturgeon season (CPUE = 1.2 fish/100 m) than the fish community season (CPUE = 0.5 fish/100 m). Similarly, shovelnose sturgeon CPUE of trotlines was higher during the sturgeon season than during the fish community season.

The size distribution of shovelnose sturgeon captured within Segment 1 has not changed appreciably since 2006, with only adult sized fish being captured. Additionally, the length-weight relationship for Segment 1 caught shovelnose sturgeon has remained relatively constant over the seven sampling seasons.

Other native target species that were collected in 2012 include blue suckers, sauger *Sander canadense*, and sand shiners. A total of three blue suckers averaging 753.07 mm in length were captured, all of which were caught in trammel nets. Similarly, a total of three sauger were caught, averaging 426.3 mm and again all were captured in trammel nets. No western silvery minnows *Hybognathus argyritis* were caught in 2012.

During 2012 white suckers *Catostomus commersoni* were the most abundant fish sampled. A total of 760 white suckers were captured, 630 using mini fyke nets. White suckers sampled in mini fyke nets averaged 40.0 mm in length. Other gears captured adult white suckers. White suckers have been one if not the most common species collected within Segment 1 over the past seven years.

Although 2011 and 2012 were relatively high water years in Segment 1, our data do not support large increases in the populations of any species. Even though Segment 1 is upstream of the Fort Peck Spillway, flows were significantly increased during both 2011 and subsequently 2012. Montana Fish, Wildlife & Parks (FWP) management data indicated a large influx of adult walleye *S. vitreus*, sauger and northern pike *Esox lucius* in 2011 due to entrainment of fish through the Fort Peck Spillway. This increase in predators within the Missouri River could have masked any increases in production of cyprinids or catostomids by heavy predation. However, no empirical data exists to fully support this hypothesis. While pallid sturgeon responded to high flows in 2011 within Segments 2 and 3 by moving further upstream than prior years, little evidence exists to suggest mass movements of pallid sturgeon into Segment 1. Nonetheless, during 2012 FWP did collect one hatchery reared pallid sturgeon from the 1997 year class in the dredge cuts adjacent to Segment 1. This fish was collected in a gill net during FWP's annual trend netting. This was the first time a hatchery reared pallid sturgeon was documented within Segment 1. Even though it was only one fish, the presence of the first hatchery reared pallid sturgeon using waters directly downstream of Fort Peck Dam after a spill gives further support that flow enhancements out of Fort Peck Dam may benefit pallid sturgeon. This finding, coupled with the change in distribution of pallid sturgeon in Segments 2 and 3 as well as the documented use of adult fish in the Missouri River during 2011 greatly increases our knowledge of pallid sturgeon behavior during more natural Missouri River flows.

TABLE OF CONTENTS

| | |
|---------------------------------------------|----|
| Introduction..... | 1 |
| Study Area | 4 |
| Methods..... | 5 |
| Sample site selection and description | 5 |
| Sampling gear | 6 |
| Data Collection and Analysis..... | 8 |
| Results | |
| Pallid sturgeon | 12 |
| Targeted Native River Species..... | 12 |
| Missouri River Fish Community | 24 |
| Discussion | 24 |
| Acknowledgments..... | 27 |
| References | 28 |
| Appendices..... | 29 |

LIST OF TABLES

| | |
|------------------------------------------------------------------------------------------------------------------------------|----|
| Table 1. Presence/absence of all fishes collected in segments 1 through 3 of the Missouri River from 2006 through 2012. | 22 |
|------------------------------------------------------------------------------------------------------------------------------|----|

LIST OF FIGURES

| | |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 1. Map of segment 1 of the Missouri River with major tributaries, common landmarks, and historic stocking locations for pallid sturgeon. Segment 1 encompasses the Missouri River from Fort Peck Dam (River Mile 1771.5) to the confluence of the Milk River (River Mile 1760). | 10 |
| Figure 2. Hydrograph (2006-2012) and 2012 water temperature for segment 1 of the Missouri River. | 11 |
| Figure 3. Mean annual catch-per-unit-effort by season of selected species collected using trammel nets in segment 1 from 2006 through 2012. | 15 |
| Figure 4. Mean annual catch-per-unit-effort by season of selected species captured using trotlines in segment 1 from 2008 to 2012. | 16 |
| Figure 5. Mean annual catch-per-unit-effort by season of selected species collected using otter trawls in segment 1 from 2006 through 2012. | 17 |
| Figure 6. Length frequency histograms for all shovelnose sturgeon sampled in segment 1 from 2006 through 2012. | 18 |
| Figure 7. Weight-length relationship for all shovelnose sturgeon sampled in segment 1 from 2006 through 2012. | 19 |
| Figure 8. Mean annual catch-per-unit-effort by season of target and non-target species collected using mini fyke nets in segment 1 from 2006 through 2012. | 20 |
| Figure 9. Mean annual catch-per-unit-effort by season of selected non-target species collected using mini fyke nets in segment 1 from 2006 through 2012. | 21 |

LIST OF APPENDICES

| | |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----|
| Appendix A. Phylogenetic list of Missouri River fishes with corresponding letter codes used in the long-term pallid sturgeon and associated fish community sampling program. | 30 |
| Appendix B. Definitions and codes used to classify standard Missouri River habitats in the long term pallid sturgeon and associated fish community sampling program..... | 36 |
| Appendix C. List of standard and wild gears, their corresponding codes in the database, seasons deployed, years used, and catch-per-unit-effort units for collection of Missouri River fishes for the long-term pallid sturgeon and associated fish community sampling program | 37 |
| Appendix D. Stocking locations and codes for pallid sturgeon by Recovery Priority Management Area in the Missouri River Basin..... | 38 |
| Appendix E. Juvenile and adult pallid sturgeon stocking summary for segment 1 of the Missouri River (RPMA 2). | 40 |
| Appendix G. Hatchery names, locations, and abbreviations. | 46 |

Introduction

The U.S. Fish and Wildlife Service (USFWS) listed the pallid sturgeon *Scaphirhynchus albus* as endangered in 1990. In response to the listing, the USFWS issued a Biological Opinion to the U.S. Army Corps of Engineers (COE), the main water management entity responsible for the Missouri River mainstem from Fort Peck Dam and Reservoir to its confluence with the Mississippi River. Additionally, an amendment to the 2000 Biological Opinion was issued in 2003. The Amendment listed several Reasonable and Prudent Alternatives (RPA) to address the inability of pallid sturgeon to naturally reproduce and the need to be able to detect changes in their populations and ecosystem trends.

The Pallid Sturgeon Population Assessment Program (Program) is guided by the RPA's in the 2003 Amendment to the 2000 Biological Opinion. The Program is a comprehensive monitoring plan designed to assess survival, movement, distribution, habitat use, and physical characteristics of these habitats used by wild and hatchery reared juvenile pallid sturgeon (Welker and Drobish 2011). The 2000 Biological Opinion divides the Program area into river and reservoir segments and assigns high, moderate, or low priority management action areas to these segments for pallid sturgeon (Welker and Drobish 2011). The focus of the Program is on the high priority management action segments. The Missouri River from Fort Peck Dam downstream to the headwaters of Lake Sakakawea, ND is listed as a high priority action segment.

The Program has stratified the Missouri River from Fort Peck Dam to the headwaters of Lake Sakakawea into four study segments based on biological, hydrological and fluvial geomorphological characteristics. The COE has contracted Montana Fish, Wildlife & Parks (FWP) to conduct the Program sampling in the Missouri River from Fort Peck Dam downstream to its confluence with the Yellowstone River, which consists of study segments 1 through 3.

This was the seventh field season that Montana Fish, Wildlife & Parks conducted standard Program sampling in Segment 1 of the Missouri River.

The objectives of this program are as follows:

1. Document annual results and long-term trends in pallid sturgeon population abundance and geographic distribution throughout the Missouri River System.
2. Document annual results and long-term trends of habitat use of wild pallid sturgeon and hatchery stocked pallid sturgeon by season and life stage.
3. Document population structure and dynamics of pallid sturgeon in the Missouri River System.
4. Evaluate annual results and long-term trends in native target species population abundance and geographic distribution throughout the Missouri River system.
5. Document annual results and long-term trends of habitat usage of the native target species by season and life stage.
6. Document annual results and long-term trends of all non-target species population abundance and geographic distribution throughout the Missouri River system, where sample size is greater than fifty individuals.

Sampling Season and Species

The Program has two discrete seasons (sturgeon and fish community), which are primarily based on water temperatures. However, the sturgeon season is designed to sample sturgeon with gears that are temperature dependent, such as gill nets. Due to the nature of the majority of habitats in segment 1 through 3, gill nets are not an efficient gear for collecting pallid sturgeon in Montana to debris flows and swift currents and therefore they are not used within Segments 1 through 3. Trammel nets and otter trawl are standard gears used in segments 1-3 during sturgeon season, and so far appear to be an effective way to sample pallid sturgeon. Trotlines are now a standard gear used in segment 1.

The fish community season runs from the beginning of July till the end of October and is designed not only to monitor sturgeon, but also to monitor other native Missouri River fish populations. Both trammel nets and the otter trawl are still used, but to more effectively sample shallow water habitats < 1.2 m in depth, mini fyke nets are also used as a standard gear.

In addition to pallid sturgeon, the Program is designed to monitor nine other native Missouri River species labeled “target” species. These include, shovelnose sturgeon *Scaphirhynchus platyrhynchus*, blue sucker *Cycleptus elongatus*, sauger *Sander canadensis*, sturgeon chub *Macrhybopsis gelida*, sicklefin chub *M. meeki*, speckled chub *M. aestivalis*, plains minnow *Hybognathus placitus*, western silvery minnow *H. argyritis*, and sand shiner

Notropis stramineus. This suite of species was selected for various reasons. First, some species may have similar habitat requirements as pallid sturgeon and therefore by monitoring their populations we may gain further insight into pallid sturgeon habitat and how anthropomorphic and natural changes to the Missouri River affect native fish assemblages. Secondly, it is hypothesized that various chub species and other native fishes are an important component of pallid sturgeon diet. Thirdly, we wouldn't expect to see an immediate response in a long-lived species such as the pallid sturgeon when environmental conditions change from either favorable or detrimental conditions. Thus, by monitoring short-lived native fishes we may be able to correlate environmental conditions to changes in fish populations on a much shorter time interval and make inferences on how pallid sturgeon populations and their habitat are being affected.

Study Area

Segment 1 of the Missouri River begins at Fort Peck Dam and runs downstream to its confluence with the Milk River. This segment constitutes only 6% (11.5 river miles) of the entire 189.5 river miles downstream of Fort Peck Dam to the headwaters of Lake Sakakawea in North Dakota (Welker and Drobish 2011). This reach of the Missouri River is characterized by an unnatural hydrograph, thermograph, sediment dynamics, and fish community due to the influence of Fort Peck Dam, which was constructed in 1940 (Bramblett and White, 2001). Segment 1 includes the Fort Peck Dredge Cuts, a deepened and widened section of river immediately below the dam created by the dredging of earth used to construct the dam. Regulated hypolimnetic water releases from Fort Peck Reservoir have changed a once turbid sandy bottom stretch of river into a cold clear cobble dominated river. Fort Peck Reservoir has substantially reduced suspended sediment loads in the river below Fort Peck Dam when compared to its natural state (Galat et al, 2005).

Peaks in the hydrograph are related to power production and barge traffic downstream, instead of natural spring runoff and precipitation events (Galat et al, 2005). Many species native to this stretch of river such as the pallid sturgeon, sicklefin chub and sturgeon chub find the cold clear water unsuitable and are now common only farther downstream where tributaries have warmed and muddied the waters of the Missouri (Gardner and Stewart, 1987). Fish much more suited for this cold clear water such as rainbow trout *Oncorhynchus mykiss*, brown trout *Salmo trutta* and Chinook salmon *Oncorhynchus tshawytscha* have been stocked on and off from 1950 to 1990. Other nonnative species such as largemouth bass *Micropterus salmoides*, northern pike *Esox lucius*, walleye *Sander vitreus*, and yellow perch *Perca flavescens* have been stocked in the dredge cuts to increase angling opportunities. It is believed that many of these sight-feeding piscivores have out competed the native fishes in this stretch of river (Galat et al, 2005). In summary, this unique stretch of river is now vastly different from the once braided and shifting channels of the “Big Muddy” before Fort Peck Dam (Galat et al, 2005).

Methods

Sampling methods for the Pallid Sturgeon Population Assessment Program were conducted in accordance with the Standard Operating Procedures (Welker and Drobish 2011), which was established by representatives from State and Federal agencies involved with pallid sturgeon recovery on the Missouri River. For a detailed description of methodologies please see Welker and Drobish (2011). A general description of those guidelines follows.

Sampling Site Selection and Description

Montana Fish Wildlife & Parks (FWP) was contracted to sample Segment 1 from Fort Peck Dam (RM 1771.5) to the mouth of the Milk River (RM 1761), Segment 2 from the mouth of the Milk River (RM 1761) to Wolf Point (RM 1701.5) and Segment 3 from Wolf Point (RM 1701.5) to the Montana/North Dakota border (RM 1586.5). Segment 1 consists of one non-random bend at river mile 1766. Segment 1 was selected as a reference study bend to be sampled each year to facilitate comparisons of the most highly altered area of the Missouri River in Recovery Priority Management Area (RPMA) 2 to downstream areas (segments 2 through 4). By comparing data from segment 1 with downstream segments, a better understanding of how Fort Peck Dam influences the fish communities of the Missouri River might be attained.

During 2012 segment 1 was sampled on May 16th during the sturgeon season and October 1st during the fish community season. Four standard gears were used, trammel net, otter trawl, and trotlines were used during both the sturgeon and fish community seasons and mini-fyke nets during the fish community season.

While Missouri River flows were lower in 2012 when compared to 2011 (Figure 2), flows were substantially higher in 2012 when compared to 2006 to 2010. Flows out of the Fort Peck Dam powerhouse were consistently over 10,000 cfs from mid-June through September.

The Population Assessment Team developed a standard set of habitat classifications for the Missouri River (Appendix B) which consists of three distinct macrohabitats found in every bend, a main channel crossover (CHXO), main channel outside bend (OSB), and main

channel inside bend (ISB). Each sampling bend was comprised of these three main macrohabitats. Nine additional macrohabitats were identified that may or may not be present in every bend: large tributary mouths (TRML), small tributary mouths (TRMS), confluence areas (CONF), large and small secondary connected channels (SCCL& SCCS), deranged channels (DRNG), braided channels (BRAD), dendritic channels (DEND) and non-connected secondary channel (SCN). For the reference bend in segment 1, five macrohabitats were sampled, CHXO, OSB, ISB, SCCL and SCCS.

Mesohabitats were established to further define macrohabitats. Mesohabitats include bars (BARS), pools (POOL), channel border (CHNB), thalweg (TLWG) and island tip (ITIP). Channel borders are situated in areas between the deepest portions of the river up to a depth of 1.2 m. Bars are considered shallow areas (< 1.2 m) where terrestrial and aquatic habitats merge. The thalweg is the deepest portion of the river between the two channel borders where the majority of the flow is directed. Pools are directly downstream of any feature that creates scour, thus creating a habitat of deep (> 1.2 m) slower moving water. Island tips are just downstream of bars or islands where two channels meet where the water is > 1.2 m in depth. Two mesohabitats were sampled in segment 1, CHNB and BARS.

For all analysis, the sampling unit was the river bend, where every river bend has a channel crossover, inside and outside bend. The downstream border of a river bend is the beginning of the next downstream bend's channel crossover.

Sampling Gear

For specific information pertaining to the habitats gears are utilized in and physical measurements taken in accordance with sampling the various gears described below please see Welker and Drobish (2011).

Trammel Net

The standard trammel net has a length of 38.1 m, an inner mesh wall 2.4 m and two outer mesh walls 1.8 m deep. The inner mesh is made of #139 multifilament twine with a bar mesh size of 25.4 mm. The outer walls are constructed of #9 multifilament twine with a bar mesh size of 203.2 mm. The float line is a 12.7 mm diameter foam core with a lead line of

22.7 kg. Trammel nets were drifted from the bow of the boat and orientated perpendicular to the river flow for a minimum of 75 m and a maximum drift distance of 300 m.

Otter Trawl

The standard otter trawl has a length of 7.6 m, a width of 4.9 m and height of 0.9 m. The otter trawl has an inner mesh (6.35mm bar, #18 polyethylene twine) and outer mesh (38mmbar, #9 polyethylene twine) and a cod end opening of 406.4 mm. The trawl doors were made from 19.1 mm marine plywood and measured 762 mm x 381 mm. The trawl doors are used to keep the mouth of the trawl open while deployed on the riverbed. The trawl also has a 7.9 m long tickler chain attached to the bottom of the mouth of the trawl, which aids in keeping it orientated on the riverbed and protecting the mouth when snags are encountered. The otter trawl was deployed from the bow of the boat parallel to the current with two 30.5 m ropes and towed downstream slightly faster than current speed for a minimum of 75 m and a maximum distance of 300 m.

Mini-Fyke Nets

The standard mini-fyke net consists of two rectangular frames 1.2 m wide and 0.6 m high and two 0.6 m tempered steel hoops. A 4.5 m long and 0.6 m high lead is connected to the first frame. The fyke net is made of 3 mm “ace” style mesh. The lead has small floats attached to the top and lead weights on the bottom. Mini-fyke nets are set with a “T” stake on shore and extend into river as perpendicular to the shoreline as possible or angled slightly downstream where higher velocities existed. Mini-fyke nets were set overnight and checked the following morning.

Trotlines

Trotlines consisted of 32 m nylon rope attached to both upstream and downstream anchors. Octopus style circle hooks were attached to the ropes using 136 kg monofilament line and commercial fishing clips. Twenty 45.7 cm leaders were used on each trotline. Hooks consisted of 2/0 circle hooks. Each trotline used one hook size and each hook size was used at least once in each macrohabitat sampled. Trotlines were baited with night crawlers, and were set overnight then checked the following morning.

Data Collection and Analysis

A minimum of eight random subsamples with each gear were deployed in the reference bend in segment 1. At least two subsamples (when possible) were taken with each gear in each macro habitat within the bend. More than two subsamples were taken in a macrohabitat for a gear when the number of discrete macrohabitats was less than four or less than four could be effectively sampled.

All fish were measured to the nearest mm. Fork length (FL) was used for sturgeon species, while other species were measured to TL with one exception, paddlefish *Polyodon spathula*, which were measured from the eye to the fork of the caudal fin. The first 25 fish of each species in each subsample were measured, after 25 they were counted.

Time was recorded at the beginning of each sample with all gears and an end time was always recorded when pulling mini-fyke net sets. A global positioning satellite (GPS) position was taken at the beginning and end of all otter and beam trawls and trammel net drifts. One GPS location was taken for mini fyke net sets. All GPS locations were taken using a Garmin GPS 76 unit with Wide Area Augmentation System (WAAS) capability.

Sample depth was determined at the beginning, middle and end of each trawl and drift using a Lowrance X136 sonar unit. One depth was taken for mini-fyke nets at the intersection of the frame and floatline using a wading rod.

Water temperature taken near the surface was recorded at every sample using the Lowrance X136 unit for trawls and trammel net drifts and using a hand held thermometer for mini fyke nets.

Habitat samples were collected randomly for 25% of each mesohabitat within each macrohabitat sampled. Velocities (mps) were taken at three depths in the water column for habitats > 1.2 m in depth (bottom, 0.8 of bottom depth and 0.2 of the bottom depth) using either a Current AA Price Meter and sounding reel or a Marsh-McBirney Flo Mate 2000. Velocities for shallow water habitats (< 1.2 m) were taken at the bottom and 0.6 of the bottom depth using the March-McBirney Flo Mate 2000.

Turbidity was recorded in nephelometric turbidity units (NTU) using a LaMotte 2020 turbidity meter. Turbidity was taken at the midpoint of all samples, except mini-fyke sets, where it was taken at the convergence of the rectangular frame and float line.

In addition to 25% of all mesohabitats, habitat measurements were taken whenever a pallid sturgeon was captured.

Genetic Verification

Genetic verification for pallid sturgeon or potential hybrids followed the methods outlined in Welker and Drobish (2011). Two fin pectoral fin clips ($\sim 2 \text{ cm}^2$) are taken from any pallid sturgeon of unknown origin. Fin samples are then preserved in 95% non-denatured alcohol for genetic analysis. All samples are sent to the U.S. Fish and Wildlife Service's Northeast Fishery Center Conservation Genetics Lab for analysis and archiving.

Analyses

The fundamental sampling unit for the Population Assessment Program is the river bend, where sample size is equal to the number of bends sampled. Since only one river bend in Segment 1 is sampled per year, only one true sample was taken. Therefore, all CPUE data for Segment 1 are the averages of all subsamples and no error is associated with these estimates.




Figure 1. Map of segment 1 of the Missouri River with major tributaries, common landmarks, and historic stocking locations for pallid sturgeon. Segment 1 encompasses the Missouri River from Fort Peck Dam (River Mile 1771.5) to the mouth of the Milk River (River Mile 1760.0).

Missouri River Below Fort Peck Dam




Figure 2. Missouri River discharge downstream of Fort Peck Dam from 2006 to 2012.

Results

Pallid Sturgeon

No pallid sturgeon or pallid sturgeon x shovelnose sturgeon hybrids were collected in Program sampling within Segment 1 of the Missouri River during 2012. This was the seventh year of standardized sampling where no pallid sturgeon were sampled. However, a 1997 year class hatchery reared pallid sturgeon was captured by FWP during their annual fall gill netting effort in the Fort Peck Dredge Cuts. This fish was not tagged and a genetics sample was taken and subsequently sent to the USFWS Lamar Genetics Laboratory, where it was confirmed as a hatchery fish from the 1997 year class. After capture, this fish received a PIT Tag (#48686C7324) and a radio transmitter (frequency 149.760, code 64). This was the first hatchery reared pallid sturgeon captured within Segment 1 by any crew since stocking began in 1998.

Targeted Native River Species

Shovelnose Sturgeon

A total of 83 shovelnose sturgeon were sampled in segment 1 during 2012, 65 and 18 during the sturgeon and fish community seasons, respectively. During the sturgeon season, 29 shovelnose sturgeon were sampled in both trammel nets (Figure 3) and trotlines (Figure 4). Of the 18 shovelnose sturgeon sampled during the fish community season, trammel nets captured eight, trotlines 9 and the otter trawl sampled one (Figure 6).

Trammel net CPUE was higher during both the sturgeon and fish community seasons during 2012, when compared to 2011 (Figure 3). No discernible pattern has been witnessed in regards to trammel net CPUE between the sturgeon and fish community seasons over the past seven years of sampling.

Trotline CPUE of shovelnose was considerably higher in 2012 during the sturgeon season when compared to the fish community season (Figure 4). Trotline CPUE has decreased during the fish community season in each year sampled (Figure 4). Shovelnose sturgeon have been the one of, if not the most abundant species captured using trotlines in Segment 1 during each sampling occasion beginning in 2008.

Few, shovelnose sturgeon have been captured in the otter trawl over the past seven years (Figure 5). This is likely due to the fact that no shovelnose smaller than 450 mm have been captured in any gear (Figure 6).

The size structure of shovelnose sturgeon sampled in segment 1 during both seasons has not changed appreciably over the last seven sampling seasons (Figure 6). Furthermore, no juvenile shovelnose sturgeon have been captured in any gear over the past seven years of sampling. In addition to the size structure remaining very similar between years, the length to weight relationship of shovelnose sturgeon has also not changed significantly since 2006 (Figure 9).

Sturgeon Chub

No sturgeon chubs were collected in Segment 1 during sampling in 2012. Only one sturgeon chub has been collected to date in segment 1 through seven years of sampling and that fish was collected in the otter trawl during 2010 (Figure 5).

Sicklefin Chub

No sicklefin chubs have been collected through seven years of sampling Segment 1.

Sand Shiner

Two sand shiners were captured in 2012, both in mini fyke nets set during the fish community season (Figure 8). This was the third lowest year for sand shiners captured in mini fyke nets within Segment 1. No sand shiners were captured in both 2011 and 2006. The highest CPUE for sand shiners was observed during 2010, with a CPUE of 6.5/net night.

Western Silvery Minnow

No western silvery minnows were collected in 2012 in segment 1. Western silvery minnows have only been collected in two out of the seven years of sampling, with over three fish per net night in 2008 and just over one per net night in 2010 (Figure 8).

Blue Sucker

A total of three adult blue suckers were sampled with trammel nets during 2012, all with trammel nets (Figure 3). Blue suckers have been sampled every year in segment 1, although in very low numbers. Based on aging of blue suckers downstream of Gavins Point Dam, these fish are likely older than age 7 (Labay et al. 2008).

Sauger

Three sauger were sampled in Segment 1 during 2012, all using trammel nets. One sauger was captured during the sturgeon season and two during the fish community season. Overall, sauger averaged 426.3 mm in length and 645 g. Very few sauger have been sampled in segment 1 in the past seven years (Figure 3).

Segment 1 Trammel Net




Figure 3. Trammel net CPUE by season for all target species and non-target species sampled in segment 1 of the Missouri River during sturgeon and fish community season from 2006 through 2012. Target species are indicated by an asterisk. Note the difference in scale of the Y-axes.

Segment 1 Trotline




Figure 4. Trotline CPUE for all target and non-target species sampled in the sturgeon season (top panel) from 2009 through 2012, and the fish community season (bottom panel) from 2008 through 2012 in segment 1 of the Missouri River. Note that trotlines were not set in 2008 during the Sturgeon Season or in 2009 during the Fish Community Season. Shovelnose sturgeon are a target species.

Segment 1 Otter Trawl




Figure 5. Otter Trawl CPUE by season for all target species and non-target species sampled in segment 1 of the Missouri River during sturgeon and fish community season from 2006 through 2012. Target species are indicated by an asterisk. Note the difference in scale of the Y-axes.

Segment 1 Shovelnose Sturgeon




Figure 6. Length frequency histogram for all shovelnose sturgeon sampled in Segment 1 of the Missouri River from 2006 through 2012.

Segment 1 Shovelnose Sturgeon




Figure 7. Weight-length relationship for all shovelnose sturgeon sampled in Segment 1 from 2006 through 2012.

Segment 1 Mini Fyke Net




Figure 8. Mini-fyke net CPUE for target and non-target species sampled during the fish community season in segment 1 of the Missouri River from 2006 through 2012. Target species are indicated by asterisks. Note the differences in Y-axes.

Segment 1 Mini Fyke Net




Figure 9. Mini-fyke net CPUE for non-target species sampled during the fish community season in segment 1 of the Missouri River from 2006 through 2012. Note the differences in Y-axes.

Table 1. Presence absence of all species (by common name) collected in segments 1 through 3 in the Missouri River during 2006 through 2012. Boxes marked with an X indicate at least one specimen was sampled. Species in bold are native target species.

| Segments | | | Segments | | |
|----------------------------------------------|---|---|---------------------------------------------|---|---|
| 1 | 2 | 3 | 1 | 2 | 3 |
| <i>Ascipenseridae - sturgeons</i> | | | <i>Esocidae - pikes</i> | | |
| Pallid sturgeon | X | X | Northern Pike | X | X |
| Shovelnose sturgeon | X | X | <i>Osmeridae - smelts</i> | | |
| <i>Polyodontidae - paddlefishes</i> | | | Rainbow smelt | | X |
| Paddlefish | X | X | <i>Ictaluridae - catfishes</i> | | |
| <i>Hiodontidae - mooneyes</i> | | | Channel catfish | X | X |
| Goldeye | X | X | Black bullhead | X | X |
| <i>Cyprinidae - carps and minnows</i> | | | Yellow bullhead | | X |
| Common Carp | X | X | Stonecat | X | X |
| Flathead chub | X | X | <i>Salmonidae - trouts</i> | | |
| Emerald shiner | X | X | Rainbow trout | X | X |
| Lake chub | X | | Brown trout | X | X |
| Longnose dace | X | X | Lake Trout | X | |
| Northern redbelly dace | X | X | Lake whitefish | X | X |
| Plains minnow | | X | Cisco | X | X |
| Western silvery minnow | X | X | <i>Gadidae - cods</i> | | |
| <i>Brassy minnow</i> | | | Burbot | X | X |
| Sicklefin chub | X | X | <i>Gasterosteidae - sticklebacks</i> | | |
| Sturgeon chub | X | X | Brook stickleback | | X |
| Sand shiner | X | X | <i>Centrarchidae - sunfishes</i> | | |
| Spottail shiner | X | X | Green sunfish | | X |
| Fathead minnow | X | X | Pumkinseed | X | X |
| <i>Catostomidae-suckers</i> | | | White crappie | X | X |

| Segments | | | Segments | | |
|----------------------------------------|---|---|------------------------------------|---|---|
| 1 | 2 | 3 | 1 | 2 | 3 |
| Bigmouth buffalo | X | X | Smallmouth bass | X | |
| Smallmouth buffalo | X | X | <i>Percidae - perches</i> | | |
| Blue sucker | X | X | Iowa darter | | X |
| River carpsucker | X | X | Yellow perch | X | X |
| White sucker | X | X | Sauger | X | X |
| Longnose sucker | X | X | Walleye | X | X |
| Shorthead redhorse | X | X | <i>Sciaenidae - drums</i> | | |
| <i>Moronidae-temperate bass</i> | | | Freshwater drum | X | X |
| White bass | | X | <i>Lepisosteidae - Gars</i> | | |
| | | | Shortnose Gar | | X |

Missouri River Fish Community

A total of 889 fishes representing 17 different species were sampled during the standardized sampling in Segment 1 during 2012. The majority of fish (n=781) were sampled during the fish community season, while the remaining 108 were caught in the sturgeon season. Mini fyke nets caught the most fish (n = 704), followed by trotlines (n = 73), trammel nets (n = 65) and otter trawl (n = 47).

White sucker *Catostomus commersoni* were the most abundant species captured with a total of 691 sampled. Other species sampled included common carp *Cyprinus carpio*, bluegill *Lepomis macrochirus*, channel catfish *Ictalurus punctatus*, emerald shiner *Notropis atherinoides*, fathead minnow *Pimephales promelas*, longnose dace *Rhinichthys cataractae*, longnose sucker *Catostomus catostomus*, rainbow trout *Onchorhynchus mykiss*, river carpsucker *Carpionodes carpio*, smallmouth bass *Micropterus dolomieu*, smallmouth buffalo *Ictiobus bubalus* and spottail shiner *Notropis hudsonius*.

Discussion

Segment 1 of the Missouri River is a highly altered segment due to the proximity of Fort Peck Dam. Fort Peck Dam, a hypolimnetic withdraw structure, is located approximately five river miles upstream of segment 1, which creates cold summer water temperatures and low suspended sediment loads. During the sturgeon season of 2012, water temperatures averaged 8.8 C° and water turbidity averaged 5.3 NTU's. During the fish community season, temperature increased to an average of 13.0 C°, while turbidity increased to 8.6 NTU's. In addition, the benthic substrate of segment 1 is noticeably different than the substrates of downstream segments. Segment 1 is primarily composed of gravel and cobble due to the degrading stream bed, which is at least in part due to the lack of suspended sediments in the water column. Although flows in Segment 1 were lower in 2012 when compared to 2011, river discharge was higher than years prior to 2011 (Figure 2). Flows of 10,000 cfs or greater were observed for the majority of the summer.

No pallid sturgeon have been sampled during standardized sampling in Segment 1 from 2006 through 2012, despite continuous stocking from 2004 to 2008 in the confluence area of the

Milk and Missouri Rivers. However, FWP did capture a 1997 year class pallid sturgeon in the Missouri River Dredge Cuts, which is just upstream of Segment 1. This is the first documentation of hatchery reared pallid sturgeon using the Missouri River upstream of the Milk River. Furthermore, data have indicated that at least some stocked pallid sturgeon from all other stockings sites in segments 2 and 3 do move upstream.

Even with the highly altered conditions of segment 1, many native species are still occupying the habitats of this segment. However, during the past six years of sampling, a total of 10 non-native species have been found in segment 1 including, common carp *Cyprinus carpio*, rainbow trout *Onchorhynchus mykiss*, brown trout *Salmo trutta*, lake trout *Salvelinus namaycush*, lake whitefish *Coregonus clupeaformis*, ciscoe *C. artedi*, spottail shiner *Notropis hudsonius*, smallmouth bass *Micropterus dolomieu*, and yellow perch *Perca flavescens* (Table 1).

The CPUE of shovelnose sturgeon in segment 1 continues to be variable (Figure 3 and 4). Overall trammel net CPUE for shovelnose sturgeon was higher in 2012 than in 2011. In addition, sturgeon season trotline CPUE was higher in 2012 than 2011, but was lower during the fish community season. Nevertheless, since only eight subsamples are taken during each season, catch is variable and little conclusions can be drawn from the CPUE data. More importantly is the consistency in the size distribution of shovelnose in segment 1. No shovelnose under 450 mm have been collected in seven years of sampling, which indicates the population is made up of older fish and little to no juvenile rearing is occurring in the area. Past telemetry studies have also shown that the shovelnose sturgeon population in segment 1 is a resident population of fish that do not migrate far. These fish do not appear to be spawning since we don't collect any black egg females in the spring or early summer, as we do in the lower parts of segment 3. Tagging information has also shown that adult shovelnose in segment 1 either do not grow or grow at a very slow rate. Individuals that have been recaptured after 20 years at large often are the same size as they were when they were tagged.

Adult blue sucker have been captured in segment 1 during all seven sampling years. Similar to shovelnose sturgeon, these have all been large adult fish. Again, CPUE has been variable, but since so few fish have been captured there does not appear to be any differences in their relative abundance over the sampling years.

Mini fyke nets were more effective in 2012 than 2011, but the majority of fish captured were white suckers. Similar to mini fyke nets, the 2012 otter trawl catch higher than 2011, but the catch was comprised mainly of white suckers.

It doesn't appear that the higher releases from Fort Peck Dam in 2011 increased many of the native cyprinids populations that reside within Segment 1. However, we did see a dramatic increase in white suckers in 2012 when compared to 2011. With the spill that occurred over the Fort Peck Dam Spillway in 2011 a lot of predatory game fish such as walleye and northern pike were entrained into the Missouri River downstream of the Dam. This increase in predator abundance was observed in FWP's annual gill netting of the Missouri River Dredge Cuts. This increase in predators could be one reason why we did not observe higher numbers of native cyprinids, since predation was likely high.

Acknowledgments

The U.S. Army Corps of Engineers provided funding for this project. We'd like to thank Tim Welker and George Williams for supporting our work in several ways. Bob Lipscomb, James Collins and Cara Holem-Bell assisted in both the field and shop throughout the year. Steve Dalbey took care of many things while we were on the river. We'd like to thank the Dave Fuller for his assistance. Thanks to Pat Braaten of the U.S. Geological Survey for answering any type of question we may have pertaining to the Missouri River and its fishes. A special thanks to Ryan Wilson, Zack Sandness, Everett Nelson and Steve Krentz of the U.S. Fish and Wildlife Service for all the collaboration between our offices.

References

- Dattilo, J. E., R. R. Dirnberger, P. T. Horner, D. J. Niswonger, M. L. Miller and V. H. Travinchek. 2008a. Three Year Summary Age and Growth Report For Sand Shiner (*Notropis stramineus*). Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Missouri Department of Conservation. Chillicothe, MO.
- Dattilo, J. E., R. R. Dirnberger, P. T. Horner, D. J. Niswonger, M. L. Miller and V. H. Travinchek. 2008b. Three Year Summary Age and Growth Report For Plains Minnow, Western Silvery Minnow, Brassy Minnow (*Hybognathus spp.*). Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Missouri Department of Conservation. Chillicothe, MO.
- Dattilo, J. E., R. R. Dirnberger, P. T. Horner, D. J. Niswonger, M. L. Miller and V. H. Travinchek. 2008c. Three Year Summary Age and Growth Report for Sauger (*Sander canadensis*). Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Missouri Department of Conservation. Chillicothe, MO.
- Galat, D.L., C.R. Berry Jr., E.J. Peters and R.G. White. 2005. Missouri River. Pages 427-480 in A.C. Benke and C.E. Cushing (editors). Rivers of North America, Elsevier, Oxford.
- Gardner, W.M. and P.A. Stewart. 1987. The Fishery of the Lower Missouri River. Federal Aid to Fish and Wildlife Restoration Project FW-2-R Job I-b. Montana Fish, Wildlife and Parks. Helena, Montana.
- Labay, S., J. Kral and S. Stukel. 2008. Three Year Summary Age and Growth Report for Blue Sucker. Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. South Dakota Department of Game, Fish and Parks. Yankton, SD.
- Pierce, C. L., C. S. Guy, P. J. Braaten, and M.A. Pegg. 2004. Fish growth, mortality, recruitment, condition, and size structure. Volume 4. Population structure and habitat use of benthic fishes along the Missouri and lower Yellowstone Rivers. U.S. Geological Survey, Cooperative Research Units, Iowa State University, Ames Iowa.
- Steffensen, K. and M. Hamel. 2008. Four Year Summary Age and Growth Report For Shovelnose Sturgeon. Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Nebraska Game and Parks Commission. Lincoln, NE.
- Welker, T. L., and M. R. Drobish. (editors), 2011. Missouri River Standard Operating Procedures for Fish Sampling and Data Collection, Volume 1.5. U.S. Army Corps of Engineers, Omaha District, Yankton, SD.

APPENDICES

Appendix A. Phylogenetic list of Missouri River fishes with corresponding letter codes used in the long-term pallid sturgeon and associated fish community sampling program. The phylogeny follows that used by the American Fisheries Society, Common and Scientific Names of Fishes from the United States and Canada, 5th edition. Asterisks and bold type denote targeted native Missouri River species.

| Scientific name | Common name | Letter Code |
|--------------------------------------------|-----------------------------|--------------|
| CLASS CEPHALASPIDOMORPHI-LAMPREYS | | |
| ORDER PETROMYZONTIFORMES | | |
| Petromyzontidae – lampreys | | |
| <i>Ichthyomyzon castaneus</i> | Chestnut lamprey | CNLP |
| <i>Ichthyomyzon fossor</i> | Northern brook lamprey | NBLP |
| <i>Ichthyomyzon unicuspis</i> | Silver lamprey | SVLP |
| <i>Ichthyomyzon gagei</i> | Southern brook lamprey | SBLR |
| Petromyzontidae | Unidentified lamprey | ULY |
| Petromyzontidae larvae | Unidentified larval lamprey | LVLP |
| CLASS OSTEICHTHYES – BONY FISHES | | |
| ORDER ACIPENSERIFORMES | | |
| Acipenseridae – sturgeons | | |
| <i>Acipenser fulvescens</i> | Lake sturgeon | LKSG |
| <i>Scaphirhynchus</i> spp. | Unidentified Scaphirhynchus | USG |
| <i>Scaphirhynchus albus</i> | Pallid sturgeon | PDSG* |
| <i>Scaphirhynchus platyrhynchus</i> | Shovelnose sturgeon | SNSG* |
| <i>S. albus</i> X <i>S. platyrhynchus</i> | Pallid-shovelnose hybrid | SNPD |
| Polyodontidae – paddlefishes | | |
| <i>Polyodon spathula</i> | Paddlefish | PDFH |
| ORDER LEPISTOSTEIFORMES | | |
| Lepisosteidae – gars | | |
| <i>Lepisosteus oculatus</i> | Spotted gar | STGR |
| <i>Lepisosteus osseus</i> | Longnose gar | LNGR |
| <i>Lepisosteus platostomus</i> | Shortnose gar | SNGR |
| ORDER AMMIFORMES | | |
| Amiidae – bowfins | | |
| <i>Amia calva</i> | Bowfin | BWFN |
| ORDER OSTEOGLOSSIFORMES | | |
| Hiodontidae – mooneyes | | |
| <i>Hiodon alosoides</i> | Goldeye | GDEY |
| <i>Hiodon tergisus</i> | Mooneye | MNEY |
| ORDER ANGUILLIFORMES | | |
| Anguillidae – freshwater eels | | |
| <i>Anguilla rostrata</i> | American eel | AMEL |

Appendix A. (continued).

| Scientific name | Common name | Letter Code |
|--------------------------------------------------|----------------------------------------|--------------|
| ORDER CLUPEIFORMES | | |
| Clupeidae – herrings | | |
| <i>Alosa alabame</i> | Alabama shad | ALSD |
| <i>Alosa chrysochloris</i> | Skipjack herring | SJHR |
| <i>Alosa pseudoharengus</i> | Alewife | ALWF |
| <i>Dorosoma cepedianum</i> | Gizzard shad | GZSD |
| <i>Dorosoma petenense</i> | Threadfin shad | TFSD |
| <i>D. cepedianum</i> X <i>D. petenense</i> | Gizzard-threadfin shad hybrid | GSTS |
| ORDER CYPRINIFORMES | | |
| Cyprinidae – carps and minnows | | |
| <i>Campostoma anomalum</i> | Central stoneroller | CLSR |
| <i>Campostoma oligolepis</i> | Largescale stoneroller | LSSR |
| <i>Carassus auratus</i> | Goldfish | GDFH |
| <i>Carassus auratus</i> X <i>Cyprinus carpio</i> | Goldfish-Common carp hybrid | GFCC |
| <i>Couesius plumbeus</i> | Lake chub | LKCB |
| <i>Ctenopharyngodon idella</i> | Grass carp | GSCP |
| <i>Cyprinella lutrensis</i> | Red shiner | RDSN |
| <i>Cyprinella spiloptera</i> | Spotfin shiner | SFSN |
| <i>Cyprinus carpio</i> | Common carp | CARP |
| <i>Erimystax x-punctatus</i> | Gravel chub | GVCB |
| <i>Hybognathus argyritis</i> | Western silvery minnow | WSMN* |
| <i>Hybognathus hankinsoni</i> | Brassy minnow | BSMN |
| <i>Hybognathus nuchalis</i> | Mississippi silvery minnow | SVMW |
| <i>Hybognathus placitus</i> | Plains minnow | PNMW* |
| <i>Hybognathus</i> spp. | Unidentified <i>Hybognathus</i> | HBNS* |
| <i>Hypophthalmichthys molitrix</i> | Silver carp | SVCP |
| <i>Hypophthalmichthys nobilis</i> | Bighead carp | BHCP |
| <i>Luxilus chrysocephalus</i> | Striped shiner | SPSN |
| <i>Luxilus cornutus</i> | Common shiner | CMSN |
| <i>Luxilus zonatus</i> | Bleeding shiner | BDSN |
| <i>Lythrurus unbratilis</i> | Western redfin shiner | WRFS |
| <i>Macrhybopsis aestivalis</i> | Speckled chub | SKCB* |
| <i>Macrhybopsis gelida</i> | Sturgeon chub | SGCB* |
| <i>Macrhybopsis meeki</i> | Sicklefin chub | SFCB* |
| <i>Macrhybopsis storeriana</i> | Silver chub | SVCB |
| <i>M. aestivalis</i> X <i>M. gelida</i> | Speckled-Sturgeon chub hybrid | SPST |
| <i>M. gelida</i> X <i>M. meeki</i> | Sturgeon-Sicklefin chub hybrid | SCSC |
| <i>Macrhybopsis</i> spp. | Unidentified chub | UHY |
| <i>Margariscus margarita</i> | Pearl dace | PLDC |
| <i>Mylocheilus caurinus</i> | Peamouth | PEMT |
| <i>Nocomis biguttatus</i> | Hornyhead chub | HHCB |
| <i>Notemigonus crysoleucas</i> | Golden shiner | GDSN |
| <i>Notropis atherinoides</i> | Emerald shiner | ERSN |
| <i>Notropis blennioides</i> | River shiner | RVSN |
| <i>Notropis boops</i> | Bigeye shiner | BESN |
| <i>Notropis burchanani</i> | Ghost shiner | GTSN |
| <i>Notropis dorsalis</i> | Bigmouth shiner | BMSN |
| <i>Notropis greeniei</i> | Wedgespot shiner | WSSN |

Appendix A. (continued).

| Scientific name | Common name | Letter Code |
|---------------------------------------|-------------------------------------|--------------|
| Cyprinidae – carps and minnows | | |
| <i>Notropis heterolepsis</i> | Blacknose shiner | BNSN |
| <i>Notropis hudsonius</i> | Spottail shiner | STSN |
| <i>Notropis nubilus</i> | Ozark minnow | OZMW |
| <i>Notropis rubellus</i> | Rosyface shiner | RYSN |
| <i>Notropis shumardi</i> | Silverband shiner | SBSN |
| <i>Notropis stilbius</i> | Silverstripe shiner | SSPS |
| <i>Notropis stramineus</i> | Sand shiner | SNSN* |
| <i>Notropis topeka</i> | Topeka shiner | TPSN |
| <i>Notropis volucellus</i> | Mimic shiner | MMSN |
| <i>Notropis wickliffi</i> | Channel shiner | CNSN |
| <i>Notropis</i> spp. | Unidentified shiner | UNO |
| <i>Opsopoeodus emiliae</i> | Pugnose minnow | PNMW |
| <i>Phenacobius mirabilis</i> | Suckermouth minnow | SMMW |
| <i>Phoxinus eos</i> | Northern redbelly dace | NRBD |
| <i>Phoxinus erythrogaster</i> | Southern redbelly dace | SRBD |
| <i>Phoxinus neogaeus</i> | Finescale dace | FSDC |
| <i>Pimephales notatus</i> | Bluntnose minnow | BNMW |
| <i>Pimephales promelas</i> | Fathead minnow | FHMW |
| <i>Pimephales vigilas</i> | Bullhead minnow | BHMW |
| <i>Platygobio gracilis</i> | Flathead chub | FHCB |
| <i>P. gracilis</i> X <i>M. meeki</i> | Flathead-sicklefin chub hybrid | FCSC |
| <i>Rhinichthys atratulus</i> | Blacknose dace | BNDC |
| <i>Rhinichthys cataractae</i> | Longnose dace | LNDC |
| <i>Richardsonius balteatus</i> | Redside shiner | RDSS |
| <i>Scardinius erythrophthalmus</i> | Rudd | RUDD |
| <i>Semotilus atromaculatus</i> | Creek chub | CKCB |
| | Unidentified Cyprinidae | UCY |
| | Unidentified Asian Carp | UAC |
| Catostomidae - suckers | | |
| <i>Carpionodes carpio</i> | River carpsucker | RVCS |
| <i>Carpionodes cyprinus</i> | Quillback | QLBK |
| <i>Carpionodes velifer</i> | Highfin carpsucker | HFCS |
| <i>Carpionodes</i> spp. | Unidentified Carpiodes | UCS |
| <i>Catostomus catostomus</i> | Longnose sucker | LNSK |
| <i>Catostomus commersoni</i> | White sucker | WTSK |
| <i>Catostomus platyrhincus</i> | Mountain sucker | MTSK |
| <i>Catostomus</i> spp. | Unidentified <i>Catostomus</i> spp. | UCA |
| <i>Cycleptus elongates</i> | Blue sucker | BUSK* |
| <i>Hypentelium nigricans</i> | Northern hog sucker | NHSC |
| <i>Ictiobus bubalus</i> | Smallmouth buffalo | SMBF |
| <i>Ictiobus cyprinellus</i> | Bigmouth buffalo | BMBF |
| <i>Ictiobus niger</i> | Black buffalo | BKBF |
| <i>Ictiobus</i> spp. | Unidentified buffalo | UBF |
| <i>Minytrema melanops</i> | Spotted sucker | SPSK |
| <i>Moxostoma anisurum</i> | Silver redhorse | SVRH |
| <i>Moxostoma carinatum</i> | River redhorse | RVRH |
| <i>Moxostoma duquesnei</i> | Black redhorse | BKRH |
| <i>Moxostoma erythrurum</i> | Golden redhorse | GDRH |
| <i>Moxostoma macrolepidotum</i> | Shorthead redhorse | SHRH |
| <i>Moxostoma</i> spp. | Unidentified redhorse | URH |

Appendix A. (continued).

| Scientific name | Common name | Letter Code |
|------------------------------------------|------------------------------------|-------------|
| Catostomidae - suckers | Unidentified Catostomidae | UCT |
| ORDER SILURIFORMES | | |
| Ictaluridae – bullhead catfishes | | |
| <i>Ameiurus melas</i> | Black bullhead | BKBH |
| <i>Ameiurus natalis</i> | Yellow bullhead | YLBH |
| <i>Ameiurusnebulosus</i> | Brown bullhead | BRBH |
| <i>Ameiurus</i> spp. | Unidentified bullhead | UBH |
| <i>Ictalurus furcatus</i> | Blue catfish | BLCF |
| <i>Ictalurus punctatus</i> | Channel catfish | CNCF |
| <i>I. furcatus</i> X <i>I. punctatus</i> | Blue-channel catfish hybrid | BCCC |
| <i>Ictalurus</i> spp. | Unidentified <i>Ictalurus</i> spp. | UCF |
| <i>Noturus exilis</i> | Slender madtom | SDMT |
| <i>Noturus flavus</i> | Stonecat | STCT |
| <i>Noturus gyrinus</i> | Tadpole madtom | TPMT |
| <i>Noturus nocturnes</i> | Freckled madtom | FKMT |
| <i>Pylodictis olivaris</i> | Flathead catfish | FHCF |
| ORDER SALMONIFORMES | | |
| Esocidae - pikes | | |
| <i>Esox americanus vermiculatus</i> | Grass pickerel | GSPK |
| <i>Esox lucius</i> | Northern pike | NTPK |
| <i>Esox masquinongy</i> | Muskellunge | MSKG |
| <i>E. lucius</i> X <i>E. masquinongy</i> | Tiger Muskellunge | TGMG |
| Umbridae - mudminnows | | |
| <i>Umbra limi</i> | Central mudminnow | MDMN |
| Osmeridae - smelts | | |
| <i>Osmerus mordax</i> | Rainbow smelt | RBST |
| Salmonidae - trouts | | |
| <i>Coregonus artedii</i> | Lake herring or cisco | CSCO |
| <i>Coregonus clupeaformis</i> | Lake whitefish | LKWF |
| <i>Oncorhynchus aguabonita</i> | Golden trout | GDTT |
| <i>Oncorhynchus clarki</i> | Cutthroat trout | CTTT |
| <i>Oncorhynchus kisutch</i> | Coho salmon | CHSM |
| <i>Oncorhynchus mykiss</i> | Rainbow trout | RBTT |
| <i>Oncorhynchus nerka</i> | Sockeye salmon | SESM |
| <i>Oncorhynchus tshawytscha</i> | Chinook salmon | CNSM |
| <i>Prosopium cylindraceum</i> | Bonniville cisco | BVSC |
| <i>Prosopium williamsoni</i> | Mountain whitefish | MTWF |
| <i>Salmo trutta</i> | Brown trout | BNTT |
| <i>Salvelinus fontinalis</i> | Brook trout | BKTT |
| <i>Salvelinus namaycush</i> | Lake trout | LKTT |
| <i>Thymallus arcticus</i> | Arctic grayling | AMGL |

Appendix A. (continued).

| Scientific name | Common name | Letter Code |
|------------------------------------------|-------------------------------|-------------|
| ORDER PERCOPSIFORMES | | |
| Percopsidae – trout-perches | | |
| <i>Percopsis omiscomaycus</i> | Trout-perch | TTPH |
| ORDER GADIFORMES | | |
| Gadidae - cods | | |
| <i>Lota lota</i> | Burbot | BRBT |
| ORDER ATHERINIFORMES | | |
| Cyprinodontidae - killifishes | | |
| <i>Fundulus catenatus</i> | Northern studfish | NTSF |
| <i>Fundulus daphanus</i> | Banded killifish | BDKF |
| <i>Fundulus notatus</i> | Blackstripe topminnow | BSTM |
| <i>Fundulus olivaceus</i> | Blackspotted topminnow | BPTM |
| <i>Fundulus sciadicus</i> | Plains topminnow | PTMW |
| <i>Fundulus zebrinus</i> | Plains killifish | PKLF |
| Poeciliidae - livebearers | | |
| <i>Gambusia affinis</i> | Western mosquitofish | MQTF |
| Atherinidae - silversides | | |
| <i>Labidesthes sicculus</i> | Brook silverside | BKSS |
| ORDER GASTEROSTEIFORMES | | |
| Gasterosteidae - sticklebacks | | |
| <i>Culea inconstans</i> | Brook stickleback | BKSB |
| ORDER SCORPAENIFORMES | | |
| Cottidae - sculpins | | |
| <i>Cottus bairdi</i> | Mottled sculpin | MDSP |
| <i>Cottus carolinae</i> | Banded sculpin | BDSP |
| ORDER PERCIFORMES | | |
| Percichthyidae – temperate basses | | |
| <i>Morone Americana</i> | White perch | WTPH |
| <i>Morone chrysops</i> | White bass | WTBS |
| <i>Morone mississippiensis</i> | Yellow bass | YWBS |
| <i>Morone saxatilis</i> | Striped bass | SDBS |
| <i>M. saxatilis X M. chrysops</i> | Striped-white bass hybrid | SBWB |
| Centrarchidae - sunfishes | | |
| <i>Ambloplites rupestris</i> | Rock bass | RKBS |
| <i>Archoplites interruptus</i> | Sacramento perch | SOPH |
| <i>Lepomis cyanellus</i> | Green sunfish | GNSF |
| <i>Lepomis gibbosus</i> | Pumpkinseed | PNSD |
| <i>Lepomis gulosus</i> | Warmouth | WRMH |
| <i>Lepomis humilis</i> | Orangespotted sunfish | OSSF |
| <i>Lepomis macrochirus</i> | Bluegill | BLGL |
| <i>Lepomis magalotis</i> | Longear sunfish | LESF |
| <i>Lepomis microlophus</i> | Redear sunfish | RESF |
| <i>L. cyanellus X L. macrochirus</i> | Green sunfish-bluegill hybrid | GSBG |

Appendix A. (continued).

| Scientific name | Common name | Letter Code |
|------------------------------------------------|-----------------------------------------------------------------|--------------|
| Centrarchidae - sunfishes | | |
| <i>L. cyanellus</i> X <i>L. humilis</i> | Green-orangespotted sunfish hybrid | GSOS |
| <i>L. macrochirus</i> X <i>L. microlophus</i> | Bluegill-redear sunfish hybrid | BGRE |
| <i>Lepomis</i> spp. | Unidentified <i>Lepomis</i> | ULP |
| <i>Micropterus dolomieu</i> | Smallmouth bass | SMBS |
| <i>Micropterus punctatus</i> | Spotted sunfish | STBS |
| <i>Micropterus salmoides</i> | Largemouth bass | LMBS |
| <i>Micropterus</i> spp. | Unidentified <i>Micropterus</i> spp. | UMC |
| <i>Pomoxis annularis</i> | White crappie | WTCP |
| <i>Pomoxis nigromaculatus</i> | Black crappie | BKCP |
| <i>Pomoxis</i> spp. | Unidentified crappie | UCP |
| <i>P. annularis</i> X <i>P. nigromaculatus</i> | White-black crappie hybrid | WCBC |
| Centrarchidae | Unidentified centrarchid | UCN |
| Percidae - perches | | |
| <i>Ammocrypta asprella</i> | Crystal darter | CLDR |
| <i>Etheostoma blennioides</i> | Greenside darter | GS DR |
| <i>Etheostoma caeruleum</i> | Rainbow darter | RBDR |
| <i>Etheostoma exile</i> | Iowa darter | IODR |
| <i>Etheostoma flabellare</i> | Fantail darter | FTDR |
| <i>Etheostoma gracile</i> | Slough darter | SLDR |
| <i>Etheostoma microperca</i> | Least darter | LTDR |
| <i>Etheostoma nigrum</i> | Johnny darter | JYDR |
| <i>Etheostoma punctulatum</i> | Stippled darter | STPD |
| <i>Etheostoma spectabile</i> | Orangethroated darter | OTDR |
| <i>Etheostoma tetrazonum</i> | Missouri saddled darter | MSDR |
| <i>Etheostoma zonale</i> | Banded darter | BDDR |
| <i>Etheostoma</i> spp. | Unidentified <i>Etheostoma</i> spp. | UET |
| <i>Perca flavescens</i> | Yellow perch | YWPH |
| <i>Percina caproides</i> | Logperch | LGPH |
| <i>Percina cymatotaenia</i> | Bluestripe darter | BTDR |
| <i>Percina evides</i> | Gilt darter | GLDR |
| <i>Percina maculate</i> | Blackside darter | BSDR |
| <i>Percina phoxocephala</i> | Slenderhead darter | SHDR |
| <i>Percina shumardi</i> | River darter | RRDR |
| <i>Percina</i> spp. | Unidentified <i>Percina</i> spp. | UPN |
| | Unidentified darter | UDR |
| <i>Sander canadense</i> | Sauger | SGER* |
| <i>Sander vitreus</i> | Walleye | WLEY |
| <i>S. canadense</i> X <i>S. vitreus</i> | Sauger-walley hybrid/Saugeye | SGWE |
| <i>Sander</i> spp. | Unidentified <i>Sander</i> (formerly <i>Stizostedion</i>) spp. | UST |
| | Unidentified Percidae | UPC |
| Sciaenidae - drums | | |
| <i>Aplodinotus grunniens</i> | Freshwater drum | FWDM |
| NON-TAXONOMIC CATEGORIES | | |
| | Age-0/Young-of-year fish | YOYF |
| | Lab fish for identification | LAB |
| | No fish caught | NFSH |
| | Unidentified larval fish | LVFS |
| | Unidentified | UNID |
| | Net Malfunction (Did Not Fish) | NDNF |

Appendix B. Definitions and codes used to classify standard Missouri River habitats in the long-term pallid sturgeon and associated fish community sampling program. Three habitat scales were used in the hierarchical habitat classification system: Macrohabitats, Mesohabitats, and Microhabitats.

| Habitat | Scale | Definition | Code |
|-----------------------------------|-------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------|
| Braided channel | Macro | An area of the river that contains multiple smaller channels and is lacking a readily identifiable main channel (typically associated with unchannelized sections) | BRAD |
| Main channel cross over | Macro | The inflection point of the thalweg where the thalweg crosses from one concave side of the river to the other concave side of the river, (i.e., transition zone from one-bend to the next bend). The upstream CHXO for a respective bend is the one sampled. | CHXO |
| Tributary confluence | Macro | Area immediately downstream, extending up to one bend in length, from a junction of a large tributary and the main river where this tributary has influence on the physical features of the main river | CONF |
| Dendric | Macro | An area of the river where the river transitions from meandering or braided channel to more of a treelike pattern with multiple channels (typically associated with unchannelized sections) | DEND |
| Deranged | Macro | An area of the river where the river transitions from a series of multiple channels into a meandering or braided channel (typically associated with unchannelized sections) | DRNG |
| Main channel inside bend | Macro | The convex side of a river bend | ISB |
| Main channel outside bend | Macro | The concave side of a river bend | OSB |
| Secondary channel-connected large | Macro | A side channel, open on upstream and downstream ends, with less flow than the main channel, large indicates this habitat can be sampled with trammel nets and trawls based on width and/or depths > 1.2 m | SCCL |
| Secondary channel-connected small | Macro | A side channel, open on upstream and downstream ends, with less flow than the main channel, small indicates this habitat cannot be sampled with trammel nets and trawls based on width and/or on depths < 1.2 m | SCCS |
| Secondary channel-non-connected | Macro | A side channel that is blocked at one end | SCCN |
| Tributary | Macro | Any river or stream flowing in the Missouri River | TRIB |
| Tributary large mouth | Macro | Mouth of entering tributary whose mean annual discharge is > 20 m ³ /s, and the sample area extends 300 m into the tributary | TRML |
| Tributary small mouth | Macro | Mouth of entering tributary whose mean annual discharge is < 20 m ³ /s, mouth width is > 6 m wide and the sample area extends 300 m into the tributary | TRMS |
| Wild | Macro | All habitats not covered in the previous habitat descriptions | WILD |
| Bars | Meso | Sandbar or shallow bank-line areas with depth < 1.2 m | BARS |
| Pools | Meso | Areas immediately downstream from sandbars, dikes, snags, or other obstructions with a formed scour hole > 1.2 m | POOL |
| Channel border | Meso | Area in the channelized river between the toe and the thalweg, area in the unchannelized river between the toe and the maximum depth | CHNB |
| Dam Tailwaters | Meso | Area below dam | DTWT |
| Thalweg | Meso | Main channel between the channel borders conveying the majority of the flow | TLWG |
| Island tip | Meso | Area immediately downstream of a bar or island where two channels converge with water depths > 1.2 m | ITIP |

Appendix C. List of standard and wild gears (type), their corresponding codes in the database, seasons deployed (Fall-Spring, Summer, or all), years used, and catch-per-unit-effort units for collection of Missouri River fishes in segment 1 for the long-term pallid sturgeon and associated fish community sampling program. Long-term monitoring began in 2006 for segment 1.

| Gear | Code | Type | Season | Years | CPUE units |
|---------------------------------------------|-------|--------------|------------|----------------|---------------------|
| Gillnet – 4 meshes, small mesh set upstream | GN14 | Wild | Sturgeon | NOT USED | fish/net night |
| Gillnet – 4 meshes, large mesh set upstream | GN41 | Wild | Sturgeon | NOT USED | fish/net night |
| Gillnet – 8 meshes, small mesh set upstream | GN18 | Wild | Sturgeon | NOT USED | fish/net night |
| Gillnet – 8 meshes, large mesh set upstream | GN81 | Wild | Sturgeon | NOT USED | fish/net night |
| Mini-fyke net | MF | Standard | Fish Comm. | 2006 - Present | fish/net night |
| Push Trawl – 8 ft 4mm x 4mm | POT02 | Evaluation | Fish Comm. | 2006 - 2008 | fish/ 100 m trawled |
| Trammel net – 1 inch inner mesh | TN | Standard | All | 2006 - Present | fish/100 m drift |
| Trot Line – Circle hooks** | TLC1 | Experimental | Sturgeon | 2007 - 2009 | fish/hook night |
| Trot Line – Circle hooks** | TLC1 | Standard | All | 2010-present | fish/hook night |
| Trot Line – Octopus hooks** | TLO_ | Wild | Sturgeon | NOT USED | fish/hook night |
| Trot Line – O'Shaughnessy hooks** | TLS_ | Wild | Sturgeon | NOT USED | fish/hook night |
| Otter trawl – 16 ft head rope | OT16 | Standard | All | 2006 - Present | fish/100 m trawled |
| Otter trawl – 16 ft SKT 4mm x 4mm HB2 MOR | OT01 | Wild | Fish Comm. | NOT USED | fish/100 m trawled |

** Code ends with line length in feet (1 = 105 ft, 2 = 205 ft, 3 = 305 ft, 4 = 405 ft). Hooks are placed between 5 and 10 feet apart.

Appendix D. Stocking locations and codes by Recovery Priority Management Area (RPMA) in the Missouri River Basin.

| State(s) | RPMA | Site Name | Code | River | RM |
|----------|------|----------------------|------|-------------|--------|
| MT | 2 | Forsyth | FOR | Yellowstone | 253.2 |
| MT | 2 | Cartersville | CAR | Yellowstone | 235.3 |
| MT | 2 | Miles City | MIC | Yellowstone | 181.8 |
| MT | 2 | Fallon | FAL | Yellowstone | 124 |
| MT | 2 | Intake | INT | Yellowstone | 70 |
| MT | 2 | Sidney | SID | Yellowstone | 31 |
| MT | 2 | Big Sky Bend | BSB | Yellowstone | 17 |
| ND | 2 | Fairview | FRV | Yellowstone | 9 |
| MT | 2 | Milk River | MLK | Milk | 11.5 |
| MT | 2 | Mouth of Milk | MOM | Missouri | 1761.5 |
| MT | 2 | Grand Champs | GRC | Missouri | 1741 |
| MT | 2 | Wolf Point | WFP | Missouri | 1701.5 |
| MT | 2 | Poplar | POP | Missouri | 1649.5 |
| MT | 2 | Brockton | BRK | Missouri | 1678 |
| MT | 2 | Culbertson | CBS | Missouri | 1621 |
| MT | 2 | Nohly Bridge | NOB | Missouri | 1590 |
| ND | 2 | Confluence | CON | Missouri | 1581.5 |
| SD/NE | 3 | Sunshine Bottom | SUN | Missouri | 866.2 |
| SD/NE | 3 | Verdel Boat Ramp | VER | Missouri | 855 |
| SD/NE | 3 | Standing Bear Bridge | STB | Missouri | 845 |
| SD/NE | 3 | Running Water | RNW | Missouri | 840.1 |
| SD/NE | 4 | St. Helena | STH | Missouri | 799 |
| SD/NE | 4 | Mullberry Bend | MUL | Missouri | 775 |
| NE/IA | 4 | Ponca State Park | PSP | Missouri | 753 |
| NE/IA | 4 | Sioux City | SIO | Missouri | 732.6 |
| NE/IA | 4 | Sloan | SLN | Missouri | 709 |
| NE/IA | 4 | Decatur | DCT | Missouri | 691 |
| NE/IA | 4 | Boyer Chute | BYC | Missouri | 637.4 |
| NE/IA | 4 | Bellevue | BEL | Missouri | 601.4 |
| NE/IA | 4 | Rulo | RLO | Missouri | 497.9 |
| NE/MO/KS | 4 | Kansas River | KSR | Missouri | 367.5 |
| NE | 4 | Platte River | PLR | Platte | 5 |
| KA/MO | 4 | Leavenworth | LVW | Missouri | 397 |
| MO | 4 | Parkville | PKV | Missouri | 377.5 |
| MO | 4 | Kansas City | KAC | Missouri | 342 |

| State(s) | RPMA | Site Name | Code | River | RM |
|----------|------|----------------|------|----------|-------|
| MO | 4 | Miami | MIA | Missouri | 262.8 |
| MO | 4 | Grand River | GDR | Missouri | 250 |
| MO | 4 | Boonville | BOO | Missouri | 195.1 |
| MO | 4 | Overton | OVT | Missouri | 185.1 |
| MO | 4 | Hartsburg | HAR | Missouri | 160 |
| MO | 4 | Jefferson City | JEF | Missouri | 143.9 |
| MO | 4 | Mokane | MOK | Missouri | 124.7 |
| MO | 4 | Hermann | HER | Missouri | 97.6 |
| MO | 4 | Washington | WAS | Missouri | 68.5 |
| MO | 4 | St. Charles | STC | Missouri | 28.5 |

Appendix E. Juvenile pallid sturgeon stocking summary for Segment 2 of the Missouri River (RPMA 2)

| Year | Stocking Site | Number Stocked | Year Class | Stock Date | Age at Stocking ^a | Primary Mark | Secondary Mark |
|------|---------------|----------------|------------|------------|------------------------------|--------------|----------------|
| 1998 | Big Sky Bend | 255 | 1997 | 8/11/1998 | Yearling | PIT Tag | Elastomer |
| 1998 | Confluence | 40 | 1997 | 8/11/1998 | Yearling | PIT Tag | Elastomer |
| 1998 | Nohly Bridge | 255 | 1997 | 8/11/1998 | Yearling | PIT Tag | Elastomer |
| 1998 | Sidney | 230 | 1997 | 8/11/1998 | Yearling | PIT Tag | Elastomer |
| 2000 | Culbertson | 34 | 1998 | 10/11/2000 | 2 yr Old | PIT Tag | |
| 2000 | Fairview | 66 | 1998 | 10/11/2000 | 2 yr Old | PIT Tag | |
| 2000 | Sidney | 66 | 1998 | 10/11/2000 | 2 yr Old | PIT Tag | |
| 2000 | Wolf Point | 34 | 1998 | 10/11/2000 | 2 yr Old | PIT Tag | |
| 2000 | Culbertson | 89 | 1999 | 10/17/2000 | Yearling | PIT Tag | |
| 2000 | Fairview | 150 | 1999 | 10/17/2000 | Yearling | PIT Tag | |
| 2000 | Sidney | 149 | 1999 | 10/17/2000 | Yearling | PIT Tag | |
| 2000 | Wolf Point | 90 | 1999 | 10/17/2000 | Yearling | PIT Tag | |
| 2002 | Culbertson | 270 | 2001 | 7/18/2002 | Yearling | CWT | Elastomer |
| 2002 | Fairview | 270 | 2001 | 7/18/2002 | Yearling | CWT | Elastomer |
| 2002 | Intake | 199 | 2001 | 7/18/2002 | Yearling | CWT | Elastomer |
| 2002 | Sidney | 271 | 2001 | 7/18/2002 | Yearling | CWT | Elastomer |
| 2002 | Wolf Point | 269 | 2001 | 7/18/2002 | Yearling | CWT | Elastomer |
| 2002 | Culbertson | 317 | 2001 | 7/26/2002 | Yearling | PIT Tag | |
| 2002 | Fairview | 360 | 2001 | 7/26/2002 | Yearling | PIT Tag | |
| 2002 | Intake | 97 | 2001 | 7/26/2002 | Yearling | PIT Tag | |
| 2002 | Sidney | 427 | 2001 | 7/26/2002 | Yearling | PIT Tag | |
| 2002 | Wolf Point | 425 | 2001 | 7/26/2002 | Yearling | PIT Tag | |
| 2002 | Intake | 155 | 2001 | 9/18/2002 | Yearling | PIT Tag | |
| 2003 | Culbertson | 1033 | 2002 | 8/7/2003 | Yearling | PIT Tag | Elastomer |
| 2003 | Fairview | 887 | 2002 | 8/7/2003 | Yearling | PIT Tag | Elastomer |
| 2003 | Intake | 1040 | 2002 | 8/7/2003 | Yearling | PIT Tag | Elastomer |
| 2003 | Wolf Point | 926 | 2002 | 8/7/2003 | Yearling | PIT Tag | Elastomer |
| 2004 | Milk River | 821 | 2003 | 4/13/2004 | Yearling | Elastomer | |
| 2004 | Culbertson | 523 | 2003 | 8/9/2004 | Yearling | PIT Tag | Elastomer |

| Year | Stocking Site | Number Stocked | Year Class | Stock Date | Age at Stocking ^a | Primary Mark | Secondary Mark |
|------|---------------|----------------|------------|------------|------------------------------|--------------|----------------|
| 2004 | Intake | 347 | 2003 | 8/9/2004 | Yearling | PIT Tag | Elasomer |
| 2004 | Sidney | 397 | 2003 | 8/9/2004 | Yearling | PIT Tag | Elastomer |
| 2004 | Wolf Point | 379 | 2003 | 8/9/2004 | Yearling | PIT Tag | Elastomer |
| 2004 | Larval Drift | 30000 | 2004 | 7/2/2004 | Fry | | |
| 2004 | Larval Drift | 50000 | 2004 | 7/8/2004 | Fry | | |
| 2004 | Larval Drift | 25000 | 2004 | 7/20/2004 | Fry | | |
| 2004 | Larval Drift | 25000 | 2004 | 7/23/2004 | Fry | | |
| 2004 | Larval Drift | 25000 | 2004 | 7/27/2004 | Fry | | |
| 2004 | Culbertson | 3819 | 2004 | 9/10/2004 | Fingerling | CWT | Elastomer |
| 2004 | Sidney | 2991 | 2004 | 9/10/2004 | Fingerling | CWT | Elastomer |
| 2004 | Wolf Point | 4040 | 2004 | 9/10/2004 | Fingerling | CWT | Elastomer |
| 2004 | Mouth of Milk | 3482 | 2004 | 10/15/2004 | Advanced Fingerling | CWT | Elastomer |
| 2004 | Intake | 2477 | 2004 | 11/18/2004 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Culbertson | 288 | 2004 | 4/12/2005 | Yearling | CWT | Elastomer |
| 2005 | Intake | 309 | 2004 | 4/12/2005 | Yearling | CWT | Elastomer |
| 2005 | Wolf Point | 271 | 2004 | 4/12/2005 | Yearling | CWT | Elastomer |
| 2005 | Intake | 175 | 2004 | 8/19/2005 | Yearling | PIT Tag | Elastomer |
| 2005 | Brockton | 229 | 2005 | 10/5/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Culbertson | 226 | 2005 | 10/5/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Intake | 456 | 2005 | 10/5/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Milk River | 232 | 2005 | 10/5/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Sidney | 122 | 2005 | 10/5/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Wolf Point | 611 | 2005 | 10/12/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Brockton | 371 | 2005 | 10/13/2005 | Advanced | | |
| 2005 | Culbertson | 1736 | 2005 | 10/13/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Culbertson | 182 | 2005 | 10/13/2005 | Advanced Fingerling | | |
| 2005 | Intake | 313 | 2005 | 10/13/2005 | Advanced Fingerling | | |
| 2005 | Milk River | 845 | 2005 | 10/13/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Mouth of Milk | 371 | 2005 | 10/13/2005 | Advanced Fingerling | | |
| 2005 | Sidney | 105 | 2005 | 10/13/2005 | Advanced Fingerling | | |
| 2005 | Wolf Point | 1521 | 2005 | 10/13/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Wolf Point | 371 | 2005 | 10/13/2005 | Advanced Fingerling | | |

| Year | Stocking Site | Number Stocked | Year Class | Stock Date | Age at Stocking ^a | Primary Mark | Secondary Mark |
|------|---------------|----------------|------------|------------|------------------------------|--------------|----------------|
| 2005 | Culbertson | 651 | 2005 | 10/19/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Intake | 2120 | 2005 | 10/19/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Milk River | 485 | 2005 | 10/19/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Sidney | 882 | 2005 | 10/19/2005 | Advanced Fingerling | CWT | Elastomer |
| 2005 | Wolf Point | 650 | 2005 | 10/19/2005 | Advanced Fingerling | CWT | Elastomer |
| 2006 | Culbertson | 235 | 2005 | 3/28/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Intake | 327 | 2005 | 3/28/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Mouth of Milk | 134 | 2005 | 3/28/2006 | Advanced fingerling | Elastomer | |
| 2006 | Sidney | 113 | 2005 | 3/28/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Wolf Point | 232 | 2005 | 3/28/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Intake | 970 | 2005 | 4/3/2006 | Yearling | PIT Tag | Elastomer |
| 2006 | Sidney | 314 | 2005 | 4/3/2006 | Yearling | PIT Tag | Elastomer |
| 2006 | Culbertson | 844 | 2005 | 4/5/2006 | Yearling | PIT Tag | Elastomer |
| 2006 | Mouth of Milk | 1007 | 2005 | 4/5/2006 | Yearling | PIT Tag | Elastomer |
| 2006 | Wolf Point | 866 | 2005 | 4/5/2006 | Yearling | PIT Tag | Elastomer |
| 2006 | Culbertson | 669 | 2005 | 5/1/2006 | Yearling | PIT Tag | Scute Removed |
| 2006 | Intake | 765 | 2005 | 5/1/2006 | Yearling | PIT Tag | Scute Removed |
| 2006 | Mouth of Milk | 650 | 2005 | 5/1/2006 | Yearling | PIT Tag | Scute Removed |
| 2006 | Sidney | 228 | 2005 | 5/1/2006 | Yearling | PIT Tag | Scute Removed |
| 2006 | Wolf Point | 653 | 2005 | 5/1/2006 | Yearling | PIT Tag | Scute Removed |
| 2006 | | 1355 | 2005 | 5/1/2006 | Yearling | PIT Tag | Scute Removed |
| 2006 | Culbertson | 1544 | 2006 | 10/24/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Intake | 1680 | 2006 | 10/24/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Mouth Milk | 1117 | 2006 | 10/24/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Sidney | 586 | 2006 | 10/24/2006 | Advanced Fingerling | Elastomer | |
| 2006 | Wolf Point | 1553 | 2006 | 10/24/2006 | Advanced Fingerling | Elastomer | |
| 2006 | School Trust | 436 | 2006 | 11/8/2006 | Advanced Fingerling | Elastomer | |
| 2007 | Culbertson | 651 | 2006 | 4/5/2007 | Yearling | PIT Tag | Scute Removed |
| 2007 | Fallon | 491 | 2006 | 4/3/2007 | Yearling | PIT Tag | Scute Removed |
| 2007 | Forsyth | 492 | 2006 | 4/3/2007 | Yearling | PIT Tag | Scute Removed |
| 2007 | Sidney | 983 | 2006 | 4/3/2007 | Yearling | PIT Tag | Scute Removed |
| 2007 | School Trust | 639 | 2006 | 4/5/2007 | Yearling | PIT Tag | Scute Removed |

| Year | Stocking Site | Number Stocked | Year Class | Stock Date | Age at Stocking ^a | Primary Mark | Secondary Mark |
|------|---------------|----------------|------------|------------|------------------------------|--------------|----------------|
| 2007 | Wolf Point | 651 | 2006 | 4/5/2007 | Yearling | PIT Tag | Scute Removed |
| 2007 | Wolf Point | 428285 | 2007 | 7/9/2007 | Fry | | |
| 2007 | Grand Champs | 5558 | 2007 | 7/13/2007 | Fry | | |
| 2007 | Miles City | 13125 | 2007 | 7/18/2007 | Fry | | |
| 2007 | Intake | 20763 | 2007 | 8/9/2007 | Fry | | |
| 2007 | Miles City | 13675 | 2007 | 8/9/2007 | Fry | | |
| 2007 | Intake | 336 | 2007 | 8/27/2007 | Fingerling | | |
| 2007 | Miles City | 336 | 2007 | 8/27/2007 | Fingerling | | |
| 2007 | Wolf Point | 672 | 2007 | 8/27/2007 | Fingerling | | |
| 2007 | Forsyth | 690 | 2007 | 8/31/2007 | Fingerling | CWT | |
| 2007 | Intake | 615 | 2007 | 8/31/2007 | Fingerling | CWT | |
| 2007 | School Trust | 1160 | 2007 | 9/6/2007 | Fingerling | CWT | |
| 2007 | Intake | 293 | 2007 | 9/12/2007 | Fingerling | | |
| 2007 | Miles City | 293 | 2007 | 9/12/2007 | Fingerling | | |
| 2007 | Wolf Point | 586 | 2007 | 9/12/2007 | Fingerling | | |
| 2007 | Culbertson | 6455 | 2007 | 9/14/2007 | Fingerling | Elastomer | |
| 2007 | Fallon | 4827 | 2007 | 9/14/2007 | Fingerling | Elastomer | |
| 2007 | Forsyth | 5370 | 2007 | 9/14/2007 | Fingerling | Elastomer | |
| 2007 | Intake | 7812 | 2007 | 9/14/2007 | Fingerling | Elastomer | |
| 2007 | School Trust | 6096 | 2007 | 9/14/2007 | Fingerling | Elastomer | |
| 2007 | Sidney | 1934 | 2007 | 9/14/2007 | Fingerling | Elastomer | |
| 2007 | Wolf Point | 6455 | 2007 | 9/14/2007 | Fingerling | Elastomer | |
| 2008 | Culbertson | 1384 | 2007 | 5/7/2008 | Yearling | PIT Tag | Scute Removed |
| 2008 | Culbertson | 643 | 2007 | 3/26/2008 | Yearling | Elastomer | |
| 2008 | Fallon | 1307 | 2007 | 5/7/2008 | Yearling | PIT Tag | Scute Removed |
| 2008 | Forsyth | 1384 | 2007 | 5/7/2008 | Yearling | PIT Tag | Scute Removed |
| 2008 | Forsyth | 106 | 2007 | 3/26/2008 | Yearling | Elastomer | |
| 2008 | Intake | 2395 | 2007 | 5/7/2008 | Yearling | PIT Tag | Scute Removed |
| 2008 | Intake | 103 | 2007 | 3/26/2008 | Yearling | Elastomer | |
| 2008 | School Trust | 1325 | 2007 | 5/7/2008 | Yearling | PIT Tag | Scute Removed |
| 2008 | School Trust | 654 | 2007 | 3/26/2008 | Yearling | Elastomer | |
| 2008 | Sidney | 149 | 2007 | 5/7/2008 | Yearling | PIT Tag | Scute Removed |

| Year | Stocking Site | Number Stocked | Year Class | Stock Date | Age at Stocking ^a | Primary Mark | Secondary Mark |
|------|---------------|----------------|------------|------------|------------------------------|--------------|----------------|
| 2008 | Sidney | 67 | 2007 | 3/26/2008 | Yearling | Elastomer | |
| 2008 | Wolf Point | 1328 | 2007 | 5/7/2008 | Yearling | PIT Tag | Scute Removed |
| 2008 | Wolf Point | 416 | 2007 | 3/26/2008 | Yearling | Elastomer | |
| 2008 | Miles City | 4797 | 2008 | 7/30/2008 | Fry | | |
| 2008 | Grand Champs | 24395 | 2008 | 7/30/2008 | Fry | | |
| 2008 | Culbertson | 15630 | 2008 | 9/24/2008 | Fingerling | Elastomer | |
| 2008 | Fallon | 7930 | 2008 | 9/29/2008 | Fingerling | Elastomer | |
| 2008 | Forsyth | 7723 | 2008 | 9/29/2008 | Fingerling | Elastomer | |
| 2008 | Intake | 12642 | 2008 | 9/29/2008 | Fingerling | Elastomer | |
| 2008 | Sidney | 3186 | 2008 | 9/29/2008 | Fingerling | Elastomer | |
| 2008 | Wolf Point | 11717 | 2008 | 9/24/2008 | Fingerling | Elastomer | |
| 2009 | Culbertson | 1387 | 2008 | 4/13/2009 | Yearling | PIT Tag | Scute Removed |
| 2009 | Fallon | 1155 | 2008 | 4/13/2009 | Yearling | PIT Tag | Scute Removed |
| 2009 | Forsyth | 1166 | 2008 | 4/13/2009 | Yearling | PIT Tag | Scute Removed |
| 2009 | Intake | 2181 | 2008 | 4/13/2009 | Yearling | PIT Tag | Scute Removed |
| 2009 | Sidney | 710 | 2008 | 4/13/2009 | Yearling | PIT Tag | Scute Removed |
| 2009 | Wolf Point | 2162 | 2008 | 4/13/2009 | Yearling | PIT Tag | Scute Removed |
| 2009 | Miles City | 46260 | 2009 | 7/31/2009 | Fry | | |
| 2009 | Wolf Point | 26175 | 2009 | 7/22/2009 | Fry | | |
| 2009 | Culbertson | 10238 | 2009 | 9/24/2009 | Fingerling | Elastomer | |
| 2009 | Fallon | 5133 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2009 | Forsyth | 5386 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2009 | Intake | 8374 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2009 | Sidney | 1865 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2009 | Wolf Point | 9946 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2009 | Intake | 8374 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2009 | Sidney | 1865 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2009 | Wolf Point | 9946 | 2009 | 9/23/2009 | Fingerling | Elastomer | |
| 2010 | Fallon | 721 | 2009 | 4/15/2010 | Yearling | PIT Tag | Scute Removed |
| 2010 | Fallon | 268 | 2009 | 8/3/2010 | Yearling | PIT Tag | Scute Removed |
| 2010 | Fallon | 1000 | 2010 | 10/7/2010 | Fingerling | Elastomer | |
| 2010 | Forsyth | 1402 | 2009 | 4/15/2010 | Yearling | PIT Tag | Scute Removed |

| Year | Stocking Site | Number Stocked | Year Class | Stock Date | Age at Stocking ^a | Primary Mark | Secondary Mark |
|------|---------------|----------------|------------|------------|------------------------------|--------------|------------------|
| 2010 | Forsyth | 268 | 2009 | 8/3/2010 | Yearling | PIT Tag | Scute Removed |
| 2010 | Intake | 1890 | 2009 | 4/15/2010 | Yearling | PIT Tag | Scute Removed |
| 2010 | Intake | 816 | 2009 | 6/4/2010 | Yearling | Elastomer | |
| 2010 | Intake | 541 | 2009 | 8/3/2010 | Yearling | PIT Tag | Scute Removed |
| 2010 | Intake | 1000 | 2010 | 10/7/2010 | Fingerling | Elastomer | |
| 2010 | Sidney | 331 | 2009 | 4/15/2010 | Yearling | PIT Tag | Scute Removed |
| 2010 | Wolf Point | 1309 | 2009 | 4/15/2010 | Yearling | PIT Tag | Elastomer, Scute |
| 2010 | Wolf Point | 858 | 2009 | 6/4/2010 | Yearling | Elastomer | |
| 2010 | Wolf Point | 425 | 2009 | 8/3/2010 | Yearling | PIT Tag | Scute Removed |
| 2010 | Wolf Point | 1000 | 2010 | 10/7/2010 | Fingerling | Elastomer | |
| 2010 | Culbertson | 65 | 2004 | 9/21/2010 | 6 Yr Old | PIT Tag | |
| 2010 | Culbertson | 1337 | 2009 | 4/15/2010 | Yearling | PIT Tag | Elastomer, Scute |
| 2010 | Culbertson | 384 | 2009 | 6/4/2009 | Yearling | PIT Tag | Scute Removed |
| 2010 | Culbertson | 1000 | 2010 | 10/7/2010 | Fingerling | Elastomer | |
| 2010 | School Trust | 1766 | 2009 | 4/15/2010 | Yearling | PIT Tag | Elastomer, Scute |
| 2011 | Culbertson | 795 | 2010 | 5/5/2011 | Yearling | PIT Tag | Scute |
| 2011 | Wolf Point | 797 | 2010 | 5/5/2011 | Yearling | PIT Tag | Scute |
| 2011 | Fallon | 531 | 2010 | 5/5/2011 | Yearling | PIT Tag | Scute |
| 2011 | Forsyth | 545 | 2010 | 5/5/2011 | Yearling | PIT Tag | Scute |
| 2011 | Intake | 510 | 2010 | 5/5/2011 | Yearling | PIT Tag | Scute |
| 2011 | Culbertson | 262 | 2010 | 8/22/2011 | Yearling | PIT Tag | Scute |
| 2011 | Fallon | 131 | 2010 | 8/22/2011 | Yearling | PIT Tag | Scute |
| 2011 | Forsyth | 174 | 2010 | 8/22/2011 | Yearling | PIT Tag | Scute |
| 2011 | Intake | 132 | 2010 | 8/22/2011 | Yearling | PIT Tag | Scute |
| 2011 | Wolf Point | 262 | 2010 | 8/22/2011 | Yearling | PIT Tag | Scute |

^aAge of fish when stocked: Fry, Fingerling, Yearling, 1yo, 2yo, 3yo, etc...

Appendix G. Hatchery names, locations, and abbreviations.

| Hatchery | State | Abbreviation |
|-------------------------------------|-------|--------------|
| Blind Pony State Fish Hatchery | MO | BYP |
| Neosho National Fish Hatchery | MO | NEO |
| Gavins Point National Fish Hatchery | SD | GAV |
| Garrison Dam National Fish Hatchery | ND | GAR |
| Miles City State Fish Hatchery | MT | MCH |
| Blue Water State Fish Hatchery | MT | BLU |
| Bozeman Fish Technology Center | MT | BFT |
| Fort Peck State Fish Hatchery | MT | FPH |