

2013 Annual Report

Pallid Sturgeon Population Assessment and Associated Fish Community Monitoring for the Missouri River: Segment 1



**Prepared for the U. S. Army Corps of Engineers - Missouri River Recovery Program
By:**

Tyler Haddix, Landon Holte and John Hunziker

**Montana Fish, Wildlife & Parks
Government Bldg.
East Kansas Street
Fort Peck, MT 59223**

May 2014

EXECUTIVE SUMMARY

Segment 1 of the Missouri River has been sampled during each year starting in 2006 to 2013, during both the sturgeon and fish community seasons. Segment 1 is situated directly downstream of Fort Peck Dam and is the most highly altered area of the Missouri River downstream of Fort Peck Dam within Montana. Segment 1 is used as a reference area, whereas the same river bend is sampled annually.

Over the eight years of sampling Segment 1 it has become apparent that the altered nature of the river precludes several native species from thriving. No pallid sturgeon *Schaphirynchus albus* have been collected during standardized sampling. In addition, no sicklefin chub *Macrhybopsis meeki* and only one sturgeon chub *M. gelida* have been collected during the sixteen sampling events over eight years. While some target species have been absent or somewhat rare, adult shovelnose sturgeon *S. platyrhynchus*, blue suckers *Cycleptus elongatus* and sand shiners *Notropis stramineus* have consistently been collected.

During 2013 a total of 85 shovelnose sturgeon were collected in Segment 1. Shovelnose sturgeon averaged 667.6 mm in length and 1,315.3 g in weight. The smallest shovelnose sturgeon captured measured 558 mm, with the largest specimen measuring 870 mm. No young-of-the-year or age-1 sized shovelnose sturgeon have been sampled within Segment 1 through the eight years of sampling.

The relative abundance of shovelnose sturgeon has remained relatively constant through the past eight years with the exception of 2009. During 2009 trammel net catch-per-unit-effort (CPUE) was at an eight year high with 4.5 fish/100 m during the fish community season. However, during the sturgeon season of the same year, CPUE was estimated at only 0.05 fish/100m, the lowest recorded trammel net CPUE for the same eight year period. During 2013 trammel net CPUE was higher during the fish community season (CPUE = 0.5 fish/100 m) than the sturgeon season (CPUE = 0.3 fish/100 m). Similarly, shovelnose sturgeon CPUE of trotlines was higher during the sturgeon season than during the fish community season.

The size distribution of shovelnose sturgeon captured within Segment 1 has not changed appreciably since 2006, with only adult sized fish being captured. Additionally, the length-weight relationship for Segment 1 caught shovelnose sturgeon has remained relatively constant over the seven sampling seasons.

Other native target species that were collected in 2013 include blue suckers, sauger *Sander canadense*, sand shiners and western silvery minnows. A total of four blue suckers, two sauger, two western silvery minnows, and 18 sand shiners were collected.

Longnose suckers *Catostomus catostomus* were the most abundant species captured in 2013, with a total of 568 sampled. In addition to longnose suckers, 105 white suckers *Catostomus commersoni* were captured. Over the eight years of sampling, white and longnose suckers have been two of the dominant species sampled within Segment 1.

TABLE OF CONTENTS

Introduction.....	1
Study Area	4
Methods.....	5
Sample site selection and description	5
Sampling gear	6
Data Collection and Analysis.....	8
Results	
Pallid sturgeon	12
Targeted Native River Species.....	12
Missouri River Fish Community	24
Discussion	24
Acknowledgments.....	27
References.....	28
Appendices.....	29

LIST OF TABLES

Table 1. Presence/absence of all fishes collected in Segments 1 through 3 of the Missouri River from 2006 through 2013.	22
--	----

LIST OF FIGURES

Figure 1. Map of Segment 1 of the Missouri River with major tributaries, common landmarks, and historic stocking locations for pallid sturgeon. Segment 1 encompasses the Missouri River from Fort Peck Dam (River Mile 1771.5) to the confluence of the Milk River (River Mile 1760).	10
Figure 2. Hydrograph (2006-2013) for Segment 1 of the Missouri River.	11
Figure 3. Mean annual catch-per-unit-effort by season of selected species collected using trammel nets in Segment 1 from 2006 through 2013.	15
Figure 4. Mean annual catch-per-unit-effort by season of selected species captured using trotlines in Segment 1 from 2008 to 2013.	16
Figure 5. Mean annual catch-per-unit-effort by season of selected species collected using otter trawls in Segment 1 from 2006 through 2013.	17
Figure 6. Length frequency histograms for all shovelnose sturgeon sampled in Segment from 2006 through 2013	18
Figure 7. Weight-length relationship for all shovelnose sturgeon sampled in Segment 1 from 2006 through 2013.	19
Figure 8. Mean annual catch-per-unit-effort by season of target and non-target species collected using mini fyke nets in Segment 1 from 2006 through 2013.	20
Figure 9. Mean annual catch-per-unit-effort by season of selected non-target species collected using mini fyke nets in Segment 1 from 2006 through 2013.	21

LIST OF APPENDICES

Appendix A. Phylogenetic list of Missouri River fishes with corresponding letter codes used in the long-term pallid sturgeon and associated fish community sampling program.	30
Appendix B. Definitions and codes used to classify standard Missouri River habitats in the long term pallid sturgeon and associated fish community sampling program.....	36
Appendix C. List of standard and wild gears, their corresponding codes in the database, seasons deployed, years used, and catch-per-unit-effort units for collection of Missouri River fishes for the long-term pallid sturgeon and associated fish community sampling program	37
Appendix D. Stocking locations and codes for pallid sturgeon by Recovery Priority Management Area in the Missouri River Basin.....	38
Appendix E. Juvenile and adult pallid sturgeon stocking summary for Segment 1 of the Missouri River (RPMA 2).	40
Appendix G. Hatchery names, locations, and abbreviations.	46

Introduction

The U.S. Fish and Wildlife Service (USFWS) listed the pallid sturgeon *Scaphirhynchus albus* as endangered in 1990. In response to the listing, the USFWS issued a Biological Opinion to the U.S. Army Corps of Engineers (COE), the main water management entity responsible for the Missouri River mainstem from Fort Peck Dam and Reservoir to its confluence with the Mississippi River. Additionally, an amendment to the 2000 Biological Opinion was issued in 2003. The Amendment listed several Reasonable and Prudent Alternatives (RPA) to address the inability of pallid sturgeon to naturally reproduce and the need to be able to detect changes in their populations and ecosystem trends.

The Pallid Sturgeon Population Assessment Program (Program) is guided by the RPA's in the 2003 Amendment to the 2000 Biological Opinion. The Program is a comprehensive monitoring plan designed to assess survival, movement, distribution, habitat use, and physical characteristics of these habitats used by wild and hatchery reared juvenile pallid sturgeon (Welker and Drobish 2011). The 2000 Biological Opinion divides the Program area into river and reservoir Segments and assigns high, moderate, or low priority management action areas to these Segments for pallid sturgeon (Welker and Drobish 2011). The focus of the Program is on the high priority management action Segments. The Missouri River from Fort Peck Dam downstream to the headwaters of Lake Sakakawea, ND is listed as a high priority action Segment.

The Program has stratified the Missouri River from Fort Peck Dam to the headwaters of Lake Sakakawea into four study Segments based on biological, hydrological and fluvial geomorphological characteristics. The COE has contracted Montana Fish, Wildlife & Parks (FWP) to conduct the Program sampling in the Missouri River from Fort Peck Dam downstream to its confluence with the Yellowstone River, which consists of study Segments 1 through 3.

This was the eighth field season that Montana Fish, Wildlife & Parks conducted standard Program sampling in Segment 1 of the Missouri River.

The objectives of this program are as follows:

1. Document annual results and long-term trends in pallid sturgeon population abundance and geographic distribution throughout the Missouri River System.
2. Document annual results and long-term trends of habitat use of wild pallid sturgeon and hatchery stocked pallid sturgeon by season and life stage.
3. Document population structure and dynamics of pallid sturgeon in the Missouri River System.
4. Evaluate annual results and long-term trends in native target species population abundance and geographic distribution throughout the Missouri River system.
5. Document annual results and long-term trends of habitat usage of the native target species by season and life stage.
6. Document annual results and long-term trends of all non-target species population abundance and geographic distribution throughout the Missouri River system, where sample size is greater than fifty individuals.

Sampling Season and Species

The Program has two discrete seasons (sturgeon and fish community), which are primarily based on water temperatures. However, the sturgeon season is designed to sample sturgeon with gears that are temperature dependent, such as gill nets. Due to the nature of the majority of habitats in Segment 1 through 3, gill nets are not an efficient gear for collecting pallid sturgeon in Montana to debris flows and swift currents and therefore they are not used within Segments 1 through 3. Trammel nets and otter trawl are standard gears used in Segments 1-3 during sturgeon season, and so far appear to be an effective way to sample pallid sturgeon. Trotlines are now a standard gear used in Segment 1.

The fish community season runs from the beginning of July till the end of October and is designed not only to monitor sturgeon, but also to monitor other native Missouri River fish populations. Both trammel nets and the otter trawl are still used, but to more effectively sample shallow water habitats < 1.2 m in depth, mini fyke nets are also used as a standard gear.

In addition to pallid sturgeon, the Program is designed to monitor nine other native Missouri River species labeled “target” species. These include, shovelnose sturgeon *Scaphirhynchus platyrhynchus*, blue sucker *Cycleptus elongatus*, sauger *Sander canadensis*, sturgeon chub *Macrhybopsis gelida*, sicklefin chub *M. meeki*, speckled chub *M. aestivalis*, plains minnow *Hybognathus placitus*, western silvery minnow *H. argyritis*, and sand shiner

Notropis stramineus. This suite of species was selected for various reasons. First, some species may have similar habitat requirements as pallid sturgeon and therefore by monitoring their populations we may gain further insight into pallid sturgeon habitat and how anthropomorphic and natural changes to the Missouri River affect native fish assemblages. Secondly, it is hypothesized that various chub species and other native fishes are an important component of pallid sturgeon diet. Thirdly, we wouldn't expect to see an immediate response in a long-lived species such as the pallid sturgeon when environmental conditions change from either favorable or detrimental conditions. Thus, by monitoring short-lived native fishes we may be able to correlate environmental conditions to changes in fish populations on a much shorter time interval and make inferences on how pallid sturgeon populations and their habitat are being affected.

Study Area

Segment 1 of the Missouri River begins at Fort Peck Dam and runs downstream to its confluence with the Milk River. This Segment constitutes only 6% (11.5 river miles) of the entire 189.5 river miles downstream of Fort Peck Dam to the headwaters of Lake Sakakawea in North Dakota (Welker and Drobish 2011). This reach of the Missouri River is characterized by an unnatural hydrograph, thermograph, sediment dynamics, and fish community due to the influence of Fort Peck Dam, which was constructed in 1940 (Bramblett and White, 2001). Segment 1 includes the Fort Peck Dredge Cuts, a deepened and widened section of river immediately below the dam created by the dredging of earth used to construct the dam. Regulated hypolimnetic water releases from Fort Peck Reservoir have changed a once turbid sandy bottom stretch of river into a cold clear cobble dominated river. Fort Peck Reservoir has substantially reduced suspended sediment loads in the river below Fort Peck Dam when compared to its natural state (Galat et al, 2005).

Peaks in the hydrograph are related to power production and barge traffic downstream, instead of natural spring runoff and precipitation events (Galat et al, 2005). Many species native to this stretch of river such as the pallid sturgeon, sicklefin chub and sturgeon chub find the cold clear water unsuitable and are now common only farther downstream where tributaries have warmed and muddied the waters of the Missouri (Gardner and Stewart, 1987). Fish much more suited for this cold clear water such as rainbow trout *Oncorhynchus mykiss*, brown trout *Salmo trutta* and Chinook salmon *Oncorhynchus tshawytscha* have been stocked on and off from 1950 to 1990. Other nonnative species such as largemouth bass *Micropterus salmoides*, northern pike *Esox lucius*, walleye *Sander vitreus*, and yellow perch *Perca flavescens* have been stocked in the dredge cuts to increase angling opportunities. It is believed that many of these sight-feeding piscivores have out competed the native fishes in this stretch of river (Galat et al, 2005). In summary, this unique stretch of river is now vastly different from the once braided and shifting channels of the “Big Muddy” before Fort Peck Dam (Galat et al, 2005).

Methods

Sampling methods for the Pallid Sturgeon Population Assessment Program were conducted in accordance with the Standard Operating Procedures (Welker and Drobish 2011), which was established by representatives from State and Federal agencies involved with pallid sturgeon recovery on the Missouri River. For a detailed description of methodologies please see Welker and Drobish (2011). A general description of those guidelines follows.

Sampling Site Selection and Description

Montana Fish Wildlife & Parks (FWP) was contracted to sample Segment 1 from Fort Peck Dam (RM 1771.5) to the mouth of the Milk River (RM 1761), Segment 2 from the mouth of the Milk River (RM 1761) to Wolf Point (RM 1701.5) and Segment 3 from Wolf Point (RM 1701.5) to the Montana/North Dakota border (RM 1586.5). Segment 1 consists of one non-random bend at river mile 1766. Segment 1 was selected as a reference study bend to be sampled each year to facilitate comparisons of the most highly altered area of the Missouri River in Recovery Priority Management Area (RPMA) 2 to downstream areas (Segments 2 through 4). By comparing data from Segment 1 with downstream Segments, a better understanding of how Fort Peck Dam influences the fish communities of the Missouri River might be attained.

During 2013 Segment 1 was sampled on May 20th during the sturgeon season and September 18th, 19th and October 30th during the fish community season. Four standard gears were used, trammel net, otter trawl, and trotlines were used during both the sturgeon and fish community seasons and mini-fyke nets during the fish community season.

Missouri River flows were relatively high downstream of Fort Peck Dam from May through June of 2013, when compared to the other seven sampling seasons (Figure 2). Flows were significantly lower during the fish community season sampling dates.

The Population Assessment Team developed a standard set of habitat classifications for the Missouri River (Appendix B) which consists of three distinct macrohabitats found in every bend, a main channel crossover (CHXO), main channel outside bend (OSB), and main channel inside bend (ISB). Each sampling bend was comprised of these three main

macrohabitats. Nine additional macrohabitats were identified that may or may not be present in every bend: large tributary mouths (TRML), small tributary mouths (TRMS), confluence areas (CONF), large and small secondary connected channels (SCCL& SCCS), deranged channels (DRNG), braided channels (BRAD), dendritic channels (DEND) and non-connected secondary channel (SCN). For the reference bend in Segment 1, five macrohabitats were sampled, CHXO, OSB, ISB, SCCL and SCCS.

Mesohabitats were established to further define macrohabitats. Mesohabitats include bars (BARS), pools (POOL), channel border (CHNB), thalweg (TLWG) and island tip (ITIP). Channel borders are situated in areas between the deepest portions of the river up to a depth of 1.2 m. Bars are considered shallow areas (< 1.2 m) where terrestrial and aquatic habitats merge. The thalweg is the deepest portion of the river between the two channel borders where the majority of the flow is directed. Pools are directly downstream of any feature that creates scour, thus creating a habitat of deep (> 1.2 m) slower moving water. Island tips are just downstream of bars or islands where two channels meet where the water is > 1.2 m in depth. Two mesohabitats were sampled in Segment 1, CHNB and BARS.

For all analysis, the sampling unit was the river bend, where every river bend has a channel crossover, inside and outside bend. The downstream border of a river bend is the beginning of the next downstream bend's channel crossover.

Sampling Gear

For specific information pertaining to the habitats gears are utilized in and physical measurements taken in accordance with sampling the various gears described below please see Welker and Drobish (2011).

Trammel Net

The standard trammel net has a length of 38.1 m, an inner mesh wall 2.4 m and two outer mesh walls 1.8 m deep. The inner mesh is made of #139 multifilament twine with a bar mesh size of 25.4 mm. The outer walls are constructed of #9 multifilament twine with a bar mesh size of 203.2 mm. The float line is a 12.7 mm diameter foam core with a lead line of 22.7 kg. Trammel nets were drifted from the bow of the boat and orientated perpendicular to the river flow for a minimum of 75 m and a maximum drift distance of 300 m.

Otter Trawl

The standard otter trawl has a length of 7.6 m, a width of 4.9 m and height of 0.9 m. The otter trawl has an inner mesh (6.35mm bar, #18 polyethylene twine) and outer mesh (38mmbar, #9 polyethylene twine) and a cod end opening of 406.4 mm. The trawl doors were made from 19.1 mm marine plywood and measured 762 mm x 381 mm. The trawl doors are used to keep the mouth of the trawl open while deployed on the riverbed. The trawl also has a 7.9 m long tickler chain attached to the bottom of the mouth of the trawl, which aids in keeping it orientated on the riverbed and protecting the mouth when snags are encountered. The otter trawl was deployed from the bow of the boat parallel to the current with two 30.5 m ropes and towed downstream slightly faster than current speed for a minimum of 75 m and a maximum distance of 300 m.

Mini-Fyke Nets

The standard mini-fyke net consists of two rectangular frames 1.2 m wide and 0.6 m high and two 0.6 m tempered steel hoops. A 4.5 m long and 0.6 m high lead is connected to the first frame. The fyke net is made of 3 mm “ace” style mesh. The lead has small floats attached to the top and lead weights on the bottom. Mini-fyke nets are set with a “T” stake on shore and extend into river as perpendicular to the shoreline as possible or angled slightly downstream where higher velocities existed. Mini-fyke nets were set overnight and checked the following morning.

Trotlines

Trotlines consisted of 32 m nylon rope attached to both upstream and downstream anchors. Octopus style circle hooks were attached to the ropes using 136 kg monofilament line and commercial fishing clips. Twenty 45.7 cm leaders were used on each trotline. Hooks consisted of 2/0 circle hooks. Each trotline used one hook size and each hook size was used at least once in each macrohabitat sampled. Trotlines were baited with night crawlers, and were set overnight then checked the following morning.

Data Collection and Analysis

A minimum of eight random subsamples with each gear were deployed in the reference bend in Segment 1. At least two subsamples (when possible) were taken with each gear in each macro habitat within the bend. More than two subsamples were taken in a macrohabitat for a gear when the number of discrete macrohabitats was less than four or less than four could be effectively sampled.

All fish were measured to the nearest mm. Fork length (FL) was used for sturgeon species, while other species were measured to TL with one exception, paddlefish *Polyodon spathula*, which were measured from the eye to the fork of the caudal fin. The first 25 fish of each species in each subsample were measured, after 25 they were counted.

Time was recorded at the beginning of each sample with all gears and an end time was always recorded when pulling mini-fyke net sets. A global positioning satellite (GPS) position was taken at the beginning and end of all otter and beam trawls and trammel net drifts. One GPS location was taken for mini fyke net sets. All GPS locations were taken using a Garmin GPS 76 unit with Wide Area Augmentation System (WAAS) capability.

Sample depth was determined at the beginning, middle and end of each trawl and drift using a Lowrance X136 sonar unit. One depth was taken for mini-fyke nets at the intersection of the frame and floatline using a wading rod.

Water temperature taken near the surface was recorded at every sample using the Lowrance X136 unit for trawls and trammel net drifts and using a hand held thermometer for mini fyke nets.

Habitat samples were collected randomly for 25% of each mesohabitat within each macrohabitat sampled. Velocities (mps) were taken at three depths in the water column for habitats > 1.2 m in depth (bottom, 0.8 of bottom depth and 0.2 of the bottom depth) using either a Current AA Price Meter and sounding reel or a Marsh-McBirney Flo Mate 2000. Velocities for shallow water habitats (< 1.2 m) were taken at the bottom and 0.6 of the bottom depth using the March-McBirney Flo Mate 2000.

Turbidity was recorded in nephelometric turbidity units (NTU) using a LaMotte 2020 turbidity meter. Turbidity was taken at the midpoint of all samples, except mini-fyke sets, where it was taken at the convergence of the rectangular frame and float line.

In addition to 25% of all mesohabitats, habitat measurements were taken whenever a pallid sturgeon was captured.

Genetic Verification

Genetic verification for pallid sturgeon or potential hybrids followed the methods outlined in Welker and Drobish (2011). Two fin pectoral fin clips ($\sim 2 \text{ cm}^2$) are taken from any pallid sturgeon of unknown origin. Fin samples are then preserved in 95% non-denatured alcohol for genetic analysis. All samples are sent to the U.S. Fish and Wildlife Service's Northeast Fishery Center Conservation Genetics Lab for analysis and archiving.

Analyses

The fundamental sampling unit for the Population Assessment Program is the river bend, where sample size is equal to the number of bends sampled. Since only one river bend in Segment 1 is sampled per year, only one true sample was taken. Therefore, all CPUE data for Segment 1 are the averages of all subsamples and no error is associated with these estimates.




Figure 1. Map of Segment 1 of the Missouri River with major tributaries, common landmarks, and historic stocking locations for pallid sturgeon. Segment 1 encompasses the Missouri River from Fort Peck Dam (River Mile 1771.5) to the mouth of the Milk River (River Mile 1760.0).

Missouri River Below Fort Peck Dam




Figure 2. Missouri River discharge downstream of Fort Peck Dam from 2006 to 2013.
Source: USGS.

Results

Pallid Sturgeon

No pallid sturgeon or pallid sturgeon x shovelnose sturgeon hybrids were collected in Program sampling within Segment 1 of the Missouri River during 2013. This was the eighth year of standardized sampling where no pallid sturgeon were sampled. However, during 2012 a 1997 year class hatchery reared pallid sturgeon was captured by FWP during their annual fall gill netting effort in the Fort Peck Dredge Cuts. This fish was not tagged and a genetics sample was taken and subsequently sent to the USFWS Lamar Genetics Laboratory, where it was confirmed as a hatchery fish from the 1997 year class. After capture, this fish received a PIT Tag (#48686C7324) and a radio transmitter (frequency 149.760, code 64). This was the first hatchery reared pallid sturgeon captured within Segment 1 by any crew since stocking began in 1998.

Targeted Native River Species

Shovelnose Sturgeon

A total of 85 shovelnose sturgeon were sampled in Segment 1 during 2013, 52 and 33 during the sturgeon and fish community seasons, respectively. However, only 46 were sampled in standardized sampling, while the other 39 were captured in targeted trotline sets in adjacent areas. During the sturgeon season, 46 shovelnose sturgeon were sampled using trotlines (Figure 3), while 4 were captured using trammel nets and 2 using the otter trawl (Figure 4). During the fish community season, 30 shovelnose sturgeon were captured using trammel nets, while only 2 were sampled using the otter trawl, and one using trotlines (Figure 6).

Shovelnose sturgeon CPUE in trammel nets was lower in 2013 when compared to 2012 during the sturgeon season, but was similar for both years for the fish community season (Figure x). No discernible pattern has been witnessed in regards to trammel net CPUE between the sturgeon and fish community seasons over the past eight years of sampling.

Trotline CPUE of shovelnose was considerably lower during 2013 for both seasons, when compared to the majority of previous sampling years (Figure 4). Trotline CPUE has decreased during the fish community season in each year sampled (Figure 4). Shovelnose

sturgeon have been the one of, if not the most abundant species captured using trotlines in Segment 1 during each sampling occasion beginning in 2008.

Few shovelnose sturgeon have been captured in the otter trawl over the past eight years (Figure 5). This is likely due to the fact that no shovelnose smaller than 450 mm have been captured in any gear (Figure 6).

The size structure of shovelnose sturgeon sampled in Segment 1 during both seasons has not changed appreciably over the last seven sampling seasons (Figure 6). Furthermore, no juvenile shovelnose sturgeon under 450 mm in length have been captured in any gear over the past eight years of sampling. In addition to the size structure remaining very similar between years, the length to weight relationship of shovelnose sturgeon has also not changed significantly since 2006 (Figure 9).

Sturgeon Chub

No sturgeon chubs were collected in Segment 1 during sampling in 2013. Only one sturgeon chub has been collected to date in Segment 1 through seven years of sampling and that fish was collected in the otter trawl during 2010 (Figure 5).

Sicklefin Chub

No sicklefin chubs were captured in 2013 and none have been collected through the eight years of sampling Segment 1.

Sand Shiner

Eighteen sand shiners were captured in 2013, all in mini fyke nets set during the fish community season (Figure 8). This was the fourth highest year for sand shiners captured in mini fyke nets within Segment 1. No sand shiners were captured in both 2011 and 2006. The highest CPUE for sand shiners was observed during 2010, with a CPUE of 6.5/net night.

Western Silvery Minnow

Two western silvery minnows were collected in 2013 in Segment 1. Western silvery minnows have only been collected in three out of the eight years of sampling, with over three fish per net night in 2008 and just over one per net night in 2010 (Figure 8).

Blue Sucker

A total of four adult blue suckers were sampled within Segment 1 during 2013, three with trammel nets (Figure 3) and one on trotlines (Figure 4). Blue suckers have been sampled every year in Segment 1, although in very low numbers. Based on aging of blue suckers downstream of Gavins Point Dam, these fish are likely older than age 7 (Labay et al. 2008).

Sauger

Two sauger were sampled in Segment 1 during 2013, one in the otter trawl and one in a mini fyke net. Both sauger were sampled during the fish community season. Very few sauger have been sampled in Segment 1 in the past seven years (Figures 3, 4 5, 8 and 9)).

Segment 1 Trammel Net




Figure 3. Trammel net CPUE by season for all target species and non-target species sampled in Segment 1 of the Missouri River during sturgeon and fish community season from 2006 through 2013. Target species are indicated by an asterisk. Note the difference in scale of the Y-axes.

Segment 1 Trotline CPUE




Figure 4. Trotline CPUE for all target and non-target species sampled in the sturgeon season (top panel) from 2009 through 2013, and the fish community season (bottom panel) from 2008 through 2013 in Segment 1 of the Missouri River. Note that trotlines were not set in 2008 during the Sturgeon Season or in 2009 during the Fish Community Season. Shovelnose sturgeon are a target species.

Segment 1 Otter Trawl




Figure 5. Otter Trawl CPUE by season for all target species and non-target species sampled in Segment 1 of the Missouri River during sturgeon and fish community season from 2006 through 2013. Target species are indicated by an asterisk. Note the difference in scale of the Y-axes.

Segment 1 Shovelnose Sturgeon




Figure 6. Length frequency histogram for all shovelnose sturgeon sampled in Segment 1 of the Missouri River from 2006 through 2013.

Segment 1 Shovelnose Sturgeon




Figure 7. Weight-length relationship for all shovelnose sturgeon sampled in Segment 1 from 2006 through 2013.

Segment 1 Mini Fyke Net




Figure 8. Mini-fyke net CPUE for target and non-target species sampled during the fish community season in Segment 1 of the Missouri River from 2006 through 2013. Target species are indicated by asterisks. Note the differences in Y-axes.

Segment 1 Mini Fyke Net




Figure 9. Mini-fyke net CPUE for non-target species sampled during the fish community season in Segment 1 of the Missouri River from 2006 through 2013. Note the differences in Y-axes.

Table 1. Presence absence of all species (by common name) collected in Segments 1 through 3 in the Missouri River during 2006 through 2013. Boxes marked with an X indicate at least one specimen was sampled. Species in bold are native target species.

Segments			Segments		
1	2	3	1	2	3
<i>Ascipenseridae - sturgeons</i>			<i>Esocidae - pikes</i>		
Pallid sturgeon	X	X	Northern Pike	X	X
Shovelnose sturgeon	X	X	<i>Osmeridae - smelts</i>		
<i>Polyodontidae - paddlefishes</i>			Rainbow smelt		X
Paddlefish	X	X	<i>Ictaluridae - catfishes</i>		
<i>Hiodontidae - mooneyes</i>			Channel catfish	X	X
Goldeye	X	X	Black bullhead		X
<i>Cyprinidae - carps and minnows</i>			Yellow bullhead		X
Common Carp	X	X	Stonecat		X
Flathead chub	X	X	<i>Salmonidae - trouts</i>		
Emerald shiner	X	X	Rainbow trout	X	X
Lake chub	X		Brown trout	X	X
Longnose dace	X	X	Lake Trout	X	
Northern redbelly dace	X	X	Lake whitefish	X	X
Plains minnow		X	Cisco	X	X
Western silvery minnow	X	X	<i>Gadidae - cods</i>		
<i>Brassy minnow</i>			Burbot	X	X
Sicklefin chub		X	<i>Gasterosteidae - sticklebacks</i>		
Sturgeon chub	X	X	Brook stickleback		X
Sand shiner	X	X	<i>Centrarchidae - sunfishes</i>		
Spottail shiner	X	X	Green sunfish		X
Fathead minnow	X	X	Pumkinseed		X
<i>Catostomidae-suckers</i>			White crappie	X	X

Segments			Segments		
1	2	3	1	2	3
Bigmouth buffalo	X	X	Smallmouth bass	X	
Smallmouth buffalo	X	X	<i>Percidae - perches</i>		
Blue sucker	X	X	Iowa darter		X
River carpsucker	X	X	Yellow perch	X	X
White sucker	X	X	Sauger	X	X
Longnose sucker	X	X	Walleye	X	X
Shorthead redhorse	X	X	<i>Sciaenidae - drums</i>		
<i>Moronidae-temperate bass</i>			Freshwater drum		X
White bass		X	<i>Lepisosteidae - Gars</i>		
			Shortnose Gar	X	X

Missouri River Fish Community

A total of 1,162 fish comprising 18 species were collected within Segment 1 during 2013. Longnose suckers *Catostomus catostomus* were the most abundant fish captured, with 568 being captured. Fathead minnows *Pimephales promelas* were the second most abundant species captured with 259 collected, while white suckers *C. commersoni* were the third most abundant species with 105 sampled. The fourth most abundant fish was rainbow trout *Oncorhynchus mykiss* with 81 young-of-the-year captured in the mini fyke nets. Other species that were in very low abundance included common carp *Cyprinus carpio* (n = 3), burbot *Lota lota* (n = 2), channel catfish *Ictalurus punctatus* (n = 3), lake whitefish *Coregonus clupeaformis* (n = 6), longnose dace *Rhinichthys cataractae* (n = 7), northern pike *Esox lucius* (n = 1), shorthead redhorse *Moxostoma macrolepidotum* (n = 8), smallmouth buffalo *Ictiobus bubalus* (n = 6), walleye *Sander vitreus* (n = 2).

Discussion

Segment 1 of the Missouri River is a highly altered Segment due to the proximity of Fort Peck Dam. Fort Peck Dam, a hypolimnetic withdraw structure, is located approximately five river miles upstream of Segment 1, which creates cold summer water temperatures and low suspended sediment loads. Water temperatures averaged 7.1 C° and turbidity averaged 3.4 NTU's during the sturgeon season and 12.7 C° and 5.0 NTU's during the fish community season. In addition, the benthic substrate of Segment 1 is noticeably different than the substrates of downstream Segments. Segment 1 is primarily composed of gravel and cobble due to the degrading stream bed, which is at least in part due to the lack of suspended sediments in the water column.

Although flows throughout the summer were lower in Segment 1 when compared to the previous two years, it was higher than years prior to 2011 (Figure 2).

No pallid sturgeon have been sampled during standardized sampling in Segment 1 from 2006 through 2013, despite continuous stocking from 2004 to 2008 in the confluence area of the

Milk and Missouri Rivers. However, FWP did capture a 1997 year class pallid sturgeon in the Missouri River Dredge Cuts, which is just upstream of Segment 1 during 2012. This was the first documentation of hatchery reared pallid sturgeon using the Missouri River upstream of the Milk River. Furthermore, data have indicated that at least some stocked pallid sturgeon from all other stockings sites in Segments 2 and 3 do move upstream.

Even with the highly altered conditions of Segment 1, many native species are still occupying the habitats of this Segment. However, during the past eight years of sampling, a total of 10 non-native species have been found in Segment 1 including, common carp, rainbow trout, brown trout *Salmo trutta*, lake trout *Salvelinus namaycush*, lake whitefish, ciscoe *C. artedi*, spottail shiner *Notropis hudsonius*, smallmouth bass *Micropterus dolomieu*, and yellow perch *Perca flavescens* (Table 1).

The CPUE of shovelnose sturgeon in Segment 1 continues to be variable (Figure 3 and 4). Overall trammel net CPUE for shovelnose sturgeon was lower in 2013 than in 2012. In addition, sturgeon and fish community season trotline CPUE was lower in 2013 than 2012. Nevertheless, since only eight subsamples are taken during each season, catch is variable and little conclusions can be drawn from the CPUE data. More importantly is the consistency in the size distribution of shovelnose in Segment 1. No shovelnose under 450 mm have been collected in eight years of sampling, which indicates the population is made up of older fish and little to no juvenile rearing is occurring in the area. Past telemetry studies have also shown that the shovelnose sturgeon population in Segment 1 is a resident population of fish that do not migrate far. These fish do not appear to be spawning since we don't collect any black egged females in the spring or early summer, as we do in the lower parts of Segment 3. Tagging information has also shown that adult shovelnose in Segment 1 either do not grow or grow at a very slow rate. Individuals that have been recaptured after 20 years at large often are the same size as they were when they were tagged.

Adult blue sucker have been captured in Segment 1 during all eight sampling years. Similar to shovelnose sturgeon, these have all been large adult fish. Again, CPUE has been variable, but since so few fish have been captured there does not appear to be any differences in their relative abundance over the sampling years.

The mini fyke net catch in Segment 1 has been highly variable over the eight years of sampling (Figures 8 and 9). The last three years of sampling has produced relatively few fish when compared to 2007 and 2008. During those years, relatively low flows were being

discharged out of Fort Peck Dam. We still do not know if the abundance of small bodied minnows is tied to Dam operations, or possibly the abundance of predators in the area.

It doesn't appear that the higher releases from Fort Peck Dam in 2011 and 2012 increased many of the native cyprinids populations that reside within Segment 1. However, we did see a dramatic increase in longnose suckers in 2013 and white suckers in 2012 when compared to 2011.

Continued sampling of Segment 1 may be important as hatchery reared pallid sturgeon in the Missouri River downstream of Fort Peck Dam mature. Monitoring this area will allow us to identify if pallid sturgeon begin using this stretch of the river. Currently, there is very little evidence suggesting that the habitat within Segment 1 is being sought out by pallid sturgeon.

Acknowledgments

The U.S. Army Corps of Engineers provided funding for this project. We'd like to thank Tim Welker and George Williams for supporting our work in several ways. Raina Whitherite, Eddie Bossert-Lomeli, Amanda Bryson and Cara Holem-Bell assisted in both the field and shop throughout the year. Steve Dalbey took care of many things while we were on the river. We'd like to thank the Dave Fuller for his assistance. Thanks to Pat Braaten of the U.S. Geological Survey for answering any type of question we may have pertaining to the Missouri River and its fishes. A special thanks to Ryan Wilson, Zack Sandness, Everett Nelson and Steve Krentz of the U.S. Fish and Wildlife Service for all the collaboration between our offices.

References

- Dattilo, J. E., R. R. Dirnberger, P. T. Horner, D. J. Niswonger, M. L. Miller and V. H. Travinchek. 2008a. Three Year Summary Age and Growth Report For Sand Shiner (*Notropis stramineus*). Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Missouri Department of Conservation. Chillicothe, MO.
- Dattilo, J. E., R. R. Dirnberger, P. T. Horner, D. J. Niswonger, M. L. Miller and V. H. Travinchek. 2008b. Three Year Summary Age and Growth Report For Plains Minnow, Western Silvery Minnow, Brassy Minnow (*Hybognathus spp.*). Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Missouri Department of Conservation. Chillicothe, MO.
- Dattilo, J. E., R. R. Dirnberger, P. T. Horner, D. J. Niswonger, M. L. Miller and V. H. Travinchek. 2008c. Three Year Summary Age and Growth Report for Sauger (*Sander canadensis*). Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Missouri Department of Conservation. Chillicothe, MO.
- Galat, D.L., C.R. Berry Jr., E.J. Peters and R.G. White. 2005. Missouri River. Pages 427-480 in A.C. Benke and C.E. Cushing (editors). Rivers of North America, Elsevier, Oxford.
- Gardner, W.M. and P.A. Stewart. 1987. The Fishery of the Lower Missouri River. Federal Aid to Fish and Wildlife Restoration Project FW-2-R Job I-b. Montana Fish, Wildlife and Parks. Helena, Montana.
- Labay, S., J. Kral and S. Stukel. 2008. Three Year Summary Age and Growth Report for Blue Sucker. Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. South Dakota Department of Game, Fish and Parks. Yankton, SD.
- Pierce, C. L., C. S. Guy, P. J. Braaten, and M.A. Pegg. 2004. Fish growth, mortality, recruitment, condition, and size structure. Volume 4. Population structure and habitat use of benthic fishes along the Missouri and lower Yellowstone Rivers. U.S. Geological Survey, Cooperative Research Units, Iowa State University, Ames Iowa.
- Steffensen, K. and M. Hamel. 2008. Four Year Summary Age and Growth Report For Shovelnose Sturgeon. Pallid Sturgeon Population Assessment Project and Associated Fish Community Monitoring for the Missouri River. Nebraska Game and Parks Commission. Lincoln, NE.
- Welker, T. L., and M. R. Drobish. (editors), 2011. Missouri River Standard Operating Procedures for Fish Sampling and Data Collection, Volume 1.5. U.S. Army Corps of Engineers, Omaha District, Yankton, SD.

APPENDICES

Appendix A. Phylogenetic list of Missouri River fishes with corresponding letter codes used in the long-term pallid sturgeon and associated fish community sampling program. The phylogeny follows that used by the American Fisheries Society, Common and Scientific Names of Fishes from the United States and Canada, 5th edition. Asterisks and bold type denote targeted native Missouri River species.

Scientific name	Common name	Letter Code
CLASS CEPHALASPIDOMORPHI-LAMPREYS		
ORDER PETROMYZONTIFORMES		
Petromyzontidae – lampreys		
<i>Ichthyomyzon castaneus</i>	Chestnut lamprey	CNLP
<i>Ichthyomyzon fossor</i>	Northern brook lamprey	NBLP
<i>Ichthyomyzon unicuspis</i>	Silver lamprey	SVLP
<i>Ichthyomyzon gagei</i>	Southern brook lamprey	SBLR
Petromyzontidae	Unidentified lamprey	ULY
Petromyzontidae larvae	Unidentified larval lamprey	LVLP
CLASS OSTEICHTHYES – BONY FISHES		
ORDER ACIPENSERIFORMES		
Acipenseridae – sturgeons		
<i>Acipenser fulvescens</i>	Lake sturgeon	LKSG
<i>Scaphirhynchus</i> spp.	Unidentified Scaphirhynchus	USG
<i>Scaphirhynchus albus</i>	Pallid sturgeon	PDSG*
<i>Scaphirhynchus platyrhynchus</i>	Shovelnose sturgeon	SNSG*
<i>S. albus</i> X <i>S. platyrhynchus</i>	Pallid-shovelnose hybrid	SNPD
Polyodontidae – paddlefishes		
<i>Polyodon spathula</i>	Paddlefish	PDFH
ORDER LEPISTOSTEIFORMES		
Lepisosteidae – gars		
<i>Lepisosteus oculatus</i>	Spotted gar	STGR
<i>Lepisosteus osseus</i>	Longnose gar	LNGR
<i>Lepisosteus platostomus</i>	Shortnose gar	SNGR
ORDER AMMIFORMES		
Amiidae – bowfins		
<i>Amia calva</i>	Bowfin	BWFN
ORDER OSTEOGLOSSIFORMES		
Hiodontidae – mooneyes		
<i>Hiodon alosoides</i>	Goldeye	GDEY
<i>Hiodon tergisus</i>	Mooneye	MNEY
ORDER ANGUILLIFORMES		
Anguillidae – freshwater eels		
<i>Anguilla rostrata</i>	American eel	AMEL

Appendix A. (continued).

Scientific name	Common name	Letter Code
ORDER CLUPEIFORMES		
Clupeidae – herrings		
<i>Alosa alabame</i>	Alabama shad	ALSD
<i>Alosa chrysochloris</i>	Skipjack herring	SJHR
<i>Alosa pseudoharengus</i>	Alewife	ALWF
<i>Dorosoma cepedianum</i>	Gizzard shad	GZSD
<i>Dorosoma petenense</i>	Threadfin shad	TFSD
<i>D. cepedianum</i> X <i>D. petenense</i>	Gizzard-threadfin shad hybrid	GSTS
ORDER CYPRINIFORMES		
Cyprinidae – carps and minnows		
<i>Campostoma anomalum</i>	Central stoneroller	CLSR
<i>Campostoma oligolepis</i>	Largescale stoneroller	LSSR
<i>Carassus auratus</i>	Goldfish	GDFH
<i>Carassus auratus</i> X <i>Cyprinus carpio</i>	Goldfish-Common carp hybrid	GFCC
<i>Couesius plumbeus</i>	Lake chub	LKCB
<i>Ctenopharyngodon idella</i>	Grass carp	GSCP
<i>Cyprinella lutrensis</i>	Red shiner	RDSN
<i>Cyprinella spiloptera</i>	Spotfin shiner	SFSN
<i>Cyprinus carpio</i>	Common carp	CARP
<i>Erimystax x-punctatus</i>	Gravel chub	GVCB
<i>Hybognathus argyritus</i>	Western silvery minnow	WSMN*
<i>Hybognathus hankinsoni</i>	Brassy minnow	BSMN
<i>Hybognathus nuchalis</i>	Mississippi silvery minnow	SVMW
<i>Hybognathus placitus</i>	Plains minnow	PNMW*
<i>Hybognathus</i> spp.	Unidentified <i>Hybognathus</i>	HBNS*
<i>Hypophthalmichthys molitrix</i>	Silver carp	SVCP
<i>Hypophthalmichthys nobilis</i>	Bighead carp	BHCP
<i>Luxilus chrysocephalus</i>	Striped shiner	SPSN
<i>Luxilus cornutus</i>	Common shiner	CMSN
<i>Luxilus zonatus</i>	Bleeding shiner	BDSN
<i>Lythrurus unbratilis</i>	Western redfin shiner	WRFS
<i>Macrhybopsis aestivalis</i>	Speckled chub	SKCB*
<i>Macrhybopsis gelida</i>	Sturgeon chub	SGCB*
<i>Macrhybopsis meeki</i>	Sicklefin chub	SFCB*
<i>Macrhybopsis storeriana</i>	Silver chub	SVCB
<i>M. aestivalis</i> X <i>M. gelida</i>	Speckled-Sturgeon chub hybrid	SPST
<i>M. gelida</i> X <i>M. meeki</i>	Sturgeon-Sicklefin chub hybrid	SCSC
<i>Macrhybopsis</i> spp.	Unidentified chub	UHY
<i>Margariscus margarita</i>	Pearl dace	PLDC
<i>Mylocheilus caurinus</i>	Peamouth	PEMT
<i>Nocomis biguttatus</i>	Hornyhead chub	HHCB
<i>Notemigonus crysoleucas</i>	Golden shiner	GDSN
<i>Notropis atherinoides</i>	Emerald shiner	ERSN
<i>Notropis blennioides</i>	River shiner	RVSN
<i>Notropis boops</i>	Bigeye shiner	BESN
<i>Notropis burchanani</i>	Ghost shiner	GTSN
<i>Notropis dorsalis</i>	Bigmouth shiner	BMSN
<i>Notropis greeni</i>	Wedgespot shiner	WSSN

Appendix A. (continued).

Scientific name	Common name	Letter Code
Cyprinidae – carps and minnows		
<i>Notropis heterolepsis</i>	Blacknose shiner	BNSN
<i>Notropis hudsonius</i>	Spottail shiner	STSN
<i>Notropis nubilus</i>	Ozark minnow	OZMW
<i>Notropis rubellus</i>	Rosyface shiner	RYSN
<i>Notropis shumardi</i>	Silverband shiner	SBSN
<i>Notropis stilbius</i>	Silverstripe shiner	SSPS
<i>Notropis stramineus</i>	Sand shiner	SNSN*
<i>Notropis topeka</i>	Topeka shiner	TPSN
<i>Notropis volucellus</i>	Mimic shiner	MMSN
<i>Notropis wickliffi</i>	Channel shiner	CNSN
<i>Notropis</i> spp.	Unidentified shiner	UNO
<i>Opsopoeodus emiliae</i>	Pugnose minnow	PNMW
<i>Phenacobius mirabilis</i>	Suckermouth minnow	SMMW
<i>Phoxinus eos</i>	Northern redbelly dace	NRBD
<i>Phoxinus erythrogaster</i>	Southern redbelly dace	SRBD
<i>Phoxinus neogaeus</i>	Finescale dace	FSDC
<i>Pimephales notatus</i>	Bluntnose minnow	BNMW
<i>Pimephales promelas</i>	Fathead minnow	FHMW
<i>Pimephales vigilas</i>	Bullhead minnow	BHMW
<i>Platygobio gracilis</i>	Flathead chub	FHCB
<i>P. gracilis</i> X <i>M. meeki</i>	Flathead-sicklefin chub hybrid	FCSC
<i>Rhinichthys atratulus</i>	Blacknose dace	BNDC
<i>Rhinichthys cataractae</i>	Longnose dace	LNDC
<i>Richardsonius balteatus</i>	Redside shiner	RDSS
<i>Scardinius erythrophthalmus</i>	Rudd	RUDD
<i>Semotilus atromaculatus</i>	Creek chub	CKCB
	Unidentified Cyprinidae	UCY
	Unidentified Asian Carp	UAC
Catostomidae - suckers		
<i>Carpionodes carpio</i>	River carpsucker	RVCS
<i>Carpionodes cyprinus</i>	Quillback	QLBK
<i>Carpionodes velifer</i>	Highfin carpsucker	HFCS
<i>Carpionodes</i> spp.	Unidentified Carpiodes	UCS
<i>Catostomus catostomus</i>	Longnose sucker	LNSK
<i>Catostomus commersoni</i>	White sucker	WTSK
<i>Catostomus platyrhincus</i>	Mountain sucker	MTSK
<i>Catostomus</i> spp.	Unidentified <i>Catostomus</i> spp.	UCA
<i>Cycleptus elongates</i>	Blue sucker	BUSK*
<i>Hypentelium nigricans</i>	Northern hog sucker	NHSC
<i>Ictiobus bubalus</i>	Smallmouth buffalo	SMBF
<i>Ictiobus cyprinellus</i>	Bigmouth buffalo	BMBF
<i>Ictiobus niger</i>	Black buffalo	BKBF
<i>Ictiobus</i> spp.	Unidentified buffalo	UBF
<i>Minytrema melanops</i>	Spotted sucker	SPSK
<i>Moxostoma anisurum</i>	Silver redhorse	SVRH
<i>Moxostoma carinatum</i>	River redhorse	RVRH
<i>Moxostoma duquesnei</i>	Black redhorse	BKRH
<i>Moxostoma erythrurum</i>	Golden redhorse	GDRH
<i>Moxostoma macrolepidotum</i>	Shorthead redhorse	SHRH
<i>Moxostoma</i> spp.	Unidentified redhorse	URH

Appendix A. (continued).

Scientific name	Common name	Letter Code
Catostomidae - suckers	Unidentified Catostomidae	UCT
ORDER SILURIFORMES		
Ictaluridae – bullhead catfishes		
<i>Ameiurus melas</i>	Black bullhead	BKBH
<i>Ameiurus natalis</i>	Yellow bullhead	YLBH
<i>Ameiurusnebulosus</i>	Brown bullhead	BRBH
<i>Ameiurus</i> spp.	Unidentified bullhead	UBH
<i>Ictalurus furcatus</i>	Blue catfish	BLCF
<i>Ictalurus punctatus</i>	Channel catfish	CNCF
<i>I. furcatus</i> X <i>I. punctatus</i>	Blue-channel catfish hybrid	BCCC
<i>Ictalurus</i> spp.	Unidentified <i>Ictalurus</i> spp.	UCF
<i>Noturus exilis</i>	Slender madtom	SDMT
<i>Noturus flavus</i>	Stonecat	STCT
<i>Noturus gyrinus</i>	Tadpole madtom	TPMT
<i>Noturus nocturnes</i>	Freckled madtom	FKMT
<i>Pylodictis olivaris</i>	Flathead catfish	FHCF
ORDER SALMONIFORMES		
Esocidae - pikes		
<i>Esox americanus vermiculatus</i>	Grass pickerel	GSPK
<i>Esox lucius</i>	Northern pike	NTPK
<i>Esox masquinongy</i>	Muskellunge	MSKG
<i>E. lucius</i> X <i>E. masquinongy</i>	Tiger Muskellunge	TGMG
Umbridae - mudminnows		
<i>Umbra limi</i>	Central mudminnow	MDMN
Osmeridae - smelts		
<i>Osmerus mordax</i>	Rainbow smelt	RBST
Salmonidae - trouts		
<i>Coregonus artedii</i>	Lake herring or cisco	CSCO
<i>Coregonus clupeaformis</i>	Lake whitefish	LKWF
<i>Oncorhynchus aguabonita</i>	Golden trout	GDTT
<i>Oncorhynchus clarki</i>	Cutthroat trout	CTTT
<i>Oncorhynchus kisutch</i>	Coho salmon	CHSM
<i>Oncorhynchus mykiss</i>	Rainbow trout	RBTT
<i>Oncorhynchus nerka</i>	Sockeye salmon	SESM
<i>Oncorhynchus tshawytscha</i>	Chinook salmon	CNSM
<i>Prosopium cylindraceum</i>	Bonniville cisco	BVSC
<i>Prosopium williamsoni</i>	Mountain whitefish	MTWF
<i>Salmo trutta</i>	Brown trout	BNTT
<i>Salvelinus fontinalis</i>	Brook trout	BKTT
<i>Salvelinus namaycush</i>	Lake trout	LKTT
<i>Thymallus arcticus</i>	Arctic grayling	AMGL

Appendix A. (continued).

Scientific name	Common name	Letter Code
ORDER PERCOPSIFORMES		
Percopsidae – trout-perches		
<i>Percopsis omiscomaycus</i>	Trout-perch	TTPH
ORDER GADIFORMES		
Gadidae - cods		
<i>Lota lota</i>	Burbot	BRBT
ORDER ATHERINIFORMES		
Cyprinodontidae - killifishes		
<i>Fundulus catenatus</i>	Northern studfish	NTSF
<i>Fundulus daphanus</i>	Banded killifish	BDKF
<i>Fundulus notatus</i>	Blackstripe topminnow	BSTM
<i>Fundulus olivaceus</i>	Blackspotted topminnow	BPTM
<i>Fundulus sciadicus</i>	Plains topminnow	PTMW
<i>Fundulus zebrinus</i>	Plains killifish	PKLF
Poeciliidae - livebearers		
<i>Gambusia affinis</i>	Western mosquitofish	MQTF
Atherinidae - silversides		
<i>Labidesthes sicculus</i>	Brook silverside	BKSS
ORDER GASTEROSTEIFORMES		
Gasterosteidae - sticklebacks		
<i>Culea inconstans</i>	Brook stickleback	BKSB
ORDER SCORPAENIFORMES		
Cottidae - sculpins		
<i>Cottus bairdi</i>	Mottled sculpin	MDSP
<i>Cottus carolinae</i>	Banded sculpin	BDSP
ORDER PERCIFORMES		
Percichthyidae – temperate basses		
<i>Morone Americana</i>	White perch	WTPH
<i>Morone chrysops</i>	White bass	WTBS
<i>Morone mississippiensis</i>	Yellow bass	YWBS
<i>Morone saxatilis</i>	Striped bass	SDBS
<i>M. saxatilis</i> X <i>M. chrysops</i>	Striped-white bass hybrid	SBWB
Centrarchidae - sunfishes		
<i>Ambloplites rupestris</i>	Rock bass	RKBS
<i>Archoplites interruptus</i>	Sacramento perch	SOPH
<i>Lepomis cyanellus</i>	Green sunfish	GNSF
<i>Lepomis gibbosus</i>	Pumpkinseed	PNSD
<i>Lepomis gulosus</i>	Warmouth	WRMH
<i>Lepomis humilis</i>	Orangespotted sunfish	OSSF
<i>Lepomis macrochirus</i>	Bluegill	BLGL
<i>Lepomis magalotis</i>	Longear sunfish	LESF
<i>Lepomis microlophus</i>	Redear sunfish	RESF
<i>L. cyanellus</i> X <i>L. macrochirus</i>	Green sunfish-bluegill hybrid	GSBG

Appendix A. (continued).

Scientific name	Common name	Letter Code
Centrarchidae - sunfishes		
<i>L. cyanellus</i> X <i>L. humilis</i>	Green-orangespotted sunfish hybrid	GSOS
<i>L. macrochirus</i> X <i>L. microlophus</i>	Bluegill-redear sunfish hybrid	BGRE
<i>Lepomis</i> spp.	Unidentified <i>Lepomis</i>	ULP
<i>Micropterus dolomieu</i>	Smallmouth bass	SMBS
<i>Micropterus punctatus</i>	Spotted sunfish	STBS
<i>Micropterus salmoides</i>	Largemouth bass	LMBS
<i>Micropterus</i> spp.	Unidentified <i>Micropterus</i> spp.	UMC
<i>Pomoxis annularis</i>	White crappie	WTCP
<i>Pomoxis nigromaculatus</i>	Black crappie	BKCP
<i>Pomoxis</i> spp.	Unidentified crappie	UCP
<i>P. annularis</i> X <i>P. nigromaculatus</i>	White-black crappie hybrid	WCBC
Centrarchidae	Unidentified centrarchid	UCN
Percidae - perches		
<i>Ammocrypta asprella</i>	Crystal darter	CLDR
<i>Etheostoma blennioides</i>	Greenside darter	GS DR
<i>Etheostoma caeruleum</i>	Rainbow darter	RBDR
<i>Etheostoma exile</i>	Iowa darter	IODR
<i>Etheostoma flabellare</i>	Fantail darter	FTDR
<i>Etheostoma gracile</i>	Slough darter	SLDR
<i>Etheostoma microperca</i>	Least darter	LTDR
<i>Etheostoma nigrum</i>	Johnny darter	JYDR
<i>Etheostoma punctulatum</i>	Stippled darter	STPD
<i>Etheostoma spectabile</i>	Orangethroated darter	OTDR
<i>Etheostoma tetrazonum</i>	Missouri saddled darter	MSDR
<i>Etheostoma zonale</i>	Banded darter	BDDR
<i>Etheostoma</i> spp.	Unidentified <i>Etheostoma</i> spp.	UET
<i>Perca flavescens</i>	Yellow perch	YWPH
<i>Percina caproides</i>	Logperch	LGPH
<i>Percina cymatotaenia</i>	Bluestripe darter	BTDR
<i>Percina evides</i>	Gilt darter	GLDR
<i>Percina maculate</i>	Blackside darter	BSDR
<i>Percina phoxocephala</i>	Slenderhead darter	SHDR
<i>Percina shumardi</i>	River darter	RRDR
<i>Percina</i> spp.	Unidentified <i>Percina</i> spp.	UPN
	Unidentified darter	UDR
<i>Sander canadense</i>	Sauger	SGER*
<i>Sander vitreus</i>	Walleye	WLEY
<i>S. canadense</i> X <i>S. vitreus</i>	Sauger-walley hybrid/Saugeye	SGWE
<i>Sander</i> spp.	Unidentified <i>Sander</i> (formerly <i>Stizostedion</i>) spp.	UST
	Unidentified Percidae	UPC
Sciaenidae - drums		
<i>Aplodinotus grunniens</i>	Freshwater drum	FWDM
NON-TAXONOMIC CATEGORIES		
	Age-0/Young-of-year fish	YOYF
	Lab fish for identification	LAB
	No fish caught	NFSH
	Unidentified larval fish	LVFS
	Unidentified	UNID
	Net Malfunction (Did Not Fish)	NDNF

Appendix B. Definitions and codes used to classify standard Missouri River habitats in the long-term pallid sturgeon and associated fish community sampling program. Three habitat scales were used in the hierarchical habitat classification system: Macrohabitats, Mesohabitats, and Microhabitats.

Habitat	Scale	Definition	Code
Braided channel	Macro	An area of the river that contains multiple smaller channels and is lacking a readily identifiable main channel (typically associated with unchannelized sections)	BRAD
Main channel cross over	Macro	The inflection point of the thalweg where the thalweg crosses from one concave side of the river to the other concave side of the river, (i.e., transition zone from one-bend to the next bend). The upstream CHXO for a respective bend is the one sampled.	CHXO
Tributary confluence	Macro	Area immediately downstream, extending up to one bend in length, from a junction of a large tributary and the main river where this tributary has influence on the physical features of the main river	CONF
Dendric	Macro	An area of the river where the river transitions from meandering or braided channel to more of a treelike pattern with multiple channels (typically associated with unchannelized sections)	DEND
Deranged	Macro	An area of the river where the river transitions from a series of multiple channels into a meandering or braided channel (typically associated with unchannelized sections)	DRNG
Main channel inside bend	Macro	The convex side of a river bend	ISB
Main channel outside bend	Macro	The concave side of a river bend	OSB
Secondary channel-connected large	Macro	A side channel, open on upstream and downstream ends, with less flow than the main channel, large indicates this habitat can be sampled with trammel nets and trawls based on width and/or depths > 1.2 m	SCCL
Secondary channel-connected small	Macro	A side channel, open on upstream and downstream ends, with less flow than the main channel, small indicates this habitat cannot be sampled with trammel nets and trawls based on width and/or on depths < 1.2 m	SCCS
Secondary channel-non-connected	Macro	A side channel that is blocked at one end	SCCN
Tributary	Macro	Any river or stream flowing in the Missouri River	TRIB
Tributary large mouth	Macro	Mouth of entering tributary whose mean annual discharge is > 20 m ³ /s, and the sample area extends 300 m into the tributary	TRML
Tributary small mouth	Macro	Mouth of entering tributary whose mean annual discharge is < 20 m ³ /s, mouth width is > 6 m wide and the sample area extends 300 m into the tributary	TRMS
Wild	Macro	All habitats not covered in the previous habitat descriptions	WILD
Bars	Meso	Sandbar or shallow bank-line areas with depth < 1.2 m	BARS
Pools	Meso	Areas immediately downstream from sandbars, dikes, snags, or other obstructions with a formed scour hole > 1.2 m	POOL
Channel border	Meso	Area in the channelized river between the toe and the thalweg, area in the unchannelized river between the toe and the maximum depth	CHNB
Dam Tailwaters	Meso	Area below dam	DTWT
Thalweg	Meso	Main channel between the channel borders conveying the majority of the flow	TLWG
Island tip	Meso	Area immediately downstream of a bar or island where two channels converge with water depths > 1.2 m	ITIP

Appendix C. List of standard and wild gears (type), their corresponding codes in the database, seasons deployed (Fall-Spring, Summer, or all), years used, and catch-per-unit-effort units for collection of Missouri River fishes in Segment 1 for the long-term pallid sturgeon and associated fish community sampling program. Long-term monitoring began in 2006 for Segment 1.

Gear	Code	Type	Season	Years	CPUE units
Gillnet – 4 meshes, small mesh set upstream	GN14	Wild	Sturgeon	NOT USED	fish/net night
Gillnet – 4 meshes, large mesh set upstream	GN41	Wild	Sturgeon	NOT USED	fish/net night
Gillnet – 8 meshes, small mesh set upstream	GN18	Wild	Sturgeon	NOT USED	fish/net night
Gillnet – 8 meshes, large mesh set upstream	GN81	Wild	Sturgeon	NOT USED	fish/net night
Mini-fyke net	MF	Standard	Fish Comm.	2006 - Present	fish/net night
Push Trawl – 8 ft 4mm x 4mm	POT02	Evaluation	Fish Comm.	2006 - 2008	fish/ 100 m trawled
Trammel net – 1 inch inner mesh	TN	Standard	All	2006 - Present	fish/100 m drift
Trot Line – Circle hooks**	TLC1	Experimental	Sturgeon	2007 - 2009	fish/hook night
Trot Line – Circle hooks**	TLC1	Standard	All	2010-present	fish/hook night
Trot Line – Octopus hooks**	TLO_	Wild	Sturgeon	NOT USED	fish/hook night
Trot Line – O'Shaughnessy hooks**	TLS_	Wild	Sturgeon	NOT USED	fish/hook night
Otter trawl – 16 ft head rope	OT16	Standard	All	2006 - Present	fish/100 m trawled
Otter trawl – 16 ft SKT 4mm x 4mm HB2 MOR	OT01	Wild	Fish Comm.	NOT USED	fish/100 m trawled

** Code ends with line length in feet (1 = 105 ft, 2 = 205 ft, 3 = 305 ft, 4 = 405 ft). Hooks are placed between 5 and 10 feet apart.

Appendix D. Stocking locations and codes by Recovery Priority Management Area (RPMA) in the Missouri River Basin.

State(s)	RPMA	Site Name	Code	River	RM
MT	2	Forsyth	FOR	Yellowstone	253.2
MT	2	Cartersville	CAR	Yellowstone	235.3
MT	2	Miles City	MIC	Yellowstone	181.8
MT	2	Fallon	FAL	Yellowstone	124
MT	2	Intake	INT	Yellowstone	70
MT	2	Sidney	SID	Yellowstone	31
MT	2	Big Sky Bend	BSB	Yellowstone	17
ND	2	Fairview	FRV	Yellowstone	9
MT	2	Milk River	MLK	Milk	11.5
MT	2	Mouth of Milk	MOM	Missouri	1761.5
MT	2	Grand Champs	GRC	Missouri	1741
MT	2	Wolf Point	WFP	Missouri	1701.5
MT	2	Poplar	POP	Missouri	1649.5
MT	2	Brockton	BRK	Missouri	1678
MT	2	Culbertson	CBS	Missouri	1621
MT	2	Nohly Bridge	NOB	Missouri	1590
ND	2	Confluence	CON	Missouri	1581.5
SD/NE	3	Sunshine Bottom	SUN	Missouri	866.2
SD/NE	3	Verdel Boat Ramp	VER	Missouri	855
SD/NE	3	Standing Bear Bridge	STB	Missouri	845
SD/NE	3	Running Water	RNW	Missouri	840.1
SD/NE	4	St. Helena	STH	Missouri	799
SD/NE	4	Mullberry Bend	MUL	Missouri	775
NE/IA	4	Ponca State Park	PSP	Missouri	753
NE/IA	4	Sioux City	SIO	Missouri	732.6
NE/IA	4	Sloan	SLN	Missouri	709
NE/IA	4	Decatur	DCT	Missouri	691
NE/IA	4	Boyer Chute	BYC	Missouri	637.4
NE/IA	4	Bellevue	BEL	Missouri	601.4
NE/IA	4	Rulo	RLO	Missouri	497.9
NE/MO/KS	4	Kansas River	KSR	Missouri	367.5
NE	4	Platte River	PLR	Platte	5
KA/MO	4	Leavenworth	LVW	Missouri	397
MO	4	Parkville	PKV	Missouri	377.5
MO	4	Kansas City	KAC	Missouri	342

State(s)	RPMA	Site Name	Code	River	RM
MO	4	Miami	MIA	Missouri	262.8
MO	4	Grand River	GDR	Missouri	250
MO	4	Boonville	BOO	Missouri	195.1
MO	4	Overton	OVT	Missouri	185.1
MO	4	Hartsburg	HAR	Missouri	160
MO	4	Jefferson City	JEF	Missouri	143.9
MO	4	Mokane	MOK	Missouri	124.7
MO	4	Hermann	HER	Missouri	97.6
MO	4	Washington	WAS	Missouri	68.5
MO	4	St. Charles	STC	Missouri	28.5

Appendix E. Juvenile pallid sturgeon stocking summary for Segment 2 of the Missouri River (RPMA 2)

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
1998	Big Sky Bend	255	1997	8/11/1998	Yearling	PIT Tag	Elastomer
1998	Confluence Nohly Bridge	40	1997	8/11/1998	Yearling	PIT Tag	Elastomer
1998	Sidney	255	1997	8/11/1998	Yearling	PIT Tag	Elastomer
1998	Sidney	230	1997	8/11/1998	Yearling	PIT Tag	Elastomer
2000	Culbertson	34	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Fairview	66	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Sidney	66	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Wolf Point	34	1998	10/11/2000	2 yr Old	PIT Tag	
2000	Culbertson	89	1999	10/17/2000	Yearling	PIT Tag	
2000	Fairview	150	1999	10/17/2000	Yearling	PIT Tag	
2000	Sidney	149	1999	10/17/2000	Yearling	PIT Tag	
2000	Wolf Point	90	1999	10/17/2000	Yearling	PIT Tag	
2002	Culbertson	270	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Fairview	270	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Intake	199	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Sidney	271	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Wolf Point	269	2001	7/18/2002	Yearling	CWT	Elastomer
2002	Culbertson	317	2001	7/26/2002	Yearling	PIT Tag	
2002	Fairview	360	2001	7/26/2002	Yearling	PIT Tag	
2002	Intake	97	2001	7/26/2002	Yearling	PIT Tag	
2002	Sidney	427	2001	7/26/2002	Yearling	PIT Tag	
2002	Wolf Point	425	2001	7/26/2002	Yearling	PIT Tag	
2002	Intake	155	2001	9/18/2002	Yearling	PIT Tag	
2003	Culbertson	1033	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2003	Fairview	887	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2003	Intake	1040	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2003	Wolf Point	926	2002	8/7/2003	Yearling	PIT Tag	Elastomer
2004	Milk River	821	2003	4/13/2004	Yearling	Elastomer	

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2004	Culbertson	523	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Intake	347	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Sidney	397	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Wolf Point	379	2003	8/9/2004	Yearling	PIT Tag	Elastomer
2004	Larval Drift	30000	2004	7/2/2004	Fry		
2004	Larval Drift	50000	2004	7/8/2004	Fry		
2004	Larval Drift	25000	2004	7/20/2004	Fry		
2004	Larval Drift	25000	2004	7/23/2004	Fry		
2004	Larval Drift	25000	2004	7/27/2004	Fry		
2004	Culbertson	3819	2004	9/10/2004	Fingerling	CWT	Elastomer
2004	Sidney	2991	2004	9/10/2004	Fingerling	CWT	Elastomer
2004	Wolf Point	4040	2004	9/10/2004	Fingerling	CWT	Elastomer
2004	Mouth of Milk	3482	2004	10/15/2004	Advanced Fingerling	CWT	Elastomer
2004	Intake	2477	2004	11/18/2004	Advanced Fingerling	CWT	Elastomer
2005	Culbertson	288	2004	4/12/2005	Yearling	CWT	Elastomer
2005	Intake	309	2004	4/12/2005	Yearling	CWT	Elastomer
2005	Wolf Point	271	2004	4/12/2005	Yearling	CWT	Elastomer
2005	Intake	175	2004	8/19/2005	Yearling	PIT Tag	Elastomer
2005	Brockton	229	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Culbertson	226	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Intake	456	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Milk River	232	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Sidney	122	2005	10/5/2005	Advanced Fingerling	CWT	Elastomer
2005	Wolf Point	611	2005	10/12/2005	Advanced Fingerling	CWT	Elastomer
2005	Brockton	371	2005	10/13/2005	Advanced Fingerling		
2005	Culbertson	1736	2005	10/13/2005	Advanced Fingerling	CWT	Elastomer
2005	Culbertson	182	2005	10/13/2005	Advanced Fingerling		

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2005	Intake	313	2005	10/13/2005	Advanced Fingerling		
2005	Milk River	845	2005	10/13/2005	Advanced Fingerling	CWT	Elastomer
2005	Mouth of Milk	371	2005	10/13/2005	Advanced Fingerling		
2005	Sidney	105	2005	10/13/2005	Advanced Fingerling		
2005	Wolf Point	1521	2005	10/13/2005	Advanced Fingerling	CWT	Elastomer
2005	Wolf Point	371	2005	10/13/2005	Advanced Fingerling		
2005	Culbertson	651	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Intake	2120	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Milk River	485	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Sidney	882	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2005	Wolf Point	650	2005	10/19/2005	Advanced Fingerling	CWT	Elastomer
2006	Culbertson	235	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Intake	327	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Mouth of Milk	134	2005	3/28/2006	Advanced fingerling	Elastomer	
2006	Sidney	113	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Wolf Point	232	2005	3/28/2006	Advanced Fingerling	Elastomer	
2006	Intake	970	2005	4/3/2006	Yearling	PIT Tag	Elastomer
2006	Sidney	314	2005	4/3/2006	Yearling	PIT Tag	Elastomer
2006	Culbertson	844	2005	4/5/2006	Yearling	PIT Tag	Elastomer
2006	Mouth of Milk	1007	2005	4/5/2006	Yearling	PIT Tag	Elastomer
2006	Wolf Point	866	2005	4/5/2006	Yearling	PIT Tag	Elastomer
2006	Culbertson	669	2005	5/1/2006	Yearling	PIT Tag	Scute Removed
2006	Intake	765	2005	5/1/2006	Yearling	PIT Tag	Scute

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
							Removed
2006	Mouth of Milk	650	2005	5/1/2006	Yearling	PIT Tag	Scute
2006	Sidney	228	2005	5/1/2006	Yearling	PIT Tag	Removed Scute
2006	Wolf Point	653	2005	5/1/2006	Yearling	PIT Tag	Removed Scute
2006		1355	2005	5/1/2006	Yearling	PIT Tag	Removed Scute Removed
2006	Culbertson	1544	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	Intake	1680	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	Mouth Milk	1117	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	Sidney	586	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	Wolf Point	1553	2006	10/24/2006	Advanced Fingerling	Elastomer	
2006	School Trust	436	2006	11/8/2006	Advanced Fingerling	Elastomer	
2007	Culbertson	651	2006	4/5/2007	Yearling	PIT Tag	Scute Removed
2007	Fallon	491	2006	4/3/2007	Yearling	PIT Tag	Scute Removed
2007	Forsyth	492	2006	4/3/2007	Yearling	PIT Tag	Scute Removed
2007	Sidney	983	2006	4/3/2007	Yearling	PIT Tag	Scute Removed
2007	School Trust	639	2006	4/5/2007	Yearling	PIT Tag	Scute Removed
2007	Wolf Point	651	2006	4/5/2007	Yearling	PIT Tag	Scute Removed
2007	Wolf Point	428285	2007	7/9/2007	Fry		
2007	Grand Champs	5558	2007	7/13/2007	Fry		
2007	Miles City	13125	2007	7/18/2007	Fry		
2007	Intake	20763	2007	8/9/2007	Fry		
2007	Miles City	13675	2007	8/9/2007	Fry		
2007	Intake	336	2007	8/27/2007	Fingerling		

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2007	Miles City	336	2007	8/27/2007	Fingerling		
2007	Wolf Point	672	2007	8/27/2007	Fingerling		
2007	Forsyth	690	2007	8/31/2007	Fingerling	CWT	
2007	Intake	615	2007	8/31/2007	Fingerling	CWT	
2007	School Trust	1160	2007	9/6/2007	Fingerling	CWT	
2007	Intake	293	2007	9/12/2007	Fingerling		
2007	Miles City	293	2007	9/12/2007	Fingerling		
2007	Wolf Point	586	2007	9/12/2007	Fingerling		
2007	Culbertson	6455	2007	9/14/2007	Fingerling	Elastomer	
2007	Fallon	4827	2007	9/14/2007	Fingerling	Elastomer	
2007	Forsyth	5370	2007	9/14/2007	Fingerling	Elastomer	
2007	Intake	7812	2007	9/14/2007	Fingerling	Elastomer	
2007	School Trust	6096	2007	9/14/2007	Fingerling	Elastomer	
2007	Sidney	1934	2007	9/14/2007	Fingerling	Elastomer	
2007	Wolf Point	6455	2007	9/14/2007	Fingerling	Elastomer	
2008	Culbertson	1384	2007	5/7/2008	Yearling	PIT Tag	Scute Removed
2008	Culbertson	643	2007	3/26/2008	Yearling	Elastomer	
2008	Fallon	1307	2007	5/7/2008	Yearling	PIT Tag	Scute Removed
2008	Forsyth	1384	2007	5/7/2008	Yearling	PIT Tag	Scute Removed
2008	Forsyth	106	2007	3/26/2008	Yearling	Elastomer	
2008	Intake	2395	2007	5/7/2008	Yearling	PIT Tag	Scute Removed
2008	Intake	103	2007	3/26/2008	Yearling	Elastomer	
2008	School Trust	1325	2007	5/7/2008	Yearling	PIT Tag	Scute Removed
2008	School Trust	654	2007	3/26/2008	Yearling	Elastomer	
2008	Sidney	149	2007	5/7/2008	Yearling	PIT Tag	Scute Removed
2008	Sidney	67	2007	3/26/2008	Yearling	Elastomer	
2008	Wolf Point	1328	2007	5/7/2008	Yearling	PIT Tag	Scute Removed

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2008	Wolf Point	416	2007	3/26/2008	Yearling	Elastomer	
2008	Miles City	4797	2008	7/30/2008	Fry		
2008	Grand Champs	24395	2008	7/30/2008	Fry		
2008	Culbertson	15630	2008	9/24/2008	Fingerling	Elastomer	
2008	Fallon	7930	2008	9/29/2008	Fingerling	Elastomer	
2008	Forsyth	7723	2008	9/29/2008	Fingerling	Elastomer	
2008	Intake	12642	2008	9/29/2008	Fingerling	Elastomer	
2008	Sidney	3186	2008	9/29/2008	Fingerling	Elastomer	
2008	Wolf Point	11717	2008	9/24/2008	Fingerling	Elastomer	
2009	Culbertson	1387	2008	4/13/2009	Yearling	PIT Tag	Scute Removed
2009	Fallon	1155	2008	4/13/2009	Yearling	PIT Tag	Scute Removed
2009	Forsyth	1166	2008	4/13/2009	Yearling	PIT Tag	Scute Removed
2009	Intake	2181	2008	4/13/2009	Yearling	PIT Tag	Scute Removed
2009	Sidney	710	2008	4/13/2009	Yearling	PIT Tag	Scute Removed
2009	Wolf Point	2162	2008	4/13/2009	Yearling	PIT Tag	Scute Removed
2009	Miles City	46260	2009	7/31/2009	Fry		
2009	Wolf Point	26175	2009	7/22/2009	Fry		
2009	Culbertson	10238	2009	9/24/2009	Fingerling	Elastomer	
2009	Fallon	5133	2009	9/23/2009	Fingerling	Elastomer	
2009	Forsyth	5386	2009	9/23/2009	Fingerling	Elastomer	
2009	Intake	8374	2009	9/23/2009	Fingerling	Elastomer	
2009	Sidney	1865	2009	9/23/2009	Fingerling	Elastomer	
2009	Wolf Point	9946	2009	9/23/2009	Fingerling	Elastomer	
2009	Intake	8374	2009	9/23/2009	Fingerling	Elastomer	
2009	Sidney	1865	2009	9/23/2009	Fingerling	Elastomer	
2009	Wolf Point	9946	2009	9/23/2009	Fingerling	Elastomer	
2010	Fallon	721	2009	4/15/2010	Yearling	PIT Tag	Scute Removed
2010	Fallon	268	2009	8/3/2010	Yearling	PIT Tag	Scute Removed

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2010	Fallon	1000	2010	10/7/2010	Fingerling	Elastomer	
2010	Forsyth	1402	2009	4/15/2010	Yearling	PIT Tag	Scute Removed
2010	Forsyth	268	2009	8/3/2010	Yearling	PIT Tag	Scute Removed
2010	Intake	1890	2009	4/15/2010	Yearling	PIT Tag	Scute Removed
2010	Intake	816	2009	6/4/2010	Yearling	Elastomer	
2010	Intake	541	2009	8/3/2010	Yearling	PIT Tag	Scute Removed
2010	Intake	1000	2010	10/7/2010	Fingerling	Elastomer	
2010	Sidney	331	2009	4/15/2010	Yearling	PIT Tag	Scute Removed
2010	Wolf Point	1309	2009	4/15/2010	Yearling	PIT Tag	Elastomer, Scute
2010	Wolf Point	858	2009	6/4/2010	Yearling	Elastomer	
2010	Wolf Point	425	2009	8/3/2010	Yearling	PIT Tag	Scute Removed
2010	Wolf Point	1000	2010	10/7/2010	Fingerling	Elastomer	
2010	Culbertson	65	2004	9/21/2010	6 Yr Old	PIT Tag	
2010	Culbertson	1337	2009	4/15/2010	Yearling	PIT Tag	Elastomer, Scute
2010	Culbertson	384	2009	6/4/2009	Yearling	PIT Tag	Scute Removed
2010	Culbertson	1000	2010	10/7/2010	Fingerling	Elastomer	
2010	School Trust	1766	2009	4/15/2010	Yearling	PIT Tag	Elastomer, Scute
2011	Culbertson	795	2010	5/5/2011	Yearling	PIT Tag	Scute
2011	Wolf Point	797	2010	5/5/2011	Yearling	PIT Tag	Scute
2011	Fallon	531	2010	5/5/2011	Yearling	PIT Tag	Scute
2011	Forsyth	545	2010	5/5/2011	Yearling	PIT Tag	Scute
2011	Intake	510	2010	5/5/2011	Yearling	PIT Tag	Scute
2011	Culbertson	262	2010	8/22/2011	Yearling	PIT Tag	Scute
2011	Fallon	131	2010	8/22/2011	Yearling	PIT Tag	Scute
2011	Forsyth	174	2010	8/22/2011	Yearling	PIT Tag	Scute
2011	Intake	132	2010	8/22/2011	Yearling	PIT Tag	Scute
2011	Wolf Point	262	2010	8/22/2011	Yearling	PIT Tag	Scute

Year	Stocking Site	Number Stocked	Year Class	Stock Date	Age at Stocking ^a	Primary Mark	Secondary Mark
2013	Wolf Point	187	2012	4/23/2013	Yearling	Pit Tag	Scute
2013	Culbertson	187	2012	4/23/2013	Yearling	Pit Tag	Scute
2013	Intake	118	2012	4/23/2013	Yearling	Pit Tag	Scute
2013	Fallon	185	2012	4/23/2013	Yearling	Pit Tag	Scute

^aAge of fish when stocked: Fry, Fingerling, Yearling, 1yo, 2yo, 3yo, etc...

Appendix G. Hatchery names, locations, and abbreviations.

Hatchery	State	Abbreviation
Blind Pony State Fish Hatchery	MO	BYP
Neosho National Fish Hatchery	MO	NEO
Gavins Point National Fish Hatchery	SD	GAV
Garrison Dam National Fish Hatchery	ND	GAR
Miles City State Fish Hatchery	MT	MCH
Blue Water State Fish Hatchery	MT	BLU
Bozeman Fish Technology Center	MT	BFT
Fort Peck State Fish Hatchery	MT	FPH