

**MONTANA DEPARTMENT OF FISH, WILDLIFE AND PARKS  
FISHERIES DIVISION  
JOB PROGRESS REPORT**

STATE: MONTANA      PROJECT TITLE: STATEWIDE FISHERIES INVESTIGATION  
PROJECT NO.: F-113-R-4      STUDY TITLE: SURVEY AND INVENTORY OF COLDWATER  
AND WARMWATER ECOSYSTEMS  
JOB NO.: V-e      JOB TITLE: NORTHEAST MONTANA WARMWATER  
ECOSYSTEMS INVESTIGATIONS  
JOB PERIOD: JULY 1, 2009 THROUGH JUNE 30, 2010

**TABLE OF CONTENTS**

<b>MONTANA DEPARTMENT OF FISH, WILDLIFE AND PARKS .....</b>	<b>1</b>
<b>ABSTRACT.....</b>	<b>2</b>
<b>PROCEDURES, RESULTS, &amp; DISCUSSION.....</b>	<b>2</b>
FORT PECK RESERVOIR AND UPPER MISSOURI RIVER PADDLEFISH STOCK.....	2
HILL COUNTY FISHING WATERS.....	134
BLAINE COUNTY PONDS .....	244
Phillips County Fishing Waters.....	31
<b>RECOMMENDATIONS.....</b>	<b>400</b>
PADDLEFISH: FORT PECK STOCK .....	400
FRESNO AND NELSON RESERVOIR .....	400
BEAVER CREEK RESERVOIR .....	400
HILL, BLAINE & PHILLIPS CO. PONDS .....	400
<b>REFERENCES .....</b>	<b>411</b>
<b>WATER CODES OF WATERS REFERRED TO.....</b>	<b>422</b>

## **ABSTRACT**

Paddlefish netting and tagging was conducted on the Missouri River paddlefish population upstream of Ft. Peck Reservoir, and tagging and harvest records were maintained. A native species creel survey was conducted on a 28-mile stretch of the Missouri River. The fourth year of a four-year telemetry study was conducted on paddlefish in the Missouri River and young-of-year paddlefish surveys were conducted in the headwaters of Ft. Peck Reservoir. Overall harvest on this population has been reduced in recent years, the average size of adult fish continues to decrease, and recruitment has been poor. As a result, regulations were changed in 2007 and were again restructured in 2008. Standardized gill netting and beach seining surveys were conducted at Fresno, Nelson, and Beaver Creek Reservoirs. Select ponds were sampled through Hill, Blaine, Phillips, Daniels, McCone, Richland, Roosevelt, Sheridan, and Valley Counties to assess fish population levels, survival and recruitment. Self-creel boxes were also maintained at select ponds in Hill, Blaine, and Phillips Counties to assess fishing pressure. Results of all other sampling are presented.

## **OBJECTIVES AND DEGREE OF ATTAINMENT**

Survey and Inventory- Objective is to survey and monitor the characteristics and trends of fish populations, angler harvest and preference, and to assess habitat conditions in selected waters. Objective accomplished, data presented.

Fish Population Management- Objective is to implement fish stocking programs and/or fish eradication actions to maintain fish populations at levels consistent with habitat conditions and other limiting factors. Objective accomplished, data presented.

Technical Guidance- To review projects by federal, state and local government agencies and private parties that have the potential to affect fisheries resources, and to provide technical advice or decisions to mitigate impacts on these resources. Provide landowners and other private parties with technical advice and information to sustain and enhance fisheries resources. Objective accomplished: ten 310 projects were reviewed and six 124 projects were reviewed with state and local agencies; attended six walleye unlimited meetings and helped with four school programs and fishing events related to the “Hooked on Fishing” program.

Angler Education- To enhance the public’s understanding, awareness and support of the state’s fishery and aquatic resources and to assist young people to develop angling skills and to appreciate the aquatic environment. Objective accomplished through staff participation in the “Hooked on Fishing” programs with local grade school children, planning and conducting of fall and winter fishing trips with area grade school and junior high children. Public presentations were also given on area fisheries in Havre and Malta. Staff also attended Walleye Unlimited meetings in Havre and Malta to provide information.

## **PROCEDURES, RESULTS, & DISCUSSION**

### **Fort Peck Reservoir and Upper Missouri River Paddlefish Stock**

The Ft. Peck stock was isolated from the Yellowstone-Sakakawea population upon closure of the Ft. Peck Dam in June of 1937. Completion of the Ft. Peck Dam isolated the Ft. Peck stock but it

also created productive rearing habitat. This resulted in increased numbers and size of paddlefish and in turn created a valuable recreational snag fishery in the Missouri River.

However, the alteration of flows from upstream dams combined with drought conditions in the Upper Missouri River and low reservoir levels have resulted in poor reproductive success of paddlefish over the last 10 to 15 years as well as reduced sizes and thus fecundity of females. In addition, the popularity of the fishery has increased dramatically during this time period bringing into question whether or not natural reproduction and recruitment is adequate for current harvest levels.

The goal of current management is to provide a stable recreational fishery while maintaining the population size and historical age structure of the spawning stock. To meet this goal, regulations were changed during the 2007-snagging season. New regulations limited harvest to one paddlefish per person and a choice where snaggers would fish (Missouri River above Ft. Peck Reservoir; Ft. Peck Dredge Cuts; lower Yellowstone River/Missouri River below Ft. Peck Reservoir). To distribute harvest and reduce size selective harvest, mandatory catch and release days (Sunday, Monday, & Thursday), and mandatory harvest days (Friday, Saturday, Tuesday, & Wednesday) were enacted. Additionally snagging was limited to 6 am to 9 pm to help reduce potential illegal take of paddlefish and make enforcement of the regulations more manageable.

Based on the results of the 2007 season, additional changes were made during the 2008 season. A quota of 500 fish was established, a season was set (May 1<sup>st</sup> to June 15<sup>th</sup>), hook size restriction were set, mandatory catch and release and harvest days were eliminated, and immediate release was further defined for paddlefish.

### **Data Collection Methods**

For more effective management of the Ft. Peck paddlefish stock, a thorough understanding of several key aspects of their life history was necessary. Data collected includes population size, harvest rates, spawning periodicity, age-structure, reproductive success, spawning locations, movements, and habitat use of paddlefish during their spawning migrations.

Baseline data on the paddlefish population above Ft. Peck Reservoir has been collected since the early 1970s. In 1993, a standardized monitoring program was established to assess population size, harvest rates, spawning periodicity, and gather baseline data on movement patterns and spawning sites. To gather this information, sampling occurs in the Missouri River during the time when paddlefish are staging around the Fred Robinson Bridge. Sampling typically occurs from April through May on the ascending arm of the hydrograph, typically at or above 8,000 cfs. Adult paddlefish are collected using drifted floating gill nets measuring 100 to 150 ft long, 6ft deep, with 4 inch mesh. Collected paddlefish are weighed, measured (eye-fork length), sexed, and tagged with an individually numbered jaw tag that is either a metal or plastic chicken leg band.

To assess angler pressure, catch and harvest rates of paddlefish, information on the ratio of tagged fish harvested, the size, sex, and age of fish harvested we conduct on-site roving creel surveys on a 28 mile stretch of the Missouri River downstream of the Fred Robinson Bridge. In addition a phone creel survey has been conducted since 2003 to assess harvest outside of the creel areas and as a check on the accuracy of the on-site creel surveys.

Beginning in 1996, concern over low flows and recruitment prompted the establishment of visual count surveys as a means of producing an annual index of recruitment of young of year (YOY) paddlefish. Visual counts have been found to be the most effective means to survey YOY paddlefish. Counts are conducted from an open bow powerboat using standardized methods and fixed transects. Observed YOY paddlefish are divided into age groups based on estimated length.

To increase our specific knowledge of the spawning locations, movements, and habitat use of paddlefish above Ft. Peck Reservoir, a contract with the University of Idaho was initiated in 2006 to conduct a four-year telemetry and egg sampling study. The telemetry study will continue through 2010.

## Adult Paddlefish Monitoring and Tagging

In 2009, paddlefish netting, tagging, and implantation of transmitters in paddlefish started on April 30<sup>th</sup>. Tagging efforts were continued until May 26<sup>th</sup> when we tagged our 324<sup>th</sup> paddlefish. Since tagging was initiated in 1977, 5,914 paddlefish have been tagged and 625-tagged paddlefish have been recaptured during annual gill netting surveys. Approximately 10.6 percent of the annual catch is comprised of recaptured fish. In 2009, 10% of the paddlefish captured were recaptured fish. Based on the tagging and recapture data, the reproductive periodicity of male paddlefish is one to two years and for females it is every two to three years.

Since tagging was initiated in 1977, a total of 687-tagged paddlefish have been reported as harvested, which is about 12% of all tagged paddlefish. While snaggers are encouraged to report catches of tagged fish, reporting rates are low in years when on-site creel surveys are not conducted. In 2009, 26-tagged paddlefish were reported as harvested and 3-tagged paddlefish were reported as snagged and released. Anglers harvested four paddlefish tagged in 2009.

## Preliminary population estimates

Estimates of population size of the recruited portion of the Ft. Peck stock were developed from 1993 through 2006 based on mark recapture sampling associated with gill netting and tagging conducted prior to and during the paddlefish snagging season. Point estimates and confidence intervals were developed using two approaches for estimate verification purposes: a modified Schnabel estimate and a modified Peterson estimate. Modifications, which allow for multiple years of marking and recapture data, were necessary because only a fraction of the recruited paddlefish stock matures and thus migrates upstream in a given year.

While the preliminary estimates were run for the entire tagging period, the most reliable Peterson estimates were developed from 1993 to 1999, when all five succeeding years of net catches could be used to assemble recaptures and high numbers of paddlefish were tagged during those years (Table 1). Based on the Peterson estimates the population size was approximately 20,500 fish from 1993 to 1999. The Schnabel estimate based on tagging over the period of 1993 to 1999 was 17,373 paddlefish with a 95% confidence interval of 15,614 to 20,336 fish. Both estimates indicate that the population size of recruited adult fish is approximately 20,000 fish. The number of adults that migrate in any given year is directly affected by the annual flows, however based on the periodicity of the fish, this means that approximately 11,700 fish migrate up the Missouri River to spawn and are “fishable” per year. As a comparison, the Yellowstone-Sakakawea stock has approximately 28,548 adults that migrate up the Yellowstone & Missouri Rivers (are “fishable”) in any given year and their total population size is approximately 53,017 (Scarnecchia 2006).

Table 1. - Population estimates (N-Hat) for paddlefish using the modified Peterson method with Ft. Peck tagging data from 1993 through 2001 where M is the number of new fish marked,  $C_{t+i}$  ( $i=1,2,3,4,5$ ) is the number of marked fish in year  $t+i$ ,  $C_R$  is the number of recaptures in years from  $t+1$  to  $t+5$  with fish tagged in other years, and R is the number of recaptures in years during  $t+1$  and  $t+5$  with fish tagged in year  $t$ .

Year	M	$C_{t+1}$	$C_{t+2}$	$C_{t+3}$	$C_{t+4}$	$C_{t+5}$	$C_R$	C_Sum	R	N_Hat	SD	95% Confidence Interval	
1993	405	500	456	281	501	368	56	2,162	46	19,035.00	7.7E-06	14,802.91	to 26,655.76
1994	500	546	280	501	368	355	106	2,156	48	22,458.33	6.4E-06	17,549.21	to 31,180.64
1995	456	281	501	368	355	88	131	1,724	38	20,688.00	7.8E-06	15,739.14	to 30,176.36
1996	281	501	368	355	88	13	121	1,446	15	27,088.40	9.5E-06	18,017.64	to 54,551.91
1997	501	368	355	88	13	221	118	1,163	30	19,422.10	9.3E-06	14,352.72	to 30,027.98
1998	368	355	88	13	221	259	124	1,060	20	19,504.00	1.1E-05	13,600.02	to 34,466.37
1999	355	88	13	221	259	240	106	927	21	15,670.71	1.4E-05	11,013.73	to 27,151.17
2000	88	13	221	259	240	323	114	1,170	4	20,609.60	1.9E-05	11,556.82	to 95,118.72
2001	13	221	259	240	323	500	141	1,684	1	10,952.50	4.6E-05	5,530.75	to 555,703.76

### ***Spawning and Recruitment***

The spawning success and recruitment rate of paddlefish is directly influenced by the magnitude, timing, and duration of peak flows. Berg (1981) postulated that a minimum flow of 14,000 cfs maintained for a period of 30 days is required to trigger paddlefish to move out of their staging areas and migrate upriver to spawning locations. This requirement has been observed in the Ft. Peck stock by monitoring flows and movement patterns, and comparing those to year class strength through aging, as well as with YOY sampling.

During the 1980s and 90s only 12 of the 20 years met the requirements necessary for successful migration and spawning (Figure 1 and 2). From 2000-2007, flows did not meet the minimum flow and duration requirements (Figure 3). However, these requirements were met in 2008 and 2009. In 2009, flows increased steadily to 10,000 cfs through April. Flow increased and peaked at 20,000 cfs on June 4<sup>th</sup> and declined to 10,000 cfs by early July. Peak flows met and exceeded trigger flows (14,000 cfs; Berg 1981) for about 35 days, slightly less than the 71-year average of 45 consecutive days (USGS 2009). The peak flow experienced in 2009 was the third highest in over a decade.

Hydrograph information (Figure 1, 2, and 3) suggests that good spawning conditions have not occurred since the late 1990s. Low recruitment has been verified by YOY surveys, which have been conducted annually since 1997 (Kozfkay & Scarnecchia 2002; Bowersox 2004; Miller 2005; Miller & Scarnecchia 2006). Good production of YOY paddlefish was observed in 1997 and 1998 when flows mimicked the historical hydrograph. Since that time less than five YOY paddlefish have been observed annually (Table 2). In 2009, no YOY or yearling paddlefish were observed during the fixed transects between RM 1844 and 1858 (Table 2). Additional surveys conducted outside of the transect area yielded better results. A total of 2 YOY and three yearling paddlefish were observed at RM 1859.

The effects of low flows and reduced recruitment is also starting to show up in the age distribution of harvested fish (Leslie 2006), which indicates that a strong year class has not been formed since the 1980s and early 1990s. However, since male paddlefish do not start spawning migrations until they are 10 and females when they are 15, the low recruitment levels over the last 10 to 15 years has not completely shown up in the harvest and population estimates yet.

Figure 1. - Historical and observed Missouri River hydrograph at the USGS Virgelle, MT gauging station 1980-1989.




Figure 2. - Historical and observed Missouri River hydrograph at the USGS Virgelle, MT gauging station 1990-1999.




Figure 3. - Historical and observed Missouri River hydrograph at the USGS Virgelle, MT gauging station 2000-2010.




Table 2. - Results of visual count surveys conducted in the upper end of Ft. Peck Reservoir from 1997 to 2009.

	<b>Transect Dates</b>	<b># Stations</b>	<b>Station Locations (RM)</b>	<b>No. Transects</b>	<b>No. YOY</b>	<b>No. Yearlings</b>	<b>Collector</b>
1997				69	113	3	
1998	7/27 to 9/23	8	1888 to 1866	216	97	54	Kozfkay
1999	8/25 to 9/20	8	1888 to 1866	174	3	10	Kozfkay
2000				90	0	11	
2001				90	1	0	
2002			1862 to 1856 ?				Bowersox ?
2003			1862 to 1856 ?	54	2	4	Bowersox ?
2004			1853 to 1838	54	0	3	
2005	8/8 & 8/16	6	1853 to 1838	36	1	0	Miller
2006	7/24 & 7/30	6	1853 to 1838	36	2	1	Miller
2007	7/31 & 8/6	6	1854 to 1838	6	0	2	Miller
2008	8/6 & 8/12	6	1844 to 1858	12	4	3	Miller
2009	8/11 & 8/12	6	1844 to 1858	12	0	0	Miller

## Harvest: Native Species Creel Survey 2009

### *Methods*

A five-year native species creel survey has been ongoing since 2005. In 2005, the roving creel survey was conducted by vehicle from April 1<sup>st</sup> to June 10<sup>th</sup> from the Fred Robinson Bridge to Peggy's Bottom. In 2006, the roving creel survey was repeated from April 3<sup>rd</sup> to June 18<sup>th</sup>. In 2007, the roving creel survey was conducted from April 1<sup>st</sup> to June 19<sup>th</sup>. In 2008, the creel was conducted by vehicle and boat from April 1<sup>st</sup>-June 15<sup>th</sup>. In 2009, the creel was conducted by vehicle from April 4<sup>th</sup> to June 15<sup>th</sup>.

### *Schedule*

A roving creel survey was conducted from April through June from the Fred Robinson Bridge to Peggy's Bottom. For the first half of April and the last half of June, fishing pressure is lower and one creel clerk was employed. During this time we used a stratified random sampling design to select survey dates from two strata, weekends and weekdays. All weekends were surveyed due to the high fishing pressure and two randomly selected weekdays were surveyed. From mid April to mid June, two creel clerks were employed and all days were surveyed. During high-pressure times, Memorial Day weekend, two extra creel clerks were employed and two teams surveyed anglers.

### *Effort*

Instantaneous pressure counts were performed once a day by vehicle due to travel distance and time. Starting times were randomly chosen from the fishing day (8:00 to 21:00 in 2005 & 2006) and counts were performed from ten pre-determined vantage points and access areas. Counts were performed at all sites except when road conditions prevented travel to some of the sites. In 2006, the accuracy of the vantage point counts were double check on three occasions by conducting simultaneous counts from vantage points and by boat for the entire creel area. In 2007, the creel survey and counts were conducted by boat for the entire creel area. In 2008, the creel survey and counts were conducted by boat and vehicle and in 2009 counts were conducted entirely by vehicle.

## ***Harvest and Catch***

Angler interviews were conducted at all boat ramps and fishing access/camping sites on the north side of the Missouri River between Fred Robinson Bridge (RM 1921) and lower Peggy's Bottom (RM 1987). Interviews were conducted at each site, during each sampling day except when road conditions prevented travel to certain sites. The creel clerks attempted to interview every fisherman on the river each sampling day. When the creel clerk encountered too many anglers at a location, the clerk systematically sub sampled every  $k$ th fisherman ( $k$  = interval demanded by the number of fishermen present) to ensure that all locations were sampled. Boat fishermen who launched within the area were interviewed even if they fished out of the study reach. Survey cards were distributed or deposited on windshields of vehicles when boat anglers were not available for interviews.

During interviews one randomly selected actively fishing person per party was interviewed. This individual was asked a series of questions relating to residence, number in party, length of stay, time spent fishing, whether a boat was used, type of gear used, and primary species being sought. Fishermen were asked how many of each species were caught, released and/or kept and if any of these fish had tags and if so the type, color, and number on the tag. When tagged paddlefish were caught, the fisherman was provided with the original tagging location, date, and weight of the fish. In addition, if fishermen released a paddlefish they were asked the reason for the release to determine if size-selective harvest was occurring.

When possible, harvested fish were measured to the nearest 0.1-inch and 0.01-pound. Eye-fork (body) length measurements were taken for paddlefish and fork-length measurements for sturgeon. Age samples consisting of jaws, spines, scales, or otoliths were collected from all harvested fish. These samples were then sent to the University of Idaho for aging.

In 2005 and 2006, the creel clerks also carried Passive Integrated Transponder (PIT) tag readers to identify any previously tagged pallid sturgeon and were prepared to PIT tag any new fish they encountered. Pamphlets and brochures developed for sturgeon, sauger, and paddlefish were distributed freely to fishermen.

## ***Data Analysis***

Fishing effort (angler-hours and total effort), catch rates, and harvest estimates were calculated using the Creel Census Program (McFarland and Roche 1987), which was developed using methods outlined in Neuhold and Lu (1957). To reduce error in 2005 and 2006, the creel survey time was broken down into three periods (April, May, and June), divided out by weekend and weekday effort, and by boat or shore fishing. In 2007, the analysis for all species except paddlefish was conducted as in 2005 and 2006. In 2007, paddlefish regulations were changed to include mandatory harvest and mandatory release days. As a result, the creel survey time for paddlefish was broken down into three periods (April, May, and June), divided out by harvest and release day effort, and by boat or shore fishing. In 2008 and 2009, paddlefish regulations changed again and the creel analysis was conducted similar to 2005 and 2006. Effort was not broken down by fishing method or by species because anglers were usually fishing for more than one species during a day.

## ***Results***

In 2009, a total of 536 parties from 10 states and 32 of the 56 Montana counties were interviewed from April 1<sup>st</sup> to June 15<sup>th</sup>. In 2009, the highest percentage of anglers in Montana came from Yellowstone (16.7%), Fergus (13.6%), Gallatin (13.2%), and Cascade (10.8%) Counties. In 2009, the average party contained 4.1 anglers (range = 1 to 22 anglers), 82% of which were males. In 2009, the average length of stay was 3.4 day/trip (range = 1 to 16 days).



### ***Effort***

In 2009, angler-hours were similar to 2008 at 28,277 (Table 3). In 2009, 81.6% of the angling effort occurred from shore. In 2009, 77.7% of the angling effort occurred in May, which coincides with the peak of the paddlefish spawning migration.

### ***Species caught and released***

In 2009, a total of 9 species were caught and/or harvested from April through June. In 2009, channel catfish, paddlefish, sauger, walleye, and drum were the most common species caught by anglers (Table 5). Channel catfish, paddlefish, sauger, and walleye were the most common species harvested (Table 5). Average length, weight, and condition of harvested fish measured by the creel clerks in 2009 are presented in Table 5 respectively.

### ***Paddlefish***

In 2009, a total of 881 paddlefish were caught and reported to creel clerks. The catch rate was 0.161 paddlefish/hour (Table 4).

In 2009, harvested paddlefish ranged in length from 24.0 to 54.0 inches (eye-fork length) and ranged in weight from 16.0 to 100 pounds (Table 8). Fifty-two percent of the harvested paddlefish sexed were males. And 26 (10.5 %) of the harvested paddlefish and 3 (1.0 %) of the released paddlefish creel had jaw tags. Harvested paddlefish ranged in age from 9 to 54 years with 48% of the harvested fish being 20 to 56 years old. Twenty-six percent of the harvested fish were less than 16 years old (new recruits).

### ***Channel Catfish***

In 2009, a total of 403 channel catfish were caught and reported to creel clerks. The catch rate was 0.115 channel catfish/angler-hour (Table 4). Eighty-two percent of the channel catfish caught were harvested. In 2009, harvested channel catfish ranged in length from 10.5 to 37.8 inches and in weight from 0.25 to 29 pounds (Table 5). Seventy percent of the harvested channel catfish sexed were females. Harvested channel catfish ranged in age from 3 to 18 years old.

### ***Stonecats***

In 2009, 10 stonecats were caught and reported to the creel clerks. The catch rate was 0.003 stonecats/angler-hour (Table 4). No stonecats caught by the anglers interviewed were harvested.

### ***Sauger***

In 2009, 78 sauger were caught and reported to the creel clerks. The catch rate was 0.018 sauger/angler-hour (Table 4). Anglers harvested 68% of the sauger caught, which ranged in length from 12.5 to 28.0 inches and in weight from 0.75 to 7.0 pounds (Table 5).

### ***Walleye***

In 2009, a total of 104 walleye were caught and reported to the creel clerks. The catch rate was 0.032 walleye/angler-hour (Table 4). Anglers harvested 86% of the walleye they caught, which ranged in length from 12.0 to 29.5 inches and in weight from 0.39 to 7.75 pounds (Table 8).

### ***Pallid Sturgeon***

In 2009, 11 pallid sturgeon were caught and reported to the creel clerks. The catch rate was 0.0004 pallid sturgeon/angler-hour.

### *Shovelnose Sturgeon*

In 2009, a total of 12 shovelnose sturgeon were caught and reported to the creel clerks. The catch rate was 0.0004 shovelnose sturgeon/angler-hour (Table 4).

In 2009, one shovelnose sturgeon was harvested. The length was 32 inches and it weighed 5 pounds (Table 5).

### *Burbot*

In 2009, a total of 2 burbot caught and reported to the creel clerks. The catch rate was 0.0005 burbot/angler-hour (Table 4). Anglers harvested 50% of the burbot caught and ranged in length from 19 to 27 inches and in weight from 1.5 to 2.5 pounds (Table 5).

### *Freshwater Drum*

In 2009 a total of 66 drum were caught and reported to the creel clerks. The catch rate was 0.018 drum/angler-hour (Table 4). Anglers harvested 70% of the drum caught and ranged in length from 9.3 to 15.2 inches and in weight from 0.22 to 1.25 pounds (Table 8).

### *Goldeye*

In 2009, 56 goldeye were caught and reported to the creel clerks. The catch rate was 0.011 goldeye/angler-hour (Table 4). Anglers harvested 27% of the goldeye caught and ranged in length from 11.5 to 13.5 (Table 5).

### *Other Species*

Other native species caught included, bigmouth buffalo, river carpsucker, and shorthead redhorse. The majority of these species were released. Other non-native species caught included common carp. The specific length, weight, and structural indices for these species are reported in Table 5.

Table 3. - Estimated fishing effort (angler-hours) and standard error (SE) by month and angler type for the Missouri River (RM 1899 to 1921), April-June 2009.

	<b>Shore</b>	<b>SE</b>	<b>Boat</b>	<b>SE</b>	<b>Overall</b>
<b>April</b>	1,965.60	550.82	588.00	225.97	2,553.60
<b>May</b>	17,531.03	3,464.95	4,378.48	897.05	21,909.51
<b>June</b>	3,565.72	922.50	137.15	65.19	3,702.87
<b>Overall</b>	23,062.35		5,103.63		28,165.98

Table 4. - Estimated hourly catch rate (fish/angler-hour) by strata channel catfish (C CAT), stonecat (S CAT), sauger (SGR), walleye (WE), pallid sturgeon (PSTG), shovelnose sturgeon (SSTG), burbot (LING), goldeye (GE), freshwater drum (DRUM), northern pike (NP), paddlefish (PF), and other species (Other) in the Missouri River (RM 1899 to 1921), April-June 2009.

<b>Overall</b>						
	<b>Shore</b>	<b>SE</b>	<b>Boat</b>	<b>SE</b>	<b>Overall</b>	<b>SE</b>
C CAT	0.0655	0.0045	0.2310	0.0105	0.1153	0.0043
S CAT	0.0029	0.0017	0.0050	0.0040	0.0034	0.0016
SGR	0.0195	0.0044	0.0163	0.0488	0.0186	0.0128
WE	0.0315	0.0018	0.0349	0.0089	0.0326	0.0026
PSTG	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
SSTG	0.0000	0.0000	0.0014	0.0010	0.0004	0.0005
LING	0.0007	0.0004	0.0000	0.0000	0.0005	0.0003
GE	0.0146	0.0025	0.0014	0.0036	0.0106	0.0021
DRUM	0.0188	0.0007	0.0175	0.0005	0.0185	0.0005
NP	0.0000	0.0000	0.0000	0.0000	0.0000	0.0000
PF	0.1399	0.0026	0.2083	0.0037	0.1605	0.0022
Other	0.0013	0.0005	0.0013	0.0005	0.0013	0.0005

Table 5. – Length, weight, and condition indices of harvested paddlefish (PF), channel catfish (C CAT), sauger (SGR), walleye (WE), shovelnose sturgeon (SSTG), burbot (LING), freshwater drum (DRUM), goldeye (GE), and common carp (CARP) from anglers creeled in the middle Missouri River native species creel, April-June 2009.

<b>Species</b>	<b>Sample Size</b>	<b>Length Range (in.)</b>	<b>Length Avg</b>	<b>Length SD</b>	<b>Weight Range (lbs.)</b>	<b>Weight Avg</b>	<b>Weight SD</b>
PF	249	24.0-54	41.7	5.9	16.0-100	47.6	21.2
C CAT	163	10.5-37.8	24.8	7.4	0.25-29.0	7.7	6.7
SGR	29	12.5-28.0	17.6	3.0	0.75-7.0	1.8	1.2
WE	41	12.0-29.5	16.7	4.1	0.39-7.75	1.9	1.7
S STG	1	32.00	32.0	0.0	5.0	5.0	0.0
LING	2	19.0-27.0	23.0	5.7	1.5-2.5	2.0	0.7
DRUM	10	9.3-15.2	12.1	1.8	0.22-1.25	0.6	0.3
GE	9	11.5-13.5	12.4	0.7	0.44-0.79	0.5	0.1
CARP	2	23.0-24.0	23.5	0.7	4.00-5.00	4.5	0.7

#### ***Paddlefish Phone Creel (2003-2009)***

Vic Riggs with the Montana Fish, Wildlife & Parks (FWP) and Larry Brooks with the University of North Dakota (Riggs 2005) designed and conducted the paddlefish phone creel survey annually from 2003 to 2005. This survey was continued solely by FWP in 2006. This survey was originally conducted (1) to determine the harvest of paddlefish at sites other than the Intake Fishing Access Site, (2) as a check on the accuracy of the Intake creel survey, (3) as a possible replacement for the Intake creel survey, which would free up technician time for other data collection needs, and (4) to obtain harvest statistics for the Fort Peck population, and in 2005 and 2006, (5) to assess angler support for changes to regulations for the Fort Peck populations.

From 2003 to 2009, harvest statistics were obtained for the Fort Peck population (Table 9). In 2005, two questions were added to determine why anglers were releasing paddlefish, and to determine the support for reducing the snagging limit to one paddlefish and the possible removal of catch and release fishing.

On average approximately 2,260 anglers fish for paddlefish above Ft. Peck Reservoir representing approximately 6,342 fishing days. On average approximately 1,613 paddlefish are caught annually above Ft. Peck Reservoir (Table 6). Approximately 50% of the paddlefish caught are released. When anglers were asked why they released their paddlefish, 53% said they were catch and release fishing, either because they did not want to harvest a paddlefish or because they wanted to fish for their entire fishing trip and harvest a fish on one of their last days. Seventeen percent of paddlefish were released because they were too big, indicating that these people were releasing the larger females. Twenty-seven percent of paddlefish were released because they were too small, indicating that these people are high grading for the larger females and are trophy fishing.

Table 6. –Summary of estimates for the Fort Peck paddlefish population from the Montana paddlefish telephone creel survey (2003-2009).

<b>Missouri River Above Fort Peck</b>							
	<b>2003</b>	<b>2004</b>	<b>2005</b>	<b>2006</b>	<b>2007</b>	<b>2008</b>	<b>2009</b>
<b>Number of Anglers</b>	1,902	2,859	2,705	2,476	2,481	1,816	1,579
<b>Total Days Fished*</b>	5,757	9,172	8,385	7,565	-	4,426	2,748
<b>Total Hours Fished*</b>	27,433	44,400	42,277	39,800	-	-	-
<b>Number Caught</b>	1,583	1,102	1,516	2,290	-	845	2,342
<b>Number Harvested</b>	868	787	1,028	1,067	634	300	564
<b>Harvest Rate (fish/day)</b>	0.151	0.086	0.123	0.141	-	0.068	0.205
<b>Harvest Rate (fish/hour)</b>	0.032	0.018	0.024	0.027	-	-	-
<b>Percent Released</b>	45.17%	28.58%	32.19%	53.42%	-	64.50%	75.90%
<b>Percent Contacted by FWP Creel Clerk</b>						85.71%	62.14%

\* Includes hours spent catch and release fishing

### Discussion

The paddlefish population above Ft. Peck Reservoir has experienced declines in recruitment, growth, and overall size of adults. Flows in the Missouri River have not been consistently high enough to produce large year classes of paddlefish for the past 10 to 15 years (Leslie 2005, 2006). In addition, the average size of adults has significantly decreased over the last 30 years (Bowersox 2004). These declines, especially in growth, are believed to be the result of decreased productivity due to the aging of Ft. Peck Reservoir (nursery grounds for paddlefish) and lower Ft. Peck Reservoir levels. Paddlefish are not experiencing as high of growth rates while residing in Ft. Peck Reservoir and as a result sexually mature fish are of a smaller size than they were when Ft. Peck Reservoir was initially formed. In addition, natural mortality and fishing mortality are resulting in the decreased proportion of large fish, specifically females. Fishing pressure has nearly doubled from 1977 to 2009, and high grading allows fishermen to increase their chances of catching a fish of a desirable size. The decreased weight of adult females is of concern because it is directly related to fecundity.

The combination of a low number of successful spawning years (based on observed trigger flow occurrence and duration) and decreased size of adults is of concern. The results of reduced recruitment of YOY paddlefish will not be evident in the spawning population for a few years, however it would be prudent to consider the effects of reduced recruitment and reduced fecundity of the adult population. Since 1992, snaggers have been allowed to harvest two paddlefish per year from the upper Missouri River fishery. Snaggers are also allowed to immediately release a snagged paddlefish if they desire. Based on analysis of the fishery and public support, the fishing limit was reduced in 2007 from two paddlefish to one paddlefish annually. In 2008, the paddlefish season was reduced from a 365-day season to a 45-day season (May 1 to June 15), making monitoring total catch more feasible. Furthermore, an annual harvest cap of 500 paddlefish has been implemented since 2008.

## **Hill County Fishing Waters**

Select waters throughout Hill County were sampled to determine fish abundance using sinking multi-filament experimental gill nets measuring 125 feet in length and 6 feet deep, consisting of 25-foot panels of ¾", 1", 1 ¼", 1 ½", and 2" mesh unless otherwise specified. Voluntary creel boxes were maintained at many of the ponds to determine fishing pressure, catch rates, and satisfaction.

### **Bailey Reservoir**

Bailey Reservoir is a privately owned reservoir that was constructed in the mid-1960s. At full pool, Bailey Reservoir covers approximately 70 surface acres and has a maximum depth of 28 feet. Bailey ranked 52<sup>nd</sup> in the region for angler pressure in 2005/2006 (123 angler days; McFarland 2006). Bailey was initially managed as a rainbow trout fishery, and rainbow trout thrived within the reservoir until 1980 when northern pike were illegally introduced. In 1984, the remainder of rainbow trout winterkilled due to severe drought. Chemical rehabilitation was considered, but at the request of the landowner a cool/warm water fishery was started. And as a result, yellow perch and black crappie were introduced in 1987, largemouth bass in 1988, and walleye in 1989. Rainbow trout are stocked periodically to supplement the fishery and in 2005, 10,000 four-inch rainbow trout were stocked in late fall. In addition, 10,000 fingerling walleye were stocked in the spring of 2005 and 261 yellow perch were transplanted from the Havre Water District Ponds in Kremlin. In 2006, an additional 4,758 pre-spawn yellow perch were transplanted. Supplement plants of 3,880, 850, and 1,500 yellow perch occurred in 2007, 2008, and 2009.

Adult sport fish populations have been monitored since 1990 with two experimental gill net sets. In addition, trap netting and electrofishing occurs periodically. In 2005, 6 traps nets were set in the spring to capture adult black crappie for transporting to Home Run Pond. From 2005 to 2007, two overnight gillnet sets were also used to monitor adult fish populations. In addition a voluntary creel box was erected in the summer of 2005 and maintained through 2007 to determine angler use, catch, and satisfaction.

In 2009, gill net surveys suggest the population of all sport fish (walleye, northern pike, yellow perch, black crappie) remain well below population densities prior to 2000 (pre-drought years; Table 7). From 1999-2007 Bailey's Reservoir was used as donor source for adult yellow perch and black crappie. The extensive removal of spawning adult fish combined with low reservoir levels, and less than ideal spawning conditions could be the most likely explanation for population densities we have today.

Table 7. - Catch rate (CPUE (fish/net)), average length (TL, in.), and average weight (lb.) of northern pike, yellow perch, black crappie, rainbow trout, walleye, and white sucker in Bailey Reservoir, 1985 to 2009.

Year	Nets	Northern pike			Yellow Perch			Black Crappie			Rainbow Trout			Walleye			White Sucker		
		CPUE	Len	Wt	CPUE	Len	Wt	CPUE	Len	Wt	CPUE	Len	Wt	CPUE	Len	Wt	CPUE	Len Avg	Wt Avg
			Avg	Avg		Avg	Avg		Avg	Avg		Avg	Avg		Avg	Avg		(in.)	(lbs.)
		(#/net)	(in.)	(lbs.)	(#/net)	(in.)	(lbs.)	(#/net)	(in.)	(lbs.)	(#/net)	(in.)	(lbs.)	(#/net)	(in.)	(lbs.)	(#/net)	(in.)	(lbs.)
1985	1	17	21.44	1.13	0	--	--	0	--	--	1	12.2	0.9	--	--	--	0	--	--
1990	3	8	18.1	1.23	11.33	7.7	0.26	7	5.7	0.1	0	--	--	--	--	--	0	--	--
1991	2	3.5	24.7	3.21	29	10.1	0.56	2	8.5	0.35	0	--	--	--	--	--	0	--	--
1992	2	3	26.8	4.29	17	8.1	0.29	8	4.7	0.08	0	--	--	--	--	--	0	--	--
1993	2	1	31.8	7.55	10.5	6.6	0.15	63.5	6.7	0.12	0	--	--	--	--	--	0	--	--
1994	2	3.5	20.1	2.59	19	6	0.1	21.5	6.3	0.14	0	--	--	--	--	--	0	--	--
1995											No Netting Conducted								
1996	2	7	23.8	3.54	43	7.2	0.19	7.5	6.8	0.21	0	--	--	--	--	--	0	--	--
1997											No Netting Conducted								
1998	2	1.5	22.2	2.43	66	8	0.26	16	9	0.44	0	--	--	--	--	--	0	--	--
1999											No Netting Conducted								
2000											No Netting Conducted								
2001											No Netting Conducted								
2002	2	0	0	0	16	9.9	0.49	15.5	11.2	0.82	0	--	--	1	25.7	6.79	1	17.9	2.41
2003											No Netting Conducted								
2004											No Netting Conducted								
2005	2	3.5	17.44	1.56	1.5	9.2	0.39	1	4.05	0.03	0	--	--	--	--	--	0	--	--
2006	2	16	17.23	1.2	3.5	7.29	0.28	0	--	--	0	--	--	6.5	9.54	0.31	0	--	--
2007	2	5.5	20.8	2.05	0.5	11.3	0.9	0	--	--	0	--	--	3	12.5	0.65	0	--	--
2008											No Netting Conducted								
2009	2	2	20.6	1.97	1	13	1.38	0	--	--	0	--	--	2	18.2	2.28	1	19	3.07

### Beaver Creek Reservoir

Beaver Creek Reservoir, located south of Havre, is a 200-acre reservoir, which has a maximum depth of 90 feet. Its proximity to the city of Havre makes this reservoir a valuable local resource and it has been managed intensively in recent years for a variety of species. The statewide fishing pressure survey for 2005/2006 indicated it was the fifth most fished reservoir in Region Six (McFarland 2006).

This reservoir was established as a rainbow trout fishery in 1975. However, the illegal introduction of northern pike (1980s) and yellow perch (1980s) has resulted in the rainbow trout fishery having varying success. As a result, the fisheries management plan was expanded to include other warm water species, which were introduced to control undesirable species and enhance the fishing opportunity within the reservoir. Currently this reservoir receives annual plants of 70,000 catchable size Eagle Lake, Erwin and Arlee rainbow trout.

In an effort to maintain the balance between the rainbow trout fishery and the warm water fishery, the use of live minnows for bait has been allowed since March of 2000. The regulation is intended to increase harvest of northern pike and perhaps open up a winter fishery for walleye. Though fishermen use live minnows regularly, a winter fishery for walleye has not developed. The trout daily limit was reduced from 5/day to 3/day in March of 2002 due to increasing fishing pressure.

### Population Status of Adult and Young-of-Year Fishes

Adult fish populations were monitored at six fixed experimental gillnetting stations, which were established in 1986. Gillnetting was conducted over night utilizing three sinking and three floating experimental gill nets (6 net-days). The sinking and floating experimental gill nets measured 125 feet in length and 6 feet deep

consisting of 25-foot panels of  $\frac{3}{4}$ ", 1", 1  $\frac{1}{4}$ ", 1  $\frac{1}{2}$ ", and 2" mesh. Fish were measured for total length (TL: inches) and weighed to the nearest 0.01 pound (lb). Prior to 1986, adult fish populations were monitored, however sampling was neither uniform, nor consistent enough to develop useful trend data on game fish population size, or composition. As a result this data was excluded from analysis and is only included within the tables for reference to the illegal introduction of northern pike and yellow perch.

The abundance and reproductive success of sport and forage fishes were monitored at six predetermined stations. Beach seining was conducted in early August using a 100- x 9-foot x  $\frac{1}{4}$  inch square mesh beach seine. The fish were sorted by species and counted.

### ***Northern pike***

Since their illegal introduction in the 1980s, northern pike abundance has maintained manageable levels within Beaver Creek Reservoir (Table 8). Northern pike populations are cyclical within Beaver Creek Reservoir, YOY catch had increased significantly in 2005 and 2006 and these fish have successfully been recruited into the population (Table 8 & 9). In 2007, the YOY catch was greatly reduced and no YOY northern pike were observed in 2008. Spring spawning conditions were favorable in 2009 and YOY catch of northern pike were similar to historic averages (Table 9).

### ***Yellow perch***

Yellow perch were illegally introduced into Beaver Creek Reservoir in 1987. Since their introduction, yellow perch have thrived within the reservoir (Table 8). As a result Beaver Creek Reservoir is a popular ice fishing destination and has historically been a source of yellow perch for kids fishing ponds, such as Home Run Pond, in Glasgow.

Beaver Creek Reservoir's yellow perch population peaked in the late 1990s and while the relative abundance is good, it is currently at half the late 1990 levels due to drought and reduced spawning success (Table 8 & 9). While yellow perch populations remain low compared to 1990 levels they are steadily increasing. The population still consists of numerous quality and preferred size fish (TL > 8 in.; Table 8). Yellow perch experienced higher spawning success in 2004, 2006, and 2009 as a result of increased water levels and flooded vegetation, (Table 9). Spawning success was reduced in 2007 and 2008 (Table 9) however the population of adult fish continues to sustain a high density (Table 8).

### ***Walleye***

Walleye were initially stocked in 1987 to provide a greater diversity of fishing opportunities within the reservoir. Natural reproduction is very limited within the reservoir and as a result, approximately 10,000 1.4-inch and 5,000 3-inch walleye fingerlings are stocked annually. In 2009, 7,000 4-inch walleye fingerlings were stocked.

Walleye within Beaver Creek Reservoir have high growth rates and relative weights (Table 8) due to abundant forage. As a result these walleye can be hard to catch and elude all but the best walleye fishermen. Since their initial introduction, high quality walleye have thrived within Beaver Creek Reservoir. The average size of walleye in 2009 was 18.26 in. with high condition factors (Table 8).

### ***Smallmouth bass***

Smallmouth bass were first introduced in 1997 and were stocked annually until 2000. As a result of these efforts there is now a self-sustaining population of smallmouth bass in Beaver Creek Reservoir. While smallmouth bass had a low relative abundance during gill netting surveys (Table 8), due to selectivity of the gear, catches of 8 to 10 inch bass by anglers are common. In addition, several YOY were collected during summer seining surveys (Table 9).



Table 11.- Summary of relative abundance (catch per unit effort (CPUE)), average total length, and relative weights of fishes collected in fall gillnetting surveys in Beaver Creek Reservoir, 1974-2009.

		Rainbow Trout				Yellow Perch			Northern Pike			Smallmouth bass			Walleye			Longnose sucker		White sucker	
		CPUE	Ave TL			CPUE	Ave TL		CPUE	Ave TL		CPUE	Ave TL		CPUE	Ave TL		CPUE	Ave TL	CPUE	Ave TL
Date		Nets	(fish/net)	(in.)	Rel Wt	(fish/net)	(in.)	Rel Wt	(fish/net)	(in.)	Rel Wt	(fish/net)	(in.)	Rel Wt	(fish/net)	(in.)	Rel Wt	(fish/net)	(in.)	(fish/net)	(in.)
Sep-74	1974	3	24.00	10.91	111.26	--	--	--	--	--	--	--	--	--	--	--	--	7.33	10.49	82.33	10.23
Nov-77	1977	3	35.00	10.05	86.31	--	--	--	--	--	--	--	--	--	--	--	--	2.33	9.66	113.00	9.75
Sep-80	1980	3	23.33	10.12	81.04	--	--	--	--	--	--	--	--	--	--	--	--	1.33	6.33	156.00	8.86
Sep-81	1981	3	7.33	10.88	82.77	--	--	--	--	--	--	--	--	--	--	--	--	6.67	8.78	165.33	8.70
Oct-82	1982	3	8.33	11.78	99.67	--	--	--	2.33	15.79	109.67	--	--	--	--	--	--	3.33	9.66	109.67	9.69
Oct-83	1983	3	3.33	11.79	94.66	--	--	--	3.67	25.10	117.07	--	--	--	--	--	--	1.33	--	98.33	--
Sep-84	1984	3	3.00	11.26	95.43	--	--	--	3.67	26.64	111.21	--	--	--	--	--	--	0.67	11.00	58.33	10.50
Sep-86	1986	6	15.00	11.50	98.90	--	--	--	4.17	16.68	109.86	--	--	--	--	--	--	0.00	--	42.00	--
Sep-87	1987	6	11.33	13.61	92.06	0.33	6.30	--	5.17	22.43	91.71	--	--	--	0.00	--	--	0.00	--	18.00	--
Sep-88	1988	6	9.67	14.74	90.40	8.17	5.93	105.50	3.00	27.55	123.61	--	--	--	0.67	10.58	86.48	4.00	--	14.00	--
Sep-89	1989	6	10.67	13.15	93.45	9.17	7.59	96.04	1.17	30.31	94.56	--	--	--	0.00	--	--	2.50	--	14.33	4.13
Sep-90	1990	6	18.50	11.96	88.66	4.00	8.51	95.13	0.67	20.95	100.49	--	--	--	2.67	13.69	81.72	9.17	8.04	9.67	14.12
Sep-91	1991	6	15.50	12.78	93.26	12.00	7.39	103.98	2.33	16.57	95.37	--	--	--	5.67	13.98	90.24	2.83	--	8.17	--
Sep-92	1992	6	13.67	13.74	93.42	6.00	6.37	91.54	3.33	25.64	113.39	--	--	--	2.33	17.84	94.80	1.33	--	7.67	--
Sep-93	1993	6	3.17	16.43	94.48	12.33	7.20	109.06	2.00	27.49	100.01	--	--	--	3.33	16.75	95.36	0.00	--	8.67	--
Sep-94	1994	6	27.67	11.73	99.87	23.83	7.65	101.80	2.83	25.52	114.54	--	--	--	1.67	17.39	103.33	0.00	--	6.00	--
Sep-95	1995	6	20.17	13.42	96.73	20.00	7.71	102.97	3.50	21.66	96.62	--	--	--	2.50	17.96	90.90	0.00	--	12.83	--
Sep-96	1996	6	7.83	12.56	96.59	38.00	7.58	105.79	2.83	24.86	103.02	0.17	10.10	119.26	3.33	16.68	96.53	0.00	--	11.00	3.75
Sep-97	1997	6	6.83	13.00	91.31	39.50	7.22	94.54	4.17	21.70	99.11	0.00	--	--	2.17	17.65	96.90	0.00	--	6.17	--
Sep-98	1998	6	4.50	15.53	86.75	47.17	7.55	93.84	4.83	24.43	94.79	0.33	11.65	114.91	4.33	18.04	96.05	0.00	--	10.17	13.74
Sep-99	1999	5	4.20	12.26	104.04	40.60	8.39	93.18	2.20	24.17	105.00	0.80	8.95	119.90	4.40	15.24	95.74	0.20	17.30	4.60	13.39
Sep-00	2000	6	1.00	15.07	93.40	25.00	7.52	96.67	2.50	25.33	99.20	0.50	7.80	104.56	4.67	16.66	96.31	0.00	--	4.17	0.00
Sep-01	2001	6	14.50	12.09	92.76	30.67	7.39	100.86	1.00	27.73	96.81	0.17	10.40	108.60	4.50	13.93	93.62	0.17	17.10	8.67	14.72
Sep-02	2002	6	3.33	11.98	96.85	21.67	7.98	100.11	1.17	25.76	96.31	0.50	9.43	99.04	7.67	14.90	89.57	0.17	--	5.33	--
Sep-03	2003	5	15.80	11.46	102.26	12.20	7.94	125.10	2.00	13.90	108.18	0.20	10.40	96.53	3.60	14.74	101.16	0.00	--	2.60	--
Sep-04	2004	6	12.83	11.62	93.09	16.17	8.34	99.43	0.67	23.90	103.89	0.33	8.20	103.42	2.50	15.32	68.68	0.17	19.20	5.17	15.99
Sep-05	2005	6	5.50	13.63	97.00	12.33	8.35	102.88	0.50	29.23	104.05	0.00	--	--	3.33	15.29	96.82	0.00	--	6.00	16.57
Sep-06	2006	6	3.00	13.38	143.90	23.00	7.71	101.30	1.50	26.94	97.10	0.00	--	--	3.00	15.08	98.10	0.00	--	3.00	16.89
Sep-07	2007	6	9.00	11.80	95.70	29.33	7.90	107.00	1.67	27.50	101.50	0.17	9.20	107.20	5.17	12.80	103.80	0.00	--	17.00	17.20
Sep-08	2008	6	10.00	12.05	104.30	26.50	8.01	102.48	1.00	28.10	97.53	0.17	14.00	113.20	2.67	19.80	94.20	0.00	--	1.83	16.89
Sep-09	2009	6	4.00	11.80	100.90	20.00	8.20	100.40	2.33	26.40	95.16	0.17	15.70	124.59	3.67	18.26	104.72	0.00	--	0.83	16.90

Table 9. – Summary of young of year yellow perch (YP), white sucker (W SU), spottail shiner (SP SH), Iowa Darter (IOWA), fathead minnow (FH MN), largemouth bass (LMB), northern pike (NP), walleye (WE), and other fishes captured by beach seining in Beaver Creek Reservoir, 1980 to 2009.

Date	Sites	YP (yoy)	YP (adult)	W SU	SP SH	IOWA	FH MN	LMB	SMB (yoy)	SMB (adult)	NP (yoy)	NP (adult)	WE (yoy)	WE (adult)	Other Sp. <sup>1</sup>
Jul-80	5	--		650	--	0	42	--	--		--		--		46
Jul-81	5	--		1,671	--	0	75	12	--		--		--		38
Jul-82	5	--		7	--	0	0	54	--		0		--		0
Jun-83	5	--		46	--	0	0	5	--		5		--		0
Aug-84	7	--		189	--	10	0	4	--		0		--		0
Sep-85	5	--		2,648	--	11	0	33	--		3		--		7
May-86	4	--		1,749	0	2	0	0	--		1		--		24
Jun-86	6	--		3,132	0	2	0	0	--		1		--		1
Aug-86	6	--		134	0	8	0	2	--		9		--		0
Sep-86	6	--		1,111	0	34	29	184	--		6		--		11
Jul-87	6	1,968		2,276	1	24	3	0	--		20		11		3
Aug-87	6	2,315		973	0	59	1	16	--		19		19		5
Jun-88	6	20		17	0	6	0	0	--		1		3		0
Aug-88	6	4,973		62	1	4	0	0	--		1		2		0
Aug-89	6	50		48	603	0	0	0	--		2		4		5
Aug-90	6	42		1	93	2	0	0	--		2		0		1
Aug-91	6	8,642		348	835	0	0	0	--		17		0		4
Aug-92	6	1,888		492	156	4	0	0	--		4		0		0
Aug-93	6	42		0	355	11	0	0	--		27		0		0
Aug-94	6	707		49	181	0	0	0	--		11		0		0
Aug-95	6	7,210		6	1,438	0	0	0	--		13		0		0
Aug-96	6	51		261	248	7	0	0	0		5		7		0
Aug-97	6	17		31	193	6	0	0	8		13		2		0
Aug-98	6	872		0	141	0	0	0	41		6		1		0
Aug-99	6	592		4	87	0	0	0	16		7		2		0
Aug-00	6	402		1	190	0	1	0	12		3		23		0
Aug-01	6	357		10	216	0	0	0	8		0		3		0
Aug-02	6	333		0	592	0	0	0	7		0		93		0
Aug-03	6	557		19	2,355	2	0	0	9		15		1		0
Aug-04	6	1,545		0	0	1	0	0	5		2		2		0
Jul-05	6	185		3	1	0	0	0	0		36		12		0
Aug-06	6	1,154		8	608	0	0	0	12		32		11		0
Jul-07	6	253		0	0	0	0	0	13		4		9		0
Jul-08	6	113		0	0	0	0	0	2		0		0		0
Aug-09	6	1,177	135	0	3	0	0	0	1	1	15	1	63	1	0

<sup>1</sup> Consists of emerald shiners, northern redbelly dace, lake chub, western silvery/plains minnow, brassy minnow, and longnose dace

## **Fresno Reservoir**

Fresno Reservoir, located 12 miles northwest of Havre, was built in 1939 for irrigation purposes along the Milk River. Fresno is a highly fluctuating mainstem reservoir of 5,757 surface acres with a mean depth of 27 feet, and a maximum depth of 48 feet. Fresno was initially developed as a rainbow trout fishery in the 1940's and 50's, however an illegal introduction of northern pike in the 1940's resulted in a severe decline in the rainbow trout fishery. As a result, Fresno was developed as a warm-water fishery and walleye, yellow perch, crappie, largemouth bass, smallmouth bass, Lake Superior whitefish, emerald shiner, and spottail shiners were introduced. Over the years, kokanee salmon, brown trout, and rainbow trout have been introduced to supplement the fishery when walleye and northern pike populations were low.

The fishery in Fresno has fluctuated throughout the years due to high fluctuations in water levels. On average, water levels in Fresno fluctuate 21.1 feet per year. The timing of this fluctuation greatly impacts the reproduction and survival of forage and sport fish. The fishery in Fresno was severely decreased in 2001 and 2002 when severe drought reduced the reservoir to 8% and 4% of storage capacity, respectively. Forage fish populations were drastically reduced and abundance and condition of key sport fish was at an all time low. As a result, 170,000 pre-spawn adult yellow perch were introduced to increase population levels so that when water levels increased, forage fish populations could rebound. In addition, 100,000 walleye fingerlings have been stocked annually since 2003.

In 2004, water levels increased and flooded shoreline vegetation, allowing the successful spawning and recruitment of forage fishes. From 2005 to 2009, water levels remained high during spring spawning and early summer rearing allowing sport and forage fish populations to rebound. The continued recovery of the fishery is dependent on maintaining water levels that will allow the successful spawning and recruitment of forage and sport fishes.

### **Population Status of Young-of-Year Fishes**

The abundance and reproductive success of sport and forage fishes were monitored at 12 fixed sites, which were established in 1968. Beach seining was conducted in late summer using a 100- x 9-foot x ¼ inch square mesh beach seine. Fish were sorted by species and counted.

Historically, the abundance of YOY fishes has been dictated by the annual fluctuations in water levels within Fresno Reservoir. The timing of this fluctuation greatly impacts the reproduction and survival of forage and sport fish. The extreme draw down of Fresno in 2001 and 2002 greatly reduced the population levels of most fishes in Fresno except for sauger, which took advantage of the increased riverine habitat (Table 10).

Since 2002, YOY forage and sport fish populations have been increasing and hopefully will continue to recover. While the reservoir levels were reduced, vegetation regenerated along the shoreline and provided spawning habitat for forage and sport fish when water levels increased in 2003. As a result, forage fish such as yellow perch, emerald shiners, and spottail shiners as well as black crappie and northern pike successfully spawned (Table 10).

The yellow perch population is slowly recovering and will continue to increase if water levels are maintained during spawning and rearing stages. Northern pike YOY levels have fluctuated throughout the years, however the population appears to be on the upward trend. In 2008 and 2009, Fresno filled to capacity and flooded a substantial amount of shoreline vegetation creating prime spawning and rearing habitat. Summer seining efforts revealed walleye, northern pike, yellow perch, spottail shiner, and black crappie, all benefited from this rise in water levels with excellent reproduction and survival (Table 10).

Table 10. – A summary of forage fish and young-of-year forage and sport fish collected using a 100- x 9-foot x ¼ inch square mesh beach seine in Fresno Reservoir, 1968-2008.

Year	Seine Hauls	Sanders	Walleye	Sauger	Northern Pike	YP (yoy)	YP (adult)	Emerald Shiner	Crappie Sp.	Spottail Shiner	Sucker sp. <sup>1</sup>	Minnow sp. <sup>2</sup>	Other <sup>3</sup>
1968	12	16	--	--	6	2,909	--	147	552	0	0	161	0
1969	12	4	--	--	6	1,140	--	385	67	0	2	380	0
1970	12	27	--	--	45	10,151	--	521	883	0	1	122	0
1972	12	102	--	--	22	1,005	--	205	379	0	0	72	0
1974	12	13	--	--	59	1,583	--	29	1,355	0	0	25	0
1975	11	10	--	--	32	4,154	--	155	59	0	0	0	0
1978	12	22	--	--	42	10,684	--	12	3	0	0	0	0
1979	12	29	--	--	45	8,516	--	340	127	0	1	0	1
1982	12	102	--	--	70	8,993	--	121	166	0	0	0	3
1983	12	23	--	--	0	2,254	--	448	9	0	1	7	0
1984	12	247	--	--	0	197	--	375	0	2	40	55	0
1985	12	64	--	--	0	379	--	684	3	2	0	9	0
1986	12	0	--	--	23	6,077	--	142	2	20	1	5	1
1987 <sup>+</sup>	12	80	--	--	113	6,233	--	1,979	7	3	0	3	0
1988	12	53	--	--	4	3,122	--	182	0	20	0	1	0
1989 <sup>+</sup>	12	56	--	--	32	24,706	--	22	0	16	2	0	0
1990	12	8	--	--	57	2,033	--	7	465	44	1	2	0
1991 <sup>+</sup>	12	8	--	--	36	3,425	--	0	42	53	0	0	0
1992 <sup>+</sup>	12	45	--	--	2	6,550	--	28	0	48	0	1	0
1993 <sup>+</sup>	12	24	--	--	9	5,595	--	12	2	162	0	0	0
1994 <sup>+</sup>	12	19	--	--	19	2,960	--	3	287	1,421	1	0	0
1995	12	5	--	--	2	1,080	--	0	2	129	0	1	0
1996 <sup>+</sup>	12	52	--	--	21	3,576	--	0	1	1,484	42	0	0
1997 <sup>+</sup>	12	46	--	--	15	3,006	--	2	1	887	2	0	0
1998 <sup>+</sup>	12	44	--	--	1	1,413	--	9	0	1,041	1	3	0
1999	12	50	--	--	7	4,271	--	176	12	182	13	0	0
2000	6	29	--	--	0	1,396	--	2	2	30	2	0	1
2001	6	86*	--	--	0	39	--	3	0	3	3	1	0
2002	12	28*	--	--	2	86	--	128	400	154	4	29	0
2003 <sup>+</sup>	12	4	--	--	46	1,871	--	5,539	90	207	0	0	1
2004 <sup>+</sup>	12	--	12	2	10	2,898	--	69	48	56	0	2	1
2005 <sup>+</sup>	12	--	26	2	19	934	--	39	15	39	0	0	0
2006 <sup>+</sup>	12	--	27	0	57	2,283	--	80	5	923	0	0	0
2007 <sup>+</sup>	12	--	7	0	13	769	--	68	54	1,106	2	0	0
2008 <sup>+</sup>	12	--	65	0	1	2,329	--	5	721	287	11	0	0
2009 <sup>+</sup>	12	--	24	0	24	1,427	224	13	25	716	1	0	0

<sup>1</sup> Consists of white and longnose sucker

<sup>2</sup> Consists of silvery minnows, lake chubs, flathead chubs, and fathead minnows

<sup>3</sup> Consists of burbot, smallmouth bass, pumpkinseed sunfish, and brook sticklebacks

<sup>+</sup> Years in which walleye fry or fingerling were stocked

\* Primarily Sauger


## Population Status of Adult Fishes

Adult fish populations were monitored from 1965 to 1974, using systematic gill netting at predetermined stations. Sampling at 12 predetermined stations was resumed in 1987 to determine changes in sport fish abundance and species composition. Samples were collected over two days utilizing six sinking multi-filament experimental gill nets each day (12 net-days). The sinking multi-filament experimental gill nets measuring 125 feet in length and 6 feet deep consisting of 25-foot panels of  $\frac{3}{4}$ ", 1", 1  $\frac{1}{4}$ ", 1  $\frac{1}{2}$ ", and 2" mesh. Fish were measured for total length (TL) to the inch and weighed to the nearest 0.01 pound. Prior to 2005, scales were collected for aging from all walleye and sauger. From 2005 to 2009, otoliths were collected from walleye for aging and oxytetracycline (OTC) analysis.

### Lake Superior Whitefish

Lake Superior whitefish (whitefish) in Fresno Reservoir historically have comprised a significant portion of the gill net catch (Figure 4), but are rarely utilized by anglers. Whitefish exhibit high growth rates in the reservoir, and thereby escape predation from all but the largest walleye and northern pike. Whitefish appear to reproduce successfully in years of stable over-winter storage.

Figure 4. - Relative abundance and size structure of lake whitefish collected with sinking experimental gill nets in Fresno Reservoir, 1987-2009.




### Yellow Perch

The yellow perch fishery in Fresno has been negatively impacted by drought over the last eight years. As a result of extreme draw downs in 2001 and 2002, yellow perch were not able to successfully spawn (Table 10) and population levels were drastically reduced (Figure 5). To remedy this situation, 170,000 pre-spawn adult yellow perch were introduced from 2001 through 2004 to increase population levels so when water levels increased these forage fish populations might rebound. In 2003 and 2004, water levels increased, flooding shoreline vegetation, and allowing the successful spawning and recruitment of forage fish, as a result stocking of pre-spawn perch was discontinued in 2005.

Since stocking of adult yellow perch was discontinued in Fresno Reservoir, the abundance of adult yellow perch has decreased, however the abundance levels remain higher than most pre-drought

levels (Figure 5). In addition, the proportion of stock size fish in the population indicates that YOY fish have successfully recruited into the population. However, this number continues to decline since stocking of adult yellow perch ceased in 2005. The continued recovery of the fishery is dependent of maintaining water levels that will allow the successful spawning and recruitment of forage and sport fishes.


Figure 5. - Relative abundance, size structure, and relative weight of yellow perch collected with sinking experimental gill nets in Fresno Reservoir, 1987-2009.



### Northern pike

Since the illegal introduction of northern pike in Fresno Reservoir in the 1940s, their population has fluctuated over the years (Figure 6). Extreme drought from 2000 to 2002 reduced the abundance of northern pike however, the population rebounded in 2003 with increased water levels and flooding of shoreline vegetation. Since 2003, reproduction has steadily increased with high numbers of YOY being collected during annual beach seining surveys. However, northern pike recruitment was significantly reduced in 2008 (Table 10).

Figure 6. - Relative abundance, size structure, and relative weight of northern pike collected with sinking experimental gill nets in Fresno Reservoir, 1987-2009.




## Walleye

From 2003 to 2009 approximately 100,000 fingerlings were stocked annually in Fresno Reservoir. In 2006, 200,000 fingerling walleye were stocked due to unallocated productions at the Ft. Peck fish hatchery.

In 2007, walleye were at their highest abundance levels on record (CPUE=26.25 fish/net; Figure 7) indicates excellent survival and recruitment of stocked YOY walleye, especially the 2005 and 2006-year class. In 2008 and 2009, walleye abundance decreased slightly (CPUE=21.5 and 22.05 fish/net; Figure 7), but re-main well above abundances observed in the late 1990s and early 2000s.

Figure 7. - Relative abundance, size structure, and relative weight of walleye in Fresno Reservoir for the years 1987-2009.




## Sauger

Sauger populations have been relatively low in Fresno Reservoir since the construction of the Fresno Dam. Sauger have been captured periodically primarily in the upper reservoir where the Milk River enters Fresno Reservoir. Sauger populations increased in 2001 and 2002 when drought and extreme drawdowns resulted in an increase in riverine habitat (Table 10), however no sauger were collected in 2009.

## Black Crappie

Black crappies were most likely introduced into Fresno in the 1950s however the first record of stocking occurred in 1991. Since 1968, YOY crappie numbers have fluctuated. In 2008, YOY black crappie abundance was the highest on record since 1974 (Table 10). However the population of adult black crappie has remained relatively low (CPUE range = 0.17 to 1.33 fish/net; 1999 to 2009) and may be misconstrued due to sampling gear bias.

Figure 8. - Relative abundance and size structure of black crappie collected with sinking experimental gill nets in Fresno Reservoir, 1987-2009.




### **Blaine County Ponds**

Select waters throughout Blaine County were sampled to determine fish abundance using sinking multi-filament experimental gill nets measuring 125 feet in length and 6 feet deep consisting of 25-foot panels of ¾", 1", 1 ¼", 1 ½", and 2" mesh unless otherwise specified. Voluntary creel boxes were maintained at many of the ponds to determine fishing pressure, catch rates, and satisfaction.

#### **Butch Reservoir**

Butch reservoir is located in south Blaine County and contains a fishery for rainbow trout, largemouth bass, and bluegill. Butch also has a windmill aerator system, which assists in over winter survival of fish. In the summer of 2005, a voluntary creel box was erected at Butch reservoir to determine fishing pressure, angler success, and angler satisfaction. In 2005, anglers reported summer catch rates of rainbow trout as 4.28 fish/hour, catches of bluegills as 1.36 fish/hour, and largemouth bass as 2.47 fish/hour (n=4). Anglers reported fall catch rates of rainbow trout as 1.65 fish/hour, catches of bluegills as 0.04 fish/hour, and largemouth bass as 2.13 fish/hour (n=5). The box was destroyed by cows during the winter of 2005/2006 and not replaced.

Figure 9. - Relative abundance of rainbow trout, largemouth bass, and bluegill collected with sinking experimental gill nets in Butch Reservoir, 1981-2009.





### Cow Creek Reservoir

Cow Creek Reservoir is a privately owned 65-acre pond located in the Bears Paw Mountains. Cow Creek Reservoir has been managed as a warm water fishery since 1994. Since 1994, walleye, channel catfish, black crappie, and tiger muskie have been stocked and yellow perch were illegally introduced in 2001. In 2007, 5,000 walleye, 3,000 channel catfish, and 820 7 inch tiger musky were stocked into Cow Creek Reservoir. Starting in 2008, 5,000 walleye and 1,000 channel catfish will be stocked in alternate years. Tiger Musky will be stocked on an as need basis. The primary food sources for these sport fish are white suckers, fathead minnows, golden shiners, and northern red belly dace. In 2005, this reservoir received 103 angler days and ranked 56<sup>th</sup> in the region for angler pressure (McFarland 2006).

Table 11. – Catch rate (CPUE (fish/net)) and average length of yellow perch, channel catfish, white sucker, walleye, and tiger muskie using gill nets in Cow Creek Reservoir (1994-2008).

Date	Nets	Yellow Perch			Channel Catfish			White Sucker		Walleye			Tiger Muskie	
		CPUE	Ave. Length		CPUE	Ave. Length	Rel. Weight	CPUE	Ave. Length	CPUE	Ave. Length	Rel. Weight	CPUE	Ave. Length
Aug-94	2.0		--		0.0	--		2.0	--	23.5	7.2		0.0	--
Sep-95	1.0	0.0	--		0.0	--		2.0	--	15.0	10.0	82.5	0.0	--
Sep-96	2.0	0.0	--		5.0	9.1	116.1	1.0	--	48.0	11.1	82.3	0.0	--
Sep-97	2.0	0.0	--		9.5	10.5	118.1	1.0	--	30.5	11.9	86.9	0.0	--
Sep-98	3.0	0.0	--		6.3	13.9	107.7	7.0	14.6	11.3	13.2	87.1	0.0	--
Sep-01	2.0	0.5	5.6		4.5	17.0	103.7	0.5	--	12.5	13.3	94.7	0.5	15.7
May-03	2.0	0.0	--		11.0	19.5	115.7	8.0	15.9	1.0	13.0	97.0	1.5	19.4
Jul-05	2.0	1.0	9.8		9.0	21.3	104.3	6.0	17.6	8.0	14.7	85.5	0.0	--
Jul-06	2.0	1.5	9.6		9.5	21.5	108.4	7.0	17.6	12.0	13.0	87.1	0.0	--
Jul-07	2.0	0.5	10.3		7.0	23.5	118.8	0.0	--	7.5	11.8	92.2	1.5	21.5
Jul-08	2.0	0.0	0.0		6.0	14.4	120.4	2.5	18.1	4.5	9.3	90.5	0.0	0.0
Jun-09	2.0	0.5	10.4		8.0	22.7	111.3	1.5	15.2	13.0	10.0	96.1	0.5	19.7

### Gazob Reservoir

Gazob Reservoir is a 10-acre pond located on BLM land in south Blaine County. Largemouth bass, bluegill, and fathead minnows were introduced in 1986 and 1987. Golden shiners are also present within the reservoir, however their introduction was not recorded.

Seining was conducted in 1988 and 1990, and electrofishing was conducted in 1994. In 2005, one experimental gill net was set perpendicular to the dam. In 2005, the average size of largemouth bass captured was 10.4 inches and bluegills were 4.4 inches. In 2009, one gill net and one trap net were set overnight. The gill net collected 101 golden shiners and one largemouth bass, the trap net contained 10 bluegill and 34 golden shiners.

### Reser Reservoir

Reser reservoir is located in northwestern Blaine County. This reservoir has been managed as a fishery since 1981 and over the years has been stocked with fathead minnows, lake chub, northern redbelly dace, western silvery/plains minnows, golden shiners, largemouth bass, black crappie, bluegill, and rainbow trout. This reservoir had frequent winterkills occur in the early 1990s and as a result two windmill aeration systems were installed. Since the installation of the aeration systems only one fish kill has occurred and this was suspected to have occurred as a result of chemical runoff from surrounding fields.

Figure 10. - Relative abundance of rainbow trout, largemouth bass, yellow perch, golden shiner, black crappie, and bluegill collected with sinking experimental gill nets in Reser Reservoir, 1987-2009.





Figure 11. - Relative abundance of rainbow trout, largemouth bass, yellow perch, golden shiner, black crappie, and bluegill collected with trap nets in Reser Reservoir, 1983-2009.



### South Casady (Casady Pond/ BR019)

South Casady is a 3-acre pond located on BLM land in North Blaine County. From 1966 to 1974 Casady was managed as a rainbow trout fishery. In the 1980s, largemouth bass, bluegills, black bullheads, yellow perch, northern pike, and golden shiners were introduced. Water levels have not been favorable in Casady and as a result a black bullhead fishery is all that remains. In 2009, a gill netting survey resulted in catch rates of 27 black bullheads/net (TL=5.7 in.). One trap net was set overnight as well and captured 10 black bullhead, 25 golden shiner, and 884 fathead minnows.

### Sundance Reservoir

Sundance reservoir is small reservoir located on BLM land in south Blaine County. This reservoir was created in the mid-1980s. Stocking records do not exist and in 2005 four nine-inch largemouth bass were collected. Sampling effort (one gill and one trap net) in 2009 suggests no largemouth bass remain in Sundance, if they do it is a small population.

### Phillips County Fishing Waters

Select waters throughout Phillips County were sampled to determine fish abundance using sinking multi-filament experimental gill nets measuring 125 feet in length and 6 feet deep consisting of 25-foot panels of  $\frac{3}{4}$ ", 1", 1  $\frac{1}{4}$ ", 1  $\frac{1}{2}$ ", and 2" mesh. Voluntary creel boxes were maintained at many of the ponds to determine fishing pressure, catch rates, and satisfaction.

### Bar Island Reservoir

Bar Island Reservoir is a small reservoir located on BLM land in south Phillips County. This reservoir was sampled in 2003 and again in 2009 with gill and trap nets. Gill nets captured only white suckers in both years (CPUE= 52 fish/net) and the trap net set in 2009 contained 266 white suckers and 1,610 fathead minnows.


### Bell Ridge Reservoir

Bell Ridge is a BLM pond, which was last stocked in 1999 with channel catfish. In 2009, the deepest point within the reservoir was 4 feet and it was approximately 10 feet below the spillway. Water levels used to be supplemented with well water, however that practice was terminated. Due to low water levels, Bell Ridge is no longer suitable as a fishery.

### Cole Ponds

The Cole Ponds are a state fishing access site and consists of three ponds that are approximately 9 acres each. These ponds are old gravel pits and are very deep clear ponds. These ponds containing self-sustaining populations of largemouth bass, yellow perch, northern pike, pumpkinseed sunfish, and black crappie.

Figure 12. - Catch rates of fishes in the Cole Ponds from summer gill netting surveys (periodic sampling 1988 to 2009).



### Compton Reservoir


Compton is a 11-acre pond located on BLM and private land in northern Phillips County. This reservoir was established as a rainbow trout fishery in 1963 and was stocked with rainbow trout until 1990. In 1988, yellow perch were introduced and at some point northern pike were also introduced as well. Gill netting surveys in 2005 resulted in the catch of three tiger salamanders but no fish. In 2009, one gill net and one trap net were set overnight. No fish were captured in the gill net. However, the trap net contained 3,295 fathead minnows. Water levels were very good at the time of sampling and a reintroduction of yellow perch will take place in 2010.

### Ester Lake

Ester Lake is a 139-acre pond located on state land and has been managed as a fishery since the 1950s. In the 1960's Ester was a productive fishery with high numbers of yellow perch, black crappie, and walleye. Since the 1980s the fishery has been in decline and water levels have been marginal in recent years. Due to reduced fishing opportunities, Ester only received 33 angler days in 2005 and ranked 63<sup>rd</sup> in the region (McFarland 2006).

In 2009, Ester received 8,000 pre-spawn yellow perch to boost the forage base that has been non-existent since the early 1980s (Figure 13).

Figure 13. - Relative abundance of yellow perch, northern pike, carp, white sucker, black crappie, and walleye in Ester Lake (periodic sampling 1980 to 2009).



### Frenchman Reservoir

Frenchman reservoir is a main channel reservoir on Frenchman Creek northwest of Saco. This reservoir is approximately 806 surface acres and is surrounded by private land with public access occurring primarily around the dam. This reservoir is fairly shallow and partial winterkills are not uncommon. This reservoir is not stocked and all the fish present within the reservoir come from

Frenchman Creek. This reservoir receives a minimal amount of fishing pressure even though the fish have high growth rates and condition factors.

In 2009, two gill nets and two trap nets were set overnight to assess the fish population. The gill nets contained walleye (CPUE= 1.5 fish/net), northern pike (CPUE= 2), common carp (CPUE= 26.5), and white sucker (CPUE=17). Trap nets contained walleye (CPUE= 0.5 fish/net), northern pike (CPUE= 1), common carp (CPUE= 100.5), white sucker (CPUE=5.5), black bullhead (CPUE= 4), fathead minnow (CPUE= 1,248.5), and brassy minnow (CPUE= 77.5).

### **Gullwing Reservoir**

Gull wing is a 17-acre BLM pond located in south Phillips County. This reservoir has a windmill aeration system and good water levels. Black crappie (1,200 8-in.) were introduced into this reservoir in 2003, however this stocking was apparently not successful because no fish were collected in 2005. Numerous fathead minnow were observed in the shallows. The re-introduction of largemouth bass was re-attempted in 2006. In 2009, one gill net was set overnight and captured 34 largemouth bass (TL= 7.5 in.).

### **Karsten Coulee**

Karsten Coulee reservoir is located on BLM land in south Phillips County. Like many ponds on BLM land in south Phillips County, livestock grazing has reduced water quality and eliminated shoreline vegetation. In 2000 an aerator was installed on Karsten Coulee and largemouth bass were stocked in 1999 and 2004. Prior to restocking in the spring of 2004, a winterkill was reported. Low water levels and poor water quality were most likely responsible. In 2009, one gill net was set overnight. Largemouth bass ranging in size from 6.6 to 9.6 inches in length were collected (CPUE = 11 fish/net). Hook and line sampling was also conducted yielding catch rates of 15 fish/hour with several bass over 14 inches in total length. This reservoir would benefit from fencing of at least a portion of the shoreline to allow the pond to recover from a chronic overgrazing problem.


### **Lark Reservoir**

Lark Reservoir is a 6-acre pond located on BLM land in south Phillips County. Lark was managed as a rainbow trout fishery from 1970 to 1994. Since the mid-1990s this reservoir has been managed as a largemouth bass fishery. In 2009, a gill netting survey resulted in catch rates of 2 largemouth bass per net (TL=10.5 in.). In 2010, introductions of a forage base (fathead minnows, black crappie, or bluegill) needs to be addressed.

### **McChesney Reservoir**

McChesney Reservoir is a privately owned reservoir in south Phillips County. This reservoir has been managed as a northern pike, yellow perch, and black crappie fishery since the 1960s. There were also stockings of bluegill, however stocking records are not complete. In 2005, this reservoir received 187 angler days and ranked 42<sup>nd</sup> in Region 6 for fishing pressure (McFarland 2006).

Figure 14. Relative abundance of yellow perch, northern pike, common carp, white sucker, and bluegill in McChesney Reservoir (periodic sampling 1984 to 2009).



### Nelson Reservoir

Nelson Reservoir, located 19 miles east of Malta, is an off-channel storage reservoir constructed in 1915 for irrigation purposes. At full storage capacity, Nelson covers approximately 4,320 surface acres, has a mean depth of 14.2 feet, and a maximum depth of 50 feet. Nelson is a relatively stable reservoir, which is not affected by drought as drastically as some reservoirs in the region with an average annual fluctuation of 8.36 feet.

Nelson was established as a fishery in the 1930s & 40s with the introduction of largemouth bass, crappie, bullheads, and rainbow trout. Commercial fishing for carp, buffalo, and goldeye was conducted in the 1920s, 30s, and in the mid 60s. Nelson has approximately 26 fish species and is managed primarily as a walleye fishery. Walleye reproduce naturally in Nelson; however walleye fry are occasionally stocked to augment natural reproduction. Spawning shoals were constructed in 1993 at three locations within the reservoir to improve the spawning habitat for walleye.

### Population Status of Adult and Young-of-Year Fishes

Since 1993, adult fish populations were monitored at 10 fixed experimental gill netting stations. Gill netting was conducted over a two-day period utilizing five sinking experimental gill nets each day (10 net-days). The sinking multi-filament experimental gill nets measured 125 feet in length and 6 feet deep consisting of 25-foot panels of  $\frac{3}{4}$ ", 1", 1  $\frac{1}{4}$ ", 1  $\frac{1}{2}$ ", and 2" mesh. Fish were measured for total length (TL: inches) and weighed to the nearest 0.01 pound (lb). Otoliths were collected from walleye for aging and oxytetracycline (OTC) analysis.

Prior to 1991, adult fish populations were monitored, however sampling was neither uniform, nor consistent enough to develop useful trend data on game fish population size, or composition. As a result this data was excluded from all analysis.

The abundance and reproductive success of sport and forage fishes were monitored at nine predetermined stations. Beach seining was conducted in late July using a 100- x 9-foot x  $\frac{1}{4}$  inch square mesh beach seine. Fish were sorted by species and counted.

Table 12. - A summary of forage fish and young-of-year forage and sport fish collected conducted at nine fixed sites using a 100- x 9-foot x ¼ inch square mesh beach seine in Nelson Reservoir, 1982-2009.

	Shorline Seined (ft)	Walleye	Yellow Perch	Northern Pike	Spottail Shiner	White Sucker	Black Crappie	Goldeye	Buffalo sp <sup>1</sup>	Smallmouth Bass	Longnose Sucker
1982	660	0	4,553	3	0	202	245	0	0	0	0
1983	1,420	4	138	18	0	543	238	0	0	0	0
1984	1,530	0	133	0	0	0	0	0	0	0	0
1985	510	3	2,272	16	1	16	67	1	0	0	0
1986*	700	0	3	7	0	10	232	0	0	0	0
1987*	495	5	1,987	0	4	45	10	7	0	0	0
1988*	520	0	783	0	1	0	35	0	0	0	0
1989*	910	10	736	4	43	1,503	135	0	0	0	0
1990	1,320	7	2,631	1	56	181	21	0	0	0	0
1991*	660	8	77	1	54	33	26	0	0	0	0
1992	635	21	140	6	387	175	18	0	4	0	0
1993*	520	3	8,287	1	520	2,688	62	0	0	0	0
1994*	830	6	1,802	10	621	697	49	0	0	0	0
1995*	760	36	232	0	3,780	180	163	0	0	0	0
1996*	870	25	4,521	13	21	101	0	0	0	0	0
1997*	890	53	2,205	0	159	534	1	0	0	0	0
1998*	340	0	126	0	33	235	4	0	0	0	0
1999	750	11	1,489	2	222	497	1	0	0	0	0
2000*	440	4	449	2	189	258	5	6	0	0	0
2001	430	2	72	1	27	800	88	0	0	0	0
2002*	415	2	19	4	8	38	482	21	62	0	0
2003	530	3	361	33	49	235	6,597	0	0	3	0
2004*	443	10	1,781	0	19	195	5	1	0	10	0
2005*	754	5	423	2	34	155	278	23	5	1	0
2006*	831	3	773	8	66	319	89	0	3	1	0
2007*	489	6	586	2	75	596	5	0	12	9	0
2008*	500	10	62	0	8	272	1,237	11	94	11	0
2009*	750	4	4,522	4	3	478	20	8	2	61	14

\*Years in which walleye fry or fingerlings were stocked


<sup>1</sup> Consists of bigmouth buffalo and smallmouth buffalo

### ***Yellow Perch***

The yellow perch fishery in Nelson Reservoir has been cyclic over the last 14 years due to drought and water fluctuations. In the early 1990s and in 2000 and 2002 the relative abundance of yellow perch was significantly reduced. However, since 2003 spring and summer rains and water retention improvements by the Bureau of Reclamation (BOR) have allowed water levels to flood shoreline vegetation and remain stable during crucial spawning and rearing periods.

In 2009, yellow perch numbers increased to their highest levels since 1991, at 19.5 fish/net. Currently, the population of adult yellow perch consists mostly of stock (5-7.9 in.) and quality (8.0-9.9 in.) sized fish.

Figure 15. - Relative abundance, size structure, and relative weight of yellow perch collected with sinking experimental gill nets in Nelson Reservoir, 1991-2009.



## Walleye


Walleye fingerlings have been periodically stocked into Nelson Reservoir to augment natural reproduction. From 2002 to 2009 (with the exception to 2006), all walleye fingerlings stocked into Nelson Reservoir have been marked with 750 ppm OTC to allow the calculation of survival on stocked fish and to distinguish stocked fish from naturally reproduced fish. In 2006, only half of the walleye stocked were marked with OTC due to problems with reaction of the walleye to the chemicals.

Even with the addition of these fish, catch of YOY walleye during seining surveys remained low when compared to pre-drought levels (Table 12). However, the high proportion of stock (10-14.9 in.) and quality size (15.0-19.9 in.) walleye in the population indicates good survival of stocked YOY walleye from 2003 through 2009 (Figure 16).

The relative abundance of adult walleye has remained relatively stable over the last five years with a minor decrease in catch rates in 2005 and 2009 (Figure 16). The condition of walleye increased after low water levels in 2002 (Figure 16).




Figure 16. - Relative abundance, size structure, and relative weight of walleye collected with sinking experimental gill nets in Nelson Reservoir, 1991-2009.



### Northern pike

Historically, the abundance of adult northern pike has remained relatively low and stable consisting of a high proportion of quality, preferred, and memorable size fish (Figure 17). Northern pike populations were significantly reduced in 2002, however the population was quickly replenished with the recruitment of YOY fishes in 2003 (Figure 17). In 2009, the northern pike population increased to 3.4 fish/net and has good balance of size and year classes (Figure 17).


Figure 17. - Relative abundance, size structure, and relative weight of northern pike collected with sinking experimental mesh gill nets in Nelson Reservoir, 1991-2009.



## Lake whitefish

Lake whitefish populations have fluctuated since 1991 due to fluctuations in water levels and temperature, which have reduced recruitment of YOY fish to the population (Figure 18). In 2007, there was a massive summer kill of lake whitefish and fall gill netting surveys indicated a drastic decrease in the abundance of lake whitefish (CPUE = 0.8 fish/net; Figure 13). Gill netting surveys conducted in 2009 reveals the population is recovering and reproducing (CPUE = 2 fish/net; Figure 18).


Figure 18. - Relative abundance, size structure, and relative weight of lake whitefish collected with sinking experimental mesh gill nets in Nelson Reservoir, 1991-2009.



## Black Crappie

Historically there has been a low abundance of black crappie in Nelson Reservoir. In 2003, black crappie reproduction was the highest on record (Table 12) and recruitment of YOY crappie into the population resulted in a high abundance of adult black crappie (Figure 19). High reproductive success over the last seven years indicates the conditions within Nelson Reservoir have been favorable for black crappie, and if conditions remain stable the population will continue to remain at good levels.

Figure 19. - Relative abundance, size structure, and relative weight of black crappie collected with sinking experimental gill nets in Nelson Reservoir, 1991-2009.




## Other fishes

A variety of other fishes are found within Nelson Reservoir, however they are rarely utilized as a sport fishery due to low abundances or their non-game status. Channel catfish, stonecats, bigmouth buffalo, smallmouth buffalo, and smallmouth bass are all present at low levels within Nelson Reservoir. Spottail shiners are also present and provide an important forage base, however in recent years their populations have been reduced and adult spottail shiners have not been present in high numbers within the annual seining surveys (Table 12).

### Paleface Reservoir

Paleface is a 4.7-acre BLM pond that contains black crappie, golden shiners, and white suckers. A windmill aeration system was installed in 2000 and a voluntary creel box was erected in 2005. Paleface, like many of other reservoirs, is negatively impacted by over grazing. Cattle have destroyed most of the shoreline vegetation and are having a negative impact on the water quality of the reservoir. Gill netting surveys in 2001 yielded eight black crappie with an average length of 8.5 inches. In 2005, gill netting captured black crappie ranging in length from 6.6 to 7.4 inches (CPUE=0.47 fish/hour), golden shiners (CPUE=0.14 fish/hour), and white suckers (0.05 fish/hour).

Figure 20. Relative abundance of black crappie, golden shiner, and white sucker collected with sinking gill nets in Paleface Reservoir (Periodic sampling 2001-2009).



### PR 018

PR 018 is a 6-acre pond located on BLM land in south Phillips County. PR 018 has a windmill aeration system and was historically a warm water fishery consisting primarily of largemouth bass. In 2004, a winterkill occurred and surplus rainbow trout were stocked in the spring of 2004. Gill netting surveys indicate excellent growth and survival of rainbow trout with a catch rate of 74 fish/net. Rainbow trout ranged in size from 7.7 to 12.2 inches TL (TL=9.92 in.) and 0.07 to 0.63 lbs. (wt=0.35 in.). Population estimates in 2009 captured no fish. Re-introduction of largemouth bass occurred in 2009 (n=3,000) and fathead minnows or bluegill may be introduced in 2010.

### PR 020

PR 020 is a 7-acre pond located on BLM land in south Phillips County and is managed as a largemouth bass fishery. PR 020 was spilling over in 2004 and the reservoir was near full in 2009. Largemouth bass were last stocked in 1993 and are successfully recruiting within this reservoir. In 2009, gill net surveys resulted in one largemouth bass (TL=13.5).


### PR 054

PR 054 is a 7.6-acre BLM pond located 3-miles from the Canadian border. This pond is managed as a largemouth bass fishery and was last stocked with largemouth bass in 2003. In 2009, one gill net was set overnight and captured no fish. One trap net was set overnight as well and contained 42 fathead minnows and one largemouth bass (TL=10.2 in.).

### Sagebrush

Sagebrush reservoir is a 5.2-acre BLM pond located in south Phillips County. Sagebrush was historically managed as a rainbow trout fishery until largemouth bass were introduced in 1989, followed by bluegill in 2001 and channel catfish in 2001. Rainbow trout are stocked periodically, with the last stocking occurring in 2002. A windmill aerator system was installed in 2000 or 2001 and this reservoir is currently full. Bluegill and largemouth bass populations are self-sustaining and rainbow trout have exhibited excellent growth and survival. In 2009, one gill net and one trap net were set overnight. The gill net contained one largemouth bass (TL= 13.2 in.) and the trap net had 163 bluegill and one largemouth bass. This reservoir may need a supplemental stocking of rainbow trout in the future to augment several years of no stocking.

Figure 21. Relative abundance of rainbow trout, largemouth bass, and bluegill collected with sinking gill nets in Sagebrush Reservoir (Periodic sampling 1982-2009).



### Taint Reservoir

Taint reservoir is a 5.7-acre reservoir located on BLM land in south Phillips County. This reservoir is a popular bluegill fishery with high catch rates and good lengths of fish. Taint reservoir has had good water levels, and was full in 2009.

Figure 22. Relative abundance of rainbow trout, largemouth bass, and bluegill collected with sinking gill nets in Taint Reservoir (Periodic sampling 1982-2009).





Figure 23. Relative abundance of largemouth bass and bluegill collected with 3x4 trap nets in Taint Reservoir (Periodic sampling 2005-2009).



### Wapiti Reservoir

Wapiti is located on BLM land is managed primarily as a largemouth bass fishery. A windmill aerator system was installed in 2001 and was out of commission during the winter of 2004/2005. A few dead largemouth bass were reported along the shoreline, however this was only a partial winterkill. Water levels were 5 feet below the spillway during the spring of 2009. One gill net was set in 2009 and captured 12 largemouth bass (TL= 7.9 in.). Hook and line sampling was also conducted and yielded largemouth bass catch rates at 26 fish/hour (TL 6-10 in.). A forage species needs to be introduced to maximize largemouth bass growth in this reservoir.

### Whiteface

Whiteface is located on BLM land is managed as a largemouth bass fishery, however stocking records do not indicate when largemouth bass were introduced to this reservoir. Whiteface reservoir has had good water levels and was full in 2009. Gill netting surveys resulted in marginal catches rates (CPUE=2 fish/net) and size distribution (TL=6.4 in.) of largemouth bass.


### Daniels County Fishing Waters

Select waters throughout Daniels County were sampled to determine fish abundance using sinking multi-filament experimental gill nets measuring 125 feet in length and 6 feet deep consisting of 25-foot panels of  $\frac{3}{4}$ ", 1", 1  $\frac{1}{4}$ ", 1  $\frac{1}{2}$ ", and 2" mesh.

### Whitetail Reservoir

Whitetail Reservoir is located on both private and state land and has been maintained as multi-species fishery since 1946. Largemouth bass were introduced in 1946, followed by northern pike in 1955, rainbow trout in 1962, yellow perch in 1975, walleye in 1979, black crappie in 1997, and fathead minnows in 2002. Yellow perch were last stocked in 2004.

Figure 24. Relative abundance of northern pike and yellow perch collected with sinking gill nets in Whitetail Reservoir (Periodic sampling 2006-2009).



### Valley County Fishing Waters

Select waters throughout Valley County were sampled to determine fish abundance using sinking multi-filament experimental gill nets measuring 125 feet in length and 6 feet deep consisting of 25-foot panels of  $\frac{3}{4}$ ", 1", 1  $\frac{1}{4}$ ", 1  $\frac{1}{2}$ ", and 2" mesh.

### Big Reservoir

Big Reservoir is located on BLM land and has been managed as a fishery since 1986. Black crappie and yellow perch were both introduced in 1986 and have been naturally reproducing in this reservoir since then. In 2006, two trap nets were set for approximately 22 hours and captured 58 black crappie (CPUE = 29 fish/net), ranging in length from 5.0 to 9.0 inches (TL = 5.3 in.). In 2009, one

gillnet and one trap were set overnight. The gill net contained no fish and the trap contained one fathead minnow.

#### **Boucher Reservoir**

Boucher Reservoir is a 9 acre reservoir located on private property. In 2009, one gill net and one trap net were set overnight and captured no fish.

#### **Burke Reservoir**

Burke Reservoir is located on private land in south Valley County. In 2009, one gill net and one trap net were set overnight. The gill net contained 18 yellow perch (TL = 9.5 in.) and the trap had 480 yellow perch (TL = 2.5 in.) and 1,600 fathead minnows.

#### **Lower Base Pond**

Lower Base Pond is located on both private and state land and has been maintained as a fishery since 1961. Rainbow trout were introduced in 1961, followed by common carp and bluegills in 1969, largemouth bass in 1971, and yellow perch in 1985. In 2009, one gill net and one trap net were set overnight. The gill net contained 11 rainbow trout (TL = 8.3 in.) and no fish were captured in the trap net.

#### **Paulo Reservoir**

Paulo Reservoir is located on BLM land and has been managed as a largemouth bass and bluegill fishery. Largemouth bass were introduced in 1993 and bluegill in 2004. In June of 2006, two trap nets were each set for approximately 20 hours. A total of 270 fathead minnows (CPUE = 6.75 fish/hr) and two bluegills (TL = 6.0 in.) were collected. In July 2006, four trap nets and two gill nets were each set for approximately 17 hours to recheck species composition. A total of 2,915 fathead minnows, 24 bluegill, two largemouth bass (TL = 3.25 in.), and one common carp, (TL = 10.5 in.) were captured in the trap nets. One largemouth bass (TL = 12.5 in.) and four bluegills (TL = 6.9 in.) were collected in the gill nets. In 2009, one gill net and one trap net were set overnight. The gill net contained five bluegill (TL = 6.1 in.) and two bluegill (TL = 2.7 in.) were captured in the trap net.

#### **Upper Base Pond**

Upper Base Pond is located on both private and state land and has been maintained as a fishery since 1961. Rainbow trout were introduced in 1961, followed by common carp and bluegills in 1969, largemouth bass in 1971, and yellow perch in 1985. In 2009, one gill net and one trap net were set overnight. The gill net contained five northern pike (TL = 24.7 in.) and no fish were captured in the trap net.

#### **Valley Reservoir**

Valley Reservoir is privately owned and has been open to the public as a multi-species fishery since 1972. Rainbow trout were introduced in 1972 and stocking was discontinued in 1986 when yellow perch and largemouth bass were stocked followed by walleye in 1989. In 2009, one gill net and one trap net were set overnight. A total of 71 yellow perch (TL = 4.3) were collected using the trap net. The gill net captured a total of 72 yellow perch ( $\bar{x}$  TL = 7.1 in.). Heavy vegetation was observed around the edges of the reservoir.

### **Wards Reservoir (VR 082)**

Wards Dam is located on BLM land has been managed a multi-species fishery since 1940. Bass, crappie, and bullheads were introduced in 1940, followed by bluegill and largemouth bass in 1944, rainbow trout in 1963, northern pike in 1970, black bullheads in 1974, and fathead minnows in 1989. In 2006 two gill nets were set for approximately 17 hours each, and no fish were captured. In 2009, one gill net and one trap net were set overnight. No fish were captured in the gill net and the trap net contained 1,250 fathead minnows.

## **RECOMMENDATIONS**

### **Paddlefish: Fort Peck Stock**

Annual tagging efforts should continue with over 300 paddlefish being tagged annually. An on-site creel survey should be conducted in 2010 due to recent changes in harvest regulations. Annual collections of paddlefish jaws should continue to assist in determining the age structure of the Fort Peck Reservoir paddlefish stock. A phone survey should be conducted in 2010, using database of anglers who purchased tags to assess angler harvest of paddlefish. Telemetry study will be continued in 2010. Additionally, harvest rates should be closely monitored and the harvest cap will be implemented.

### **Fresno and Nelson Reservoir**

Standardized late-summer seining should continue to assess sport fish reproduction and forage fish abundance in Fresno and Nelson Reservoirs. Standardized sampling of adult sport fishes should be continued utilizing fall gill netting to gather recruitment information relating to walleye and other key sport and forage fish year-class strength and winter reservoir water levels. Walleye reproduction is still considered to be below optimum at Fresno and Nelson Reservoirs. Walleye fingerling should be stocked in Fresno and Nelson in 2010 to establish strong year-classes. Stocked fingerling should be marked with Oxytetracycline (OTC) so that recruitment of stocked walleye and natural reproduction can be determined.

### **Beaver Creek Reservoir**

Standardized late-summer seining should continue to assess sport fish reproduction and forage fish abundance at Beaver Creek Reservoir. Standardized sampling of adult sport fishes should be continued utilizing fall gill netting to gather recruitment information relating to sport and forage fish year-class strength and to monitor growth and survival of stocked walleye, rainbow trout, and forage availability. Spring and fall plants of walleye fingerlings and advanced fingerlings should be continued.

### **Hill, Blaine & Phillips Co. Ponds**

Sampling of adult sport fish populations should continue annually at Bailey Reservoir, Ester Reservoir, and Cow Creek Reservoir. All other ponds should be sampled every two to three years to assess adult fish populations, growth, and recruitment. In addition self-creel survey boxes will be maintained to assess the fishing pressure at these ponds. This information will allow us to tailor our management and stocking efforts to meet the needs of the public. Yellow perch should be collected from the Kremlin Water District Ponds and transferred to Cow Creek Reservoir and Ester in the spring of 2010. Northern pike will be stocked in Wild Horse. Bluegill will be trapped from Taint and Sagebrush and transferred to PR 018, Lark, Wapiti, and Karsten Coulee.



### **Daniels, Sheridan, and Valley Co. Ponds**

Sampling of adult sport fish populations should continue annually at Box Elder Reservoir. All other ponds should be sampled every two to three years to assess adult fish populations, growth, and recruitment.

### **REFERENCES**

- Berg, R. K. 1981. Fish populations of the wild and scenic Missouri River, Montana, Montana Department of Fish, Wildlife, and Parks. 242pp.
- Bowersox, B. J. 2004. An investigation of paddlefish, *Polyodon spathula*, and their prey in Fort Peck Reservoir, Montana. Master's Thesis, University of Idaho, Moscow, 95 p.
- Kozfkay, J. R. and D. L. Scarnecchia. 2002. Year-class strength and feeding ecology of age-0 and age-1 paddlefish (*Polyodon spathula*) in Fort Peck Lake, Montana, USA. Journal of Applied Ichthyology **18**: 601-607.
- Leslie, L. 2005. Statewide Fisheries Investigations, Northeast Montana warm water ecosystems investigations, survey and inventory of coldwater and warm water ecosystems. Montana Fish, Wildlife, and Parks. Project F-11-R-4. Helena. 20 pp.
- Leslie, L. 2006. Statewide Fisheries Investigations, Northeast Montana warm water ecosystems investigations, survey and inventory of coldwater and warm water ecosystems. Montana Fish, Wildlife, and Parks. Project F-11-R-4. Helena. 44 pp.
- McFarland, B. and R. Roche. 1987. Creel Census Program. Vers. 1. Computer Software. Montana Fish, Wildlife, & Park, 1987.
- McFarland, B. 2006. 2005 Statewide Angling Pressure Use Report. Montana Fish, Wildlife, & Parks, Helena, MT. Pp. 173.
- Neuhold, J. M. and H. L. Kuo. 1957. Creel Census Method. Utah State Department of Fish and Game. 36pp.
- Miller, S. E. 2005. Protocol for juvenile paddlefish visual counts and zooplankton sampling on Fort Peck Reservoir, Montana. Moscow, University of Idaho: 6.
- Miller, S. E. and D. E. Scarnecchia. 2006. Juvenile paddlefish visual counts and zooplankton sampling on Fort Peck Reservoir, Montana: Report for 2005. University of Idaho, Moscow. 5p.
- Miller, S. E., D. E. Scarnecchia, and L. Leslie. 2006. Spring migrations of adult paddlefish in the Missouri River above Fort Peck Reservoir: Progress report for 2006. University of Idaho, Moscow. 27p.
- Riggs, V. 2005. Montana Paddlefish Telephone Creel Survey 2003 and 2004. Miles City, Montana Fish, Wildlife & Parks: 20.

US Congress. 1975a. Hearings on Senate Bill 1506, a bill to amend the wild and scenic rivers act, part 2- Missouri River, Montana. US Government Printing Office, Washington, D. C. 444pp.

US Congress. 1975b. Designating a segment of the Missouri River in the state of Montana as a component of the national wild and scenic river system. Senate Report No. 94-502. 16pp.

USFWS (U.S. Fish and Wildlife Service). 1999. Endangered and threatened wildlife. United States Code of Federal Regulations, Title 50, Sections 17.11 and 17.12.

United States Geological Survey. 2008 Water resources data for Missouri River near Virgelle MT. Retrieved: January 5, 2008, from [http://waterdata.usgs.gov/ mt/](http://waterdata.usgs.gov/mt/).

### **Water Codes of Waters Referred To**

154535	Bailey Reservoir	162500	Missouri River Sec. 05
164375	Bar Island Reservoir	162520	Missouri River Sec. 06
168616	Bell Ridge Reservoir	156480	Nelson Reservoir
154570	Beaver Creek Reservoir	167399	Paleface Reservoir
154590	Big Reservoir	156598	Paulo Reservoir
154700	Boucher Reservoir	157040	PR 018
160630	Burke Reservoir	167780	PR 020
164575	Butch Reservoir	157600	PR 054
154761	Cole Ponds	158860	Reser Reservoir
154772	Compton Reservoir	168047	Sagebrush Reservoir
164789	Cow Creek Reservoir	154731	South Casady Reservoir
155120	Ester Lake	168445	Sundance Reservoir
155200	Frenchman Reservoir	168475	Taint Reservoir
165140	Fort Peck Reservoir	155340	Upper Base Pond
155240	Fresno Reservoir	159505	Valley Reservoir
165219	Gazob Reservoir	168706	Wapiti Reservoir
165385	Gullwing Reservoir	1500AS	Wards Reservoir
166155	Karsten Coulee Reservoir	168815	Whiteface Reservoir
166495	Lark Reservoir	168860	Whitetail Reservoir
155340	Lower Base Pond		
166921	McChesney Reservoir		

Key words:

Paddlefish, harvest, walleye, Lake Superior whitefish, northern pike, black crappie, yellow perch, goldeye, channel catfish, sauger, shovelnose sturgeon, pallid sturgeon, burbot, smallmouth buffalo, largemouth buffalo, age, water levels, creel census, population estimates, recruitment, tiger musky, stocking, Cisco, smelt

Prepared by: Cody Nagel

Date: March 23, 2010